

AD-A068 442

ARMY INST OF SURGICAL RESEARCH FORT SAM HOUSTON TEX

F/G 6/5

USE OF ANTISERA TO HUMAN PLASMA PROTEINS TO MEASURE SPECIFIC MO--ETC(U)

APR 79 M C POWANDA, R F BERENDT

UNCLASSIFIED

NL

1 OF 1  
ADA  
068442



END  
DATE  
FILMED

6-79

DOC



**LEVEL**

2 B.S.

6

Use of Antisera to Human Plasma Proteins to Measure Specific Monkey Plasma

Proteins

AD A 068442

10

Michael C./Powanda and Richard F./Berendt

U. S. Army Institute of Surgical Research

Brooke Army Medical Center

Fort Sam Houston, TX 78234

and

U. S. Army Medical Research Institute of Infectious Diseases

Fort Detrick, Frederick, MD 21701

Running Title: Quantitation of Monkey Plasma Proteins

DDC  
MAY 9 1979  
C

12 13p

DDC FILE COPY

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Care.

The views of the authors do not purport to reflect the positions of the Department of the Army or the Department of Defense.

11

17 Apr 1979

Cleared for public release; distribution unlimited.

404 885 LB  
79 05 09 044

USE OF ANTISERA TO HUMAN PLASMA PROTEINS  
TO MEASURE SPECIFIC MONKEY PLASMA PROTEINS<sup>1, 2</sup>

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DOC	B..ff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	and/or SPECIAL
ff	

QUANTITATION OF MONKEY PLASMA PROTEINS

79 05 09 044

Michael C. Powanda, Ph.D.

and

Richard F. Berendt, Ph.D.

1/ From the United States Army Institute of Surgical Research  
Brooke Army Medical Center, Fort Sam Houston, TX 78234, and  
the United States Army Medical Research Institute of Infectious  
Diseases, Fort Detrick, Frederick, Maryland 21701

2. We gratefully acknowledge the technical assistance of John P. Fowler.
3. In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council. The facilities are fully accredited by the American Association for Accreditation of Laboratory Animal Care. The views of the authors do not purport to reflect the positions of the Department of the Army or the Department of Defense.
4. Reprint requests to: Michael C. Powanda, Department of Biochemistry, USA Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, TX 78234.

**Key Words**

**Monkeys**

**Plasma proteins**

**Radial immunodiffusion**

SUMMARY

Antibodies against specific human plasma proteins were used to estimate the concentration of ceruloplasmin, haptoglobin, alpha<sub>1</sub> acid glycoprotein, transferrin, and alpha<sub>1</sub> antitrypsin in the plasma of rhesus, cynomolgus and squirrel monkeys.

alpha(1)

Plasma acute-phase globulins increase in concentration during injury, infection and various inflammatory diseases (1,2). In fact, certain of these proteins appear to be very sensitive indicators of the extent and persistence of injury/inflammation (3). Other plasma proteins have been used as monitors of nutritional status (4). It has been hypothesized that many of these plasma proteins interact with other aspects of host metabolism, physiology and immunology to ameliorate the consequences of disease and enhance healing (5). Monkeys, being closely related to man, have been used to investigate a wide variety of diseases and nutritional disorders (6,7).

A simple method for quantitating specific plasma proteins would allow assessment of the effects of various diseases and nutritional regimens on individual plasma proteins. To this end we tested radial immunodiffusion plates (5) designed for determination of human plasma proteins for their ability to detect and measure proteins in monkey plasma. Plasma was obtained by venipuncture of three monkeys of each of three selected species. Table 1 shows that all three species of monkeys possessed plasma proteins antigenically related to those of man. Alpha<sub>1</sub> acid glycoprotein was present in amounts too low for accurate determination. No reaction was detectable when rat plasma was used instead of monkey. Whether the monkey plasma proteins which react with antisera to human plasma proteins are functionally analogous to those of man remains to be determined, but certain of these proteins do seem to respond during infection in a manner similar to their presumed

5. Prepared by Behring Diagnostics, American Hoechst Corp.,  
Somerville, NJ.

human counterparts (7). There appears to be a fair degree of monkey-to-monkey variation so one might have to use each animal as its own control, as well as using pooled normal, i.e., healthy monkey plasma, for an additional reference.

## REFERENCES

1. Werner M: Serum protein changes during the acute phase reaction. Clin Chim Acta 25:299-305, 1969
2. Bostian K A, Blackburn B S, Wannemacher R W, Jr, et al: Sequential changes in the concentration of specific serum proteins during typhoid fever infection in man. J Lab Clin Med 87:577-585, 1976
3. Fischer C L, Gill C, Forrester M G, et al: Quantitation of "acute-phase proteins" postoperatively. Value in detection and monitoring of complications. Am J Clin Pathol 66:840-846, 1976
4. McFarlane H: Acute-phase proteins in malnutrition. In Malnutrition and the Immune Response, p 403-405, Suskind R M, Ed, Raven Press, New York, 1977
5. Powanda M C: Changes in body balances of nitrogen and other key nutrients: description and underlying mechanisms. Am J Clin Nutr 30:1254-1268, 1977
6. Bustad L K, Hegreberg G A, Padgett G A: Naturally Occurring Animal Models of Human Disease: A Bibliography. Institute of Laboratory Animal Resources, National Academy of Sciences, Washington, DC, 1976
7. Berendt R F, Knutsen G L, Powanda M C: Nonhuman primate model for the study of respiratory Klebsiella pneumoniae infection. Infect Immun 22:275-281, 1978

TABLE 1

Cross-reactivity of monkey plasma proteins with human antisera

Antiserum <sup>a</sup> Against:	mg/dl (Coefficient of variation <sup>b</sup> )		
	Rhesus ( <i>Macaca mulatta</i> )	Cynomolgus ( <i>Macaca fascicularis</i> )	Squirrel ( <i>Saimiri sciureus</i> )
Ceruloplasmin	67.1 (1.8)	47.7 (8.9)	56.4 (3.9)
Haptoglobin	222.7 (7.9)	317.6 (4.7)	448.2 (5.0)
Alpha <sub>1</sub> - antitrypsin	1021.3 (3.7)	904.0 (4.0)	724.0 (7.9)
Transferrin	499.7 (5.6)	506.0 (6.2)	626.7 (5.0)

<sup>a</sup> Ceruloplasmin and haptoglobin based upon undiluted plasma; alpha<sub>1</sub> - antitrypsin and transferrin based upon a 1:10 dilution of plasma.

<sup>b</sup> Coefficient of variation:  $CV(\%) = \frac{s}{\bar{x}} \times 100.$