

AD-A068 943

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 20/4

GALERKIN METHODS FOR MISCIBLE DISPLACEMENT PROBLEMS IN POROUS M--ETC(U)

MAR 79 R E EWING, M F WHEELER

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-1932

NL

| OF |

AD
A068943



END

DATE
FILMED

7-79

DDC

LEVEL

2

AD A068943

MRC Technical Summary Report #1932

GALERKIN METHODS FOR MISCIBLE
DISPLACEMENT PROBLEMS IN POROUS MEDIA

Richard E. Ewing and Mary Fanett Wheeler

See 1473 in back.

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

March 1979

(Received December 4, 1978)

DDC
RECEIVED
MAY 23 1979
C

Approved for public release
Distribution unlimited

Sponsored by
U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

National Science Foundation
Washington, D. C. 20550

DDC FILE COPY

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

GALERKIN METHODS FOR MISCIBLE DISPLACEMENT

PROBLEMS IN POROUS MEDIA

Richard E. Ewing¹ and Mary Fanett Wheeler²

Technical Summary Report #1932
March 1979

ABSTRACT

A priori error estimates for Galerkin methods for numerical approximation of the coupled quasilinear system for $c = c(x,t)$ and $p = p(x,t)$ given by

$$\nabla \cdot [a(x,c) \{ \nabla p - \gamma(x,c) \nabla z \}] = 0 ,$$

$$\nabla \cdot [b(x,c, \nabla p) \nabla c] - u(x,c, \nabla p) \cdot \nabla c = \phi(x) \frac{\partial c}{\partial t} ,$$

for $x \in \Omega$, $t \in (0,T]$, and appropriate Neumann boundary and initial conditions are considered. Equations of this type arise in models for the miscible displacement of one incompressible fluid by another in a porous medium.

Estimates for both continuous time and fully-discrete time Galerkin methods are presented.

AMS (MOS) Subject Classifications: 65M15, 65N15, 65N30, 76.35

Key Words: Galerkin methods, Error estimates, Fluid flow, Numerical analysis
Work Unit Number 7 (Numerical Analysis)

¹The Ohio State University, Columbus, Ohio 43210.

²Rice University, Houston, Texas 77001.

SIGNIFICANCE AND EXPLANATION

Equations of the type stated in the Abstract arise, for example, in models for the miscible displacement of one incompressible fluid by another in a porous medium as in chemical flooding of oil wells to recover more of the oil. The variable $c = c(x,t)$ corresponds to the concentration of the fluid used for the flooding process while the variable $p = p(x,t)$ corresponds to the pressure at the location x in the medium at time t .

This paper gives error analysis and error estimates for certain numerical methods for approximate solution of the coupled quasilinear system of parabolic partial differential equations in the Abstract. Both continuous-time and fully-discrete time methods are presented and analyzed.

ACCESSION for	
NTIS	White Section ☒
DDC	Buff Section ☐
UNANNOUNCED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
04	SPECIAL
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

GALERKIN METHODS FOR MISCIBLE DISPLACEMENT PROBLEMS IN POROUS MEDIA

Richard E. Ewing¹ and Mary Fanett Wheeler²

1. Introduction

We consider the numerical approximation by Galerkin methods to a problem arising in the miscible displacement of one incompressible fluid by another in a porous medium. A set of equations [6] and [9] modeling the pressure $p(x,y,t)$ and the concentration $c(x,y,t)$ is given by

$$(1.1) \quad \frac{\partial}{\partial x} \left(\frac{k_x}{\mu(c)} \left(\frac{\partial p}{\partial x} - \gamma(c) \frac{\partial z}{\partial x} \right) \right) + \frac{\partial}{\partial y} \left(\frac{k_y}{\mu(c)} \left(\frac{\partial p}{\partial y} - \gamma(c) \frac{\partial z}{\partial y} \right) \right) \equiv - \frac{\partial}{\partial x} u_x - \frac{\partial}{\partial y} u_y = q(x,y)$$

and

$$(1.2) \quad \begin{aligned} & \frac{\partial}{\partial x} (\alpha(\phi, D, c, \nabla p) \frac{\partial c}{\partial x} - u_x c) + \frac{\partial}{\partial y} (\delta(\phi, D, c, \nabla p) \frac{\partial c}{\partial y} - u_y c) \\ & + \frac{\partial}{\partial y} (\beta(\phi, D, c, \nabla p) \frac{\partial c}{\partial x}) + \frac{\partial}{\partial x} (\beta(\phi, D, c, \nabla p) \frac{\partial c}{\partial y}) \\ & = \phi(x,y) \frac{\partial c}{\partial t} + q(x,y) \hat{c}(x,y,t), \end{aligned}$$

for $(x,y) \in \Omega$, $t \in J = (0,T]$, where the reservoir Ω is a bounded domain with boundary $\partial\Omega$. Here $k_x = k_x(x,y)$ and $k_y = k_y(x,y)$ are the permeabilities in the x and y directions respectively, $\mu = \mu(c)$ is the local viscosity of the fluid, $\gamma = \gamma(c)$ is its density, $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ are the components of the dip angle, and q , the imposed total flow rate, is a linear combination of a finite number of Dirac measures, (i.e. sources and sinks). The function $\phi = \phi(x,y)$ is the porosity and $D = D(x,y)$ is the molecular diffusion level. The components of the dispersion coefficient, α , δ and β , are functions of porosity, diffusion, concentration, and the gradient of pressure. It is not uncommon in petroleum reservoir engineering for one to assume that $\beta = 0$ and $\alpha = \alpha(D)$ and $\delta = \delta(D)$ (see [9]). The function $\hat{c}$ is equal to the specified concentration at injection wells (sources) and to $c(x,y,t)$ for production wells (sinks). No flow

¹The Ohio State University, Columbus, Ohio 43210.

²Rice University, Houston, Texas 77001.

conditions are assumed on the boundary of the reservoir and an initial concentration is given throughout the reservoir. For a general discussion of the physics of miscible displacement problems in reservoir engineering the reader is referred to Peaceman [7].

In the Galerkin procedure we shall use different subspaces of $H^1(\Omega)$ to approximate pressure and concentration. We note that, if $c(x,y,t) \equiv 1$, then (1.1) and (1.2) are identical. In this case we are forced to use the same approximating subspaces for pressure and concentration. We are thus led to replacing (1.2) by the non-divergence form

$$(1.3) \quad \begin{aligned} & \frac{\partial}{\partial x} \left(\alpha \frac{\partial c}{\partial x} \right) + \frac{\partial}{\partial y} \left(\delta \frac{\partial c}{\partial y} \right) + \frac{\partial}{\partial y} \left(\beta \frac{\partial c}{\partial x} \right) + \frac{\partial}{\partial x} \left(\beta \frac{\partial c}{\partial y} \right) - u_x \frac{\partial c}{\partial x} - u_y \frac{\partial c}{\partial y} \\ & = \phi \frac{\partial c}{\partial t} + q(x,y) (\bar{c} - c), \end{aligned}$$

which is obtained by multiplying (1.1) by c and subtracting the result from (1.2).

Settari, Price, and Dupont [9] presented, without analysis, a finite element method for solving (1.1)-(1.3) with $\delta \equiv 0$. For simplicity, we shall treat the coupled quasilinear system with dependent variables $c = c(x,t)$ and $p = p(x,t)$ given by

$$(1.4) \quad \nabla \cdot [a(x,c) (\nabla p - \gamma(x,c) \nabla z)] = 0,$$

$$(1.5) \quad \nabla \cdot [b(x,c, \nabla p) \nabla c] - u(x,c, \nabla p) \cdot \nabla c = \phi(x) \frac{\partial c}{\partial t},$$

for $x \in \Omega$, $t \in J \equiv (0,T)$, where Ω is a bounded domain in $\mathbb{R}^d$, $d = 2$, with boundary $\partial\Omega$ and $u(x,c, \nabla p) = -a(x,c) (\nabla p - \gamma(x,c) \nabla z)$ is a vector in $\mathbb{R}^2$. The analysis for (1.4)-(1.5) easily extends to the case (1.1) and (1.3) where mixed $x-y$ derivatives are present. We assume that the following boundary and initial conditions hold:

$$(1.6) \quad a(x,c) \left\{ \frac{\partial p}{\partial \nu} - \gamma(x,c) \frac{\partial z}{\partial \nu} \right\} = 0, \quad x \in \partial\Omega, \quad t \in J,$$

$$(1.7) \quad b(x,c, \nabla p) \frac{\partial c}{\partial \nu} = 0, \quad x \in \partial\Omega, \quad t \in J,$$

$$(1.8) \quad c(x,0) = c_0(x), \quad x \in \Omega,$$

where $\frac{\partial f}{\partial \nu}$ is the normal derivative of f on the boundary of Ω . We note that (1.4)-(1.8) will define $p(x,t)$ only to within an arbitrary constant. We shall

normalize p by the condition that

$$(1.9) \quad \frac{1}{|\Omega|} \int_{\Omega} p(x,t) dx = 1, \quad t \in J,$$

where $|\Omega|$ is the measure of Ω . We note that for (1.4) and (1.5), we have chosen $q \equiv 0$ in (1.1) and (1.3). If q is assumed to be smooth, which corresponds to smoothly distributed sources and sinks, the analysis follows with few changes.

The paper consists of three additional sections. In Section 2, the continuous time and discrete time Galerkin procedures are formulated. Assumptions on the coefficients corresponding to the physical problem and smoothness assumptions on the solution are given. Methods are presented which allow the approximations for concentration and pressure to lie in different subspaces of $H^1(\Omega)$. In Section 3, a-priori error estimates for the continuous time approximations are given. Optimal L^2 rates of convergence are established for the case where the dispersion coefficient depends only upon molecular diffusion using subspaces of the same order of approximation. In the case that the dispersion coefficient also depends upon the Darcy velocity, we obtain an optimal L^2 rate using a subspace of one higher order for the pressure than for the concentration. In Section 4, these results are extended to the discrete time case.

2. The Finite Element Methods and a Précis of the Results

Let $(u, v) = \int_{\Omega} uv \, dx$ and $\|u\|^2 = (u, u)$. Let $W_s^k(\Omega)$ be the Sobolev space on Ω with norm

$$\|\phi\|_{W_s^k} = \left(\sum_{|\alpha| \leq k} \left\| \frac{\partial^\alpha \phi}{\partial x^\alpha} \right\|_{L^s(\Omega)}^s \right)^{\frac{1}{s}},$$

with the usual modification for $s = \infty$. When $s = 2$, let $\|\phi\|_{W_2^k} = \|\phi\|_{H^k} \equiv \|\phi\|_k$. If $\nabla f = (f_1, f_2)$ write $\|\nabla f\|_{W_s^k}$ in place of $(\|f_1\|_{W_s^k}^s + \|f_2\|_{W_s^k}^s)^{\frac{1}{s}}$. Also $H^s(\partial\Omega)$ will denote the usual Sobolev space on $\partial\Omega$.

Let $\{M_h\}$ be a family of finite-dimensional subspaces of $H^1(\Omega)$ with the following property:

For $p = 2$ or $p = \infty$, there exist an integer $r \geq 2$ and a constant K_0

such that, for $1 \leq q \leq r$ and $\phi \in W_p^q(\Omega)$:

$$(2.1) \quad \inf_{\chi \in M_h} (\|\phi - \chi\|_{W_p^0} + h \|\phi - \chi\|_{W_p^1}) \leq K_0 \|\phi\|_{W_p^q} h^q.$$

Similarly, we define a family of finite-dimensional subspaces of $H^1(\Omega)$ called $\{N_h\}$ which satisfies the same property as $\{M_h\}$ with r replaced by s . We also assume that the families $\{M_h\}$ and $\{N_h\}$ satisfy the following so-called "inverse hypotheses":

$$(2.2) \quad \begin{aligned} \text{a) } \|\phi\|_{L^\infty(\Omega)} &\leq K_0 h^{-\frac{d}{2}} \|\phi\| = K_0 h^{-1} \|\phi\|, \quad \text{and} \\ \text{b) } \|\nabla \phi\|_{L^\infty(\Omega)} &\leq K_0 h^{-1} \|\nabla \phi\|. \end{aligned}$$

Restrict Ω as follows (with S denoting the collection of restrictions):

(S) 1) Ω is H^2 -regular, i.e., if

$$-\Delta v + \theta v = \zeta, \quad x \in \Omega, \quad \theta = 0 \text{ or } 1,$$

$$\frac{\partial v}{\partial \nu} = \eta, \quad x \in \partial\Omega,$$

$$\text{and } (\zeta, 1) + \int_{\partial\Omega} \eta ds = 0 \text{ if } \theta = 0,$$

then

$$\|v\|_2 \leq K(\Omega) (\|\zeta\| + \|\eta\|_{H^{1/2}(\partial\Omega)});$$

2) $\partial\Omega$ is Lipschitz.

Let

$$b \equiv b(x, c, \nabla p) \equiv b(x, c, p_x, p_y)$$

and

$$u \equiv u(x, c, \nabla p) = (u_1(x, c, p_x), u_2(x, c, p_y)) .$$

For some $\hat{\varepsilon} > 0$ restrict the variable q_1 to lie between

$$-\hat{\varepsilon} \leq q_1 \leq 1 + \hat{\varepsilon} .$$

Assume the following regularity for a , γ , b , u , and ϕ :

(Q): 1) There exist constants a_* , b_* , ϕ_* and K_1 such that

$$(2.3) \quad \begin{aligned} a) & \quad 0 < a_* \leq a(x, q_1) \leq K_1 , \\ b) & \quad |\gamma(x, q_1)| \leq K_1 , \\ c) & \quad 0 < \phi_* \leq \phi(x) \leq K_1 , \\ d) & \quad |\nabla z(x)| \leq K_1 , \\ e) & \quad 0 < b_* \leq b(x, \tilde{q}, q_2, q_3), \quad \tilde{q}, q_2, q_3 \in \mathbb{R} , \\ f) & \quad |u_i(x, q_1, q_4)| \leq K_1 (1 + |q_4|), \quad i = 1, 2, q_4 \in \mathbb{R} , \\ g) & \quad |b(x, c, \nabla p)| \leq K_1 . \end{aligned}$$

2) There exists a constant M such that for (c, p) the solution to (1.4)-(1.8)

and for $(q_2, q_3) \in \mathbb{R}^2$, $i = 1, 2$,

$$(2.4) \quad \begin{aligned} & \left| \frac{\partial a}{\partial c}(x, q_1) \right| + \left| \frac{\partial \gamma}{\partial c}(x, q_1) \right| + \left| \frac{\partial^2 b}{\partial c^2}(x, c, \nabla p) \right| + \left| \frac{\partial u_i}{\partial q_1}(x, q_1, \nabla p) \right| \\ & + \left| \frac{\partial u_i}{\partial q_2}(x, q_1, q_2) \right| + \left| \frac{\partial^2 u_i}{\partial x_i \partial q_2}(x, c) \right| + \left| \frac{\partial^2 u_i}{\partial c^2}(x, c, \nabla p) \right| + \left| \frac{\partial^2 u_i}{\partial c \partial q_2}(x, c) \right| \\ & + \left| \frac{\partial b}{\partial q_1}(x, q_1, p_x, p_y) \right| + \left| \frac{\partial b}{\partial q_2}(x, q_1, q_2, q_3) \right| + \left| \frac{\partial b}{\partial q_3}(x, q_1, q_2, q_3) \right| \leq M . \end{aligned}$$

The assumptions made on the coefficients in (Q) are physically motivated. In the general miscible displacement formulation (1.1)-(1.3)

$$\alpha = \phi \left\{ D + \alpha_\ell \frac{u_x^2}{|||u|||} + \alpha_t \frac{u_y^2}{|||u|||} \right\} ,$$

$$\beta = \phi \left\{ D + (\alpha_\ell - \alpha_t) \frac{u_x u_y}{|||u|||} \right\} , \quad \text{and}$$

$$\delta = \phi \left\{ D + a_t \frac{u_y^2}{|||u|||} + a_x \frac{u_x^2}{|||u|||} \right\},$$

where $|||u||| = \sqrt{u_x^2 + u_y^2}$ and a_t and a_x are nonnegative constants (see [6] and [9]). One can show that a , δ , and δ satisfy the conditions assumed for b , that $k_x/\mu(c)$ and $k_y/\mu(c)$ satisfy those for a , and that u_x and u_y satisfy assumptions made on the u_1 .

Let

$$||\phi||_{L^q((a,b),X)} = |||\phi(\cdot, t)|||_X |||_{L^q(a,b)}, \quad 1 \leq q \leq \infty.$$

Let (p, c) , the solution of (1.4)-(1.8), satisfy the following regularity assumptions:

$$(R): \quad a) \quad ||c||_{L^\infty(J; H^r)} + \left\| \frac{\partial c}{\partial t} \right\|_{L^2(J; H^r)} + ||p||_{L^\infty(J; H^s)} \leq K_2,$$

$$(2.5) \quad b) \quad ||c||_{L^\infty(J; H^{2+\epsilon})} + \left\| \frac{\partial c}{\partial t} \right\|_{L^\infty(J; W_3^1)} + \left\| \frac{\partial^2 c}{\partial t^2} \right\|_{L^2(J; L^2)} + \left\| \frac{\partial p}{\partial t} \right\|_{L^\infty(J; H^1)} \leq K_2,$$

for some $\epsilon > 0$.

The analysis proceeds, following Wheeler [10], using a pair of auxiliary elliptic problems. Let $\tilde{p} \in N_h$ be the elliptic projection of p into N_h defined by

$$(2.6) \quad (a(\cdot, c(\cdot, t)) \nabla \tilde{p}, \nabla v) = (a(\cdot, c(\cdot, t)) \nabla p, \nabla v), \quad v \in N_h,$$

for each $t \in J$, where

$$(2.7) \quad \frac{1}{|\Omega|} \int_{\Omega} \tilde{p}(x, t) dx = 1, \quad \text{for each } t \in J,$$

and where (p, c) is the solution of (1.4)-(1.8). The restrictions (S) imply the following result.

Lemma 2.1. There exists $K_3 = K_3(\Omega, \varepsilon, K_0, K_1)$ such that

$$(2.8) \quad ||\tilde{p} - p|| + h ||\nabla(\tilde{p} - p)|| \leq K_3 h^s ||p||_s.$$

Let $\lambda > 0$ be chosen sufficiently large that the bilinear form

$$B(\psi, \chi) = (b(c, \nabla p) \nabla \psi, \nabla \chi) + (u(c, \nabla p) \cdot \nabla \psi, \chi) + \lambda(\psi, \chi)$$

is coercive over $H^1(\Omega)$. Let $\tilde{c} \in M_h$ be the elliptic projection of c into M_h , defined by

$$(2.9) \quad B(\tilde{c}, w) = B(c, w) = -(\phi \frac{\partial c}{\partial t}, w) + \lambda(c, w), \quad w \in M_h,$$

for each $t \in J$. Then, as in [10] and [3], we can obtain the following lemma.

Lemma 2.2. There exists $K_4 = K_4(\Omega, b_4, \lambda, K_0, K_1, K_2, M)$ such that

$$\begin{aligned} \text{a) } & \|\tilde{c} - c\|_{L^2(J; L^2)} + \left\| \frac{\partial(\tilde{c} - c)}{\partial t} \right\|_{L^2(J; L^2)} + h \|\tilde{c} - c\|_{L^2(J; H^1)} \\ (2.10) \quad & \leq K_4 h^r \left\{ \|c\|_{L^2(J; H^r)} + \left\| \frac{\partial c}{\partial t} \right\|_{L^2(J; H^r)} \right\}, \\ \text{b) } & \|\tilde{c} - c\|_{L^\infty(J; L^2)} \leq K_4 h^r \|c\|_{L^\infty(J; H^r)}. \end{aligned}$$

Assume that there exists a constant K_5 such that

$$(2.11) \quad \|\nabla \tilde{p}\|_{L^\infty(J; L^\infty)} + \|\nabla \tilde{c}\|_{L^\infty(J; L^\infty)} \leq K_5.$$

See [2] and [10] for some sufficient conditions for these assumptions.

We first consider continuous time approximations of (p, c) . Denote the approximation of p by $P : [0, T] \rightarrow N_h$ and the approximation of c by $C : [0, T] \rightarrow M_h$, where (P, C) is defined by the relations: (we suppress the dependence of the coefficients on x)

$$(2.12) \quad (a(C) \nabla P, \nabla v) = (a(C) \gamma(C) \nabla z, \nabla v), \quad v \in N_h,$$

and

$$(2.13) \quad (b(C, \nabla P) \nabla C, \nabla w) + (u(C, \nabla P) \cdot \nabla C, w) + \left(\frac{\partial C}{\partial t}, w \right) = 0, \quad w \in M_h,$$

with

$$(2.14) \quad C(x, 0) = \tilde{c}(x, 0),$$

where $\tilde{c}$ is the elliptic projection of c defined in (2.9). In Section 3, we shall obtain a priori estimates for $c - C$ and for $\nabla(p - P)$.

Next, we define a fully discrete, $O(\Delta t)$ -correct method for approximating (p, c) based on backward differencing in time. Let $\Delta t > 0$, $N = T/\Delta t \in \mathbb{Z}$, and $t^0 = 0$, $t \in \mathbb{R}$. Also, let $\psi^n = \psi^n(x) = \psi(x, t^n)$ and $d_t \psi^n = (\psi^{n+1} - \psi^n)/\Delta t$. Denote the approximation of p by $W : \{0 = t^0, t^1, \dots, t^N = T\} \rightarrow N_h$ and the approximation of c

by $Z : \{t^0, t^1, \dots, t^N\} \rightarrow M_h$. Assuming that W^n and Z^n are known, we determine W^{n+1} and Z^{n+1} as follows:

$$(2.15) \quad (\phi_d_t Z^n, \psi) + (b(Z^n, VW^n) \nabla Z^{n+1}, \nabla \psi) + (u(Z^n, VW^n) \cdot \nabla Z^{n+1}, \psi) = 0, \quad \psi \in M_h,$$

and

$$(2.16) \quad (a(Z^{n+1}) VW^{n+1}, \nabla y) = (a(Z^{n+1}) \gamma(Z^{n+1}) \nabla Z, \nabla y), \quad y \in N_h,$$

where $j = 0$ or 1 . We note that the coefficient matrix arising from the algebraic system (2.15) with $j = 0$ is symmetric. However, in many problems the transport term is large and it may be numerically advantageous to use (2.15) with $j = 1$ even though the coefficient matrix is no longer symmetric. Both discretizations are $O(\Delta t)$ in time. In (2.16) no explicit time discretization error is made. A priori error estimates will be presented for $c^n - Z^n$ and $V(P^n - W^n)$ in Section 4.

Let $f^{n+1/2} \equiv (f^{n+1} + f^n)/2$. If (2.15) is replaced by

$$(2.17) \quad (\phi_d_t \tilde{Z}^n, \psi) + (b(E\tilde{Z}^n, VEW^n) \nabla \tilde{Z}^{n+\frac{1}{2}}, \nabla \psi) + (u(E\tilde{Z}^n, VEW^n) \cdot \nabla \tilde{Z}^{n+\frac{1}{2}}, \psi) = 0, \quad \psi \in M_h,$$

where $E\tilde{Z}^n = \frac{3}{2} \tilde{Z}^n - \frac{1}{2} \tilde{Z}^{n-1}$, an analysis similar to that given in Section 4 (see [8], [10], [2] and [5]) will yield a time discretization error of size $O((\Delta t)^2)$.

When estimating the actual physical quantities, we know that the pressure is much smoother in time than the concentration. Thus, in practice, one should not use (2.16) to redetermine VW^{n+1} at each time step. Instead one value of VW^{n+1} should be used in the coefficients of (2.15) for several time steps before a new pressure is approximated. One way to view this procedure is to have two different time steps for the different equations. An analysis of this method will follow in a more general paper.

3. A Priori Error Estimates for the Continuous Time Approximations

In this section, we develop a priori bounds for the errors $(c - C)$ and $\nabla(p - P)$. We shall see that if b depends on ∇p then for the best rates of convergence one should use subspaces of piecewise polynomials of different degrees for $\{M_h\}$ and $\{N_h\}$. We first obtain an estimate for $P - \tilde{p}$ where P is defined by (2.12)-(2.14) and $\tilde{p}$ is defined by (2.6)-(2.7).

We assume that for some $\epsilon > 0$, $\epsilon \leq \hat{\epsilon}$, (see the statement preceding Q),

$$(3.0) \quad -\epsilon \leq C \leq 1 + \epsilon.$$

We shall prove that

$$\|c - C\| = O(h^r + h^{s-1}),$$

which yields

$$(3.1) \quad \|c - C\|_{L^\infty} \leq K(h^{r-1} + h^{s-2}).$$

Since $0 \leq c \leq 1$, one can deduce from (3.1) with the restriction $r \geq 2$ and $s \geq 3$ that (3.0) holds for small h .

Lemma 3.1. There exists a positive constant $K_6 = K_6(a_*, K_1, K_5, M)$ such that

$$\|\nabla(P - \tilde{p})\| \leq K_6 \|c - C\|, \quad t \in J.$$

Proof. Subtract (2.6) from (2.12) and use (1.4) and (1.6) to obtain

$$(3.2) \quad (a(C)\nabla(P - \tilde{p}), \nabla v) = ([a(c) - a(C)]\nabla \tilde{p}, \nabla v) + ([a(C)\gamma(C) - a(c)\gamma(c)]\nabla z, \nabla v), \quad v \in N_h.$$

If $v = P - \tilde{p} \in N_h$, then

$$(3.3) \quad \begin{aligned} a_* \|\nabla(P - \tilde{p})\|^2 &\leq |(a(C)\nabla(P - \tilde{p}), \nabla(P - \tilde{p}))| \leq K_7 \|c - C\| \|\nabla(P - \tilde{p})\| \\ &\leq \frac{a_*}{2} \|\nabla(P - \tilde{p})\|^2 + \frac{K_7^2}{2a_*} \|c - C\|^2, \quad t \in J, \end{aligned}$$

where K_7 depends upon uniform bounds for $|\frac{\partial a}{\partial c}|$, $|\frac{\partial \gamma}{\partial c}|$, $|\nabla z|$, and $\|\nabla \tilde{p}\|_{L^\infty(J; L^\infty)}$. The result of the lemma follows directly from (3.3).

We next compare C and $\tilde{c}$, defined in (2.12)-(2.14) and (2.9), respectively. We shall make an additional assumption on P defined in (2.12)-(2.14). Assume that there exists a positive constant K^* such that

$$(3.4) \quad \|\nabla P\|_{L^\infty(J; L^\infty)} \leq K^*.$$

Without loss of generality, assume $K^* \geq 2K_5$.

Theorem 3.2. There exists $K_8 = K_8(\lambda, b_*, \phi_*, M; K_i, i \leq 6; K^*)$ such that

$$(3.5) \quad \|C - \tilde{C}\|_{L^\infty(J; L^2)} + \|\nabla(C - \tilde{C})\|_{L^2(J; L^2)} \leq K_8(h^r + h^{s-1}).$$

If s and r from Lemmas 2.1 and 2.2 satisfy $s \geq 3$ and $r \geq 2$ and h is taken sufficiently small, then

$$(3.6) \quad \|\nabla P\|_{L^\infty(J; L^\infty)} \leq 2K_5 \leq K^*$$

and a K_8 above can be chosen which is independent of K^* .

Proof. Subtract (2.9) from (2.13) to obtain

$$(3.7) \quad \begin{aligned} & (b(C, \nabla P) \nabla(C - \tilde{C}), \nabla w) + \left(\phi \left(\frac{\partial C}{\partial t} - \frac{\partial \tilde{C}}{\partial t} \right), w \right) = ([b(C, \nabla P) - b(C, \nabla \tilde{P})] \nabla \tilde{C}, \nabla w) \\ & + \left(\phi \left(\frac{\partial C}{\partial t} - \frac{\partial \tilde{C}}{\partial t} \right), w \right) - (u(C, \nabla P) \cdot \nabla(C - \tilde{C}), w) \\ & + ([u(C, \nabla P) - u(C, \nabla \tilde{P})] \cdot \nabla \tilde{C}, w) + \lambda(\tilde{C} - C, w), \quad w \in M_h. \end{aligned}$$

Next, letting $w = C - \tilde{C} \in M_h$, integrating the left side of (3.7) from 0 to t and using (2.14), we obtain

$$(3.8) \quad \begin{aligned} & \int_0^t \left(\phi \frac{\partial}{\partial \tau} (C - \tilde{C}), (C - \tilde{C}) \right) d\tau + \int_0^t (b(C, \nabla P) \nabla(C - \tilde{C}), \nabla(C - \tilde{C})) d\tau \\ & \geq \frac{\phi_*}{2} \| (C - \tilde{C})(t) \|^2 + b_* \int_0^t \| \nabla(C - \tilde{C})(\tau) \|^2 d\tau. \end{aligned}$$

We next use (2.8), Lemma 2.2, Lemma 3.1 and Hölder's inequality to see that

$$(3.9) \quad \begin{aligned} & \left| \int_0^t ([b(C, \nabla P) - b(C, \nabla \tilde{P})] \nabla \tilde{C}, \nabla(C - \tilde{C})) d\tau \right| \\ & \leq K_9 \|\nabla \tilde{C}\|_{L^\infty(J; L^\infty)} \int_0^t (\|C - \tilde{C}\| + \|\nabla(P - \tilde{P})\|) \|\nabla(C - \tilde{C})\| d\tau \\ & \leq K_{10} \int_0^t (\|C - \tilde{C}\| + \|\tilde{C} - C\| + \|\nabla(P - \tilde{P})\|) \|\nabla(C - \tilde{C})\| d\tau \\ & \leq \frac{b_*}{4} \int_0^t \|\nabla(C - \tilde{C})(\tau)\|^2 d\tau + K_{11} \int_0^t \| (C - \tilde{C})(\tau) \|^2 d\tau \\ & + K_{12} \int_0^t \{ \| (C - \tilde{C})(\tau) \|^2 + \|\nabla(P - \tilde{P})(\tau)\|^2 \} d\tau \\ & < \frac{b_*}{4} \int_0^t \|\nabla(C - \tilde{C})(\tau)\|^2 d\tau + K_{11} \int_0^t \| (C - \tilde{C})(\tau) \|^2 d\tau + K_{13} \{ h^{2r} + h^{2s-2} \}. \end{aligned}$$

Here K_{11} depends upon M, K_1, K_5, K_6 , and b_* , while K_{13} depends upon $M, K_1, K_2, K_3, K_4, K_5, K_6$ and b_* . We can similarly obtain the bound

$$(3.10) \quad \left| \int_0^t ([u(c, \nabla p) - u(C, \nabla p)] \cdot \nabla \tilde{c}, C - \tilde{c}) d\tau \right| \leq K_{14} \int_0^t \| (C - \tilde{c})(\tau) \|^2 d\tau + K_{15} \{h^{2r} + h^{2s-2}\}.$$

Next, from (2.3), we note that

$$(3.11) \quad \left| \int_0^t (u(C, \nabla p) \cdot \nabla (C - \tilde{c}), C - \tilde{c}) d\tau \right| \leq K_{16} \{ \|\nabla p\|_{L^\infty(J; L^\infty)} + 1 \}^2 \int_0^t \| (C - \tilde{c})(\tau) \|^2 d\tau + \frac{b_*}{8} \int_0^t \|\nabla (C - \tilde{c})(\tau)\|^2 d\tau.$$

Then using Lemma 2.2, we obtain

$$(3.12) \quad \left| \int_0^t \left(\phi \left(\frac{\partial c}{\partial \tau} - \frac{\partial \tilde{c}}{\partial \tau} \right), C - \tilde{c} \right) d\tau + \lambda \int_0^t (\tilde{c} - c, C - \tilde{c}) d\tau \right| \leq K_{17} \int_0^t \| (C - \tilde{c})(\tau) \|^2 d\tau + K_{17} h^{2r}.$$

Combining like terms in the estimates (3.8)-(3.12) and the assumption (3.4), we obtain, from (3.7),

$$(3.13) \quad \begin{aligned} & \frac{\phi_*}{2} \| (C - \tilde{c})(t) \|^2 + \frac{b_*}{2} \int_0^t \|\nabla (C - \tilde{c})(\tau)\|^2 d\tau \\ & \leq K_{18} \int_0^t \| (C - \tilde{c})(\tau) \|^2 d\tau + K_{19} \{h^{2r} + h^{2s-2}\}, \end{aligned}$$

where K_{18} depends, in particular, upon K^* . Then applying Gronwall's lemma to (3.13), we obtain

$$(3.14) \quad \|C - \tilde{c}\|_{L^\infty(J; L^2)} + \|\nabla (C - \tilde{c})\|_{L^2(J; L^2)} \leq K_{20} \{h^r + h^{s-1}\},$$

where we note that

$$(3.15) \quad K_{20} \leq K_{21} \exp\{K_{22} K^{*2}\}.$$

To complete our argument, we must show that for h sufficiently small (3.6) holds.

We use Lemma 3.1, (2.2), (2.11), (3.14), and (3.15) to see that

$$\begin{aligned}
(3.16) \quad \|VP\|_{L^\infty(J;L^\infty)} &\leq \|V(P-\tilde{P})\|_{L^\infty(J;L^\infty)} + \|V\tilde{P}\|_{L^\infty(J;L^\infty)} \\
&\leq K_5 + K_0 h^{-1} \|V(P-\tilde{P})\|_{L^\infty(J;L^2)} \\
&\leq K_5 + K_0 h^{-1} K_6 (\|C-\tilde{C}\|_{L^\infty(J;L^2)} + \|\tilde{C}-C\|_{L^\infty(J;L^2)}) \\
&\leq K_5 + K_0 K_6 K_{21} \exp(K_{22} K^{*2}) \{h^{r-1} + h^{s-2}\} + K_0 K_2 K_4 K_6 h^{r-1}.
\end{aligned}$$

Then clearly if h is taken sufficiently small, and if $r \geq 2$, and $s \geq 3$; then

$$(3.17) \quad \|VP\|_{L^\infty(J;L^\infty)} \leq 2K_5 \leq K^*.$$

We see that, from (3.5), in order to get the best rates of convergence, one should use a space of piecewise polynomials of degree one greater for N_h than for M_h . For example, the use of piecewise linear polynomials ($r = 2$) for M_h and piecewise quadratic polynomials ($r = 3$) for N_h , would yield a rate of convergence of the form $O(h^2)$ for $C - \tilde{C}$.

We next combine the results of Lemma 3.1 and Theorem 3.2 with Lemmas 2.1 and 2.2 and the triangle inequality to obtain the following result.

Theorem 3.3. There exists a constant K_{23} such that

$$(3.18) \quad h \|V(P - p)\|_{L^\infty(J;L^2)} + \|C - c\|_{L^\infty(J;L^2)} + h \|V(C - c)\|_{L^2(J;L^2)} \leq K_{23} (h^r + h^{s-1}).$$

Some mathematical models for miscible displacements which are currently being used by oil companies make the assumption that the coefficient b in (1.2) does not depend upon Vp , but only upon x and c . Making this assumption, we obtain optimal order convergence rates in the L^2 norms using test spaces composed of piecewise polynomials of equal degrees. These results involve deriving a sharper estimate than obtained in (3.10).

Theorem 3.4. Assume $b = b(x, c)$ in (1.2) is independent of Vp . There exists a constant $K_{24} = K_{24}(\lambda, b_*, \phi_*, K_0, K_1, i \leq 6; K^*, M)$ such that

$$(3.19) \quad h \|V(P - p)\|_{L^\infty(J;L^2)} + \|C - c\|_{L^\infty(J;L^2)} + h \|V(C - c)\|_{L^2(J;L^2)} \leq K_{24} (h^r + h^s + h^{r+s-3}).$$

If either $r \geq 3$ or $s \geq 3$ for $r = 2$ and h is taken sufficiently small, then

$$(3.20) \quad \|\nabla p\|_{L^\infty(J; L^\infty)} \leq 2K_5 \leq K^*$$

and a K_{24} can be chosen which is independent of K^* .

Proof. Most of the estimates will follow as in the proof of Theorem 3.2. Let $\zeta = c - \tilde{c}$. The estimate (3.9) can be replaced by

$$(3.21) \quad \begin{aligned} \left| \int_0^t (b(c) - b(\tilde{c})) \nabla \tilde{c} \cdot \nabla \zeta \, d\tau \right| &\leq K_{25} \int_0^t (\|c - \tilde{c}\| + \|\zeta\|) \|\nabla \zeta\| \, d\tau \\ &\leq \frac{b_*}{8} \int_0^t \|\nabla \zeta(\tau)\|^2 \, d\tau + K_{25} \int_0^t \|\zeta(\tau)\|^2 \, d\tau + K_{26} h^{2r}. \end{aligned}$$

In order to obtain a sharper estimate for (3.10), we rewrite

$$(3.22) \quad \begin{aligned} ([u(c, \nabla p) - u(c, \nabla \tilde{p})] \cdot \nabla \tilde{c}, \zeta) &= ([u(c, \nabla p) - u(c, \nabla \tilde{p})] \cdot \nabla c, \zeta) \\ &+ ([u(c, \nabla p) - u(c, \nabla \tilde{p})] \cdot \nabla (\tilde{c} - c), \zeta) \\ &+ ([u(c, \nabla \tilde{p}) - u(c, \nabla p)] \cdot \nabla \tilde{c}, \zeta) = T_1 + T_2 + T_3. \end{aligned}$$

From Section 1, we note that for $u = u(c, q)$, $\frac{\partial u_i}{\partial q_i} = \frac{\partial u_i}{\partial q_i}(c)$ is independent of q , $i = 1, 2$.

Integrating the first term on the right-hand side of (3.22) by parts, we have

$$(3.23) \quad \begin{aligned} T_1 &= - \left(p - \tilde{p}, \frac{\partial}{\partial x_1} \left[\frac{\partial u_1}{\partial q_1}(c) \frac{\partial c}{\partial x_1} \zeta \right] + \frac{\partial}{\partial x_2} \left[\frac{\partial u_2}{\partial q_2}(c) \frac{\partial c}{\partial x_2} \zeta \right] \right) \\ &+ \int_{\partial \Omega} (p - \tilde{p}) \zeta \left[\frac{\partial u_1}{\partial q_1}(c) \frac{\partial c}{\partial x_1} v_1 + \frac{\partial u_2}{\partial q_2}(c) \frac{\partial c}{\partial x_2} v_2 \right] \, ds. \end{aligned}$$

Thus, from the assumed smoothness of u and c , we have

$$(3.24) \quad |T_1| \leq K_{27} \|p - \tilde{p}\| \|\zeta\|_1 + \left| \int_{\partial \Omega} (p - \tilde{p}) \zeta \left[\frac{\partial u_1}{\partial q_1}(c) \frac{\partial c}{\partial x_1} v_1 + \frac{\partial u_2}{\partial q_2}(c) \frac{\partial c}{\partial x_2} v_2 \right] \, ds \right|.$$

We now estimate the above boundary term by considering the following Neumann problem:

$$(3.25) \quad \begin{aligned} a) \quad & -\nabla \cdot a(x, c) \nabla \psi + \psi = 0, \quad x \in \Omega, \\ b) \quad & a(x, c) \frac{\partial \psi}{\partial \nu} = \Gamma, \quad x \in \partial \Omega, \end{aligned}$$

where

$$(3.26) \quad \Gamma = \zeta \left[\frac{\partial u_1}{\partial q_1} (c) \frac{\partial c}{\partial x_1} v_1 + \frac{\partial u_2}{\partial q_2} (c) \frac{\partial c}{\partial x_2} v_2 \right] \\ \equiv \zeta G_1 .$$

By our regularity assumptions (S) and (R), we have $\psi \in H^2(\Omega)$ and

$$(3.27) \quad \|\psi\|_2 \leq \kappa \|\Gamma\|_{H^{1/2}(\partial\Omega)} .$$

We note that since $c \in L^\infty(J; H^{2+\epsilon}(\Omega))$ and $\frac{\partial u_1}{\partial q_1} = a(c)$ is uniformly bounded, $G_1 \in L^\infty(J; H^{1/2+\epsilon}(\partial\Omega))$ and we have by Lemma 2.2 of [1],

$$(3.28) \quad \|\Gamma\|_{H^{1/2}(\partial\Omega)} \leq \kappa \|G_1\|_{L^\infty(J; H^{1/2+\epsilon}(\partial\Omega))} \|\zeta\|_{H^{1/2}(\partial\Omega)} \\ \leq \kappa \|\zeta\|_1 .$$

Multiplying (3.25.a) by $p - \tilde{p}$ and integrating by parts, we see that

$$(a(x, c) \nabla \psi, \nabla(p - \tilde{p})) + (\psi, p - \tilde{p}) - \int_{\partial\Omega} \Gamma(p - \tilde{p}) ds = 0 .$$

Using (2.6), we note that

$$(3.29) \quad \int_{\partial\Omega} \Gamma(p - \tilde{p}) ds = (a(x, c) \nabla(\psi - \psi^*), \nabla(p - \tilde{p})) + (\psi, p - \tilde{p}), \quad \psi^* \in N_h .$$

Hence, by (2.1), Lemma 2.1, (3.27), (3.28), and (3.29), we have

$$(3.30) \quad \left| \int_{\partial\Omega} \Gamma(p - \tilde{p}) ds \right| \leq \kappa_{28} \{h \|\nabla(p - \tilde{p})\| + \|p - \tilde{p}\|\} \|\psi\|_2 \leq \kappa_{29} \{h \|\nabla(p - \tilde{p})\| + \|p - \tilde{p}\|\} \|\Gamma\|_{H^{1/2}(\partial\Omega)} \\ \leq \kappa_{30} \{h \|\nabla(p - \tilde{p})\| + \|p - \tilde{p}\|\} \|\zeta\|_1 \leq \frac{b_*}{8} \|\zeta\|_1^2 + \kappa_{31} h^{2s} .$$

Combining (3.24), (3.30), and Lemma 2.1, we obtain

$$(3.31) \quad |T_1| \leq \frac{b_*}{8} \|\zeta\|_1^2 + \kappa_{32} \|\zeta\|^2 + \kappa_{33} h^{2s} .$$

Using Lemmas 2.2 and 3.1 we note that the third term on the right of (3.22) can be bounded as follows

$$(3.32) \quad |T_3| \leq \kappa_{34} \|\nabla \tilde{c}\|_{L^\infty(J; L^\infty)} \{ \|\nabla(\tilde{p} - p)\| + \kappa^* \|c - \tilde{c}\| \} \|\zeta\| \leq \kappa_{35} \|\zeta\|^2 + \kappa_{36} h^{2r} .$$

In order to bound the second term on the right side of (3.22), we use (2.2) and Lemmas 2.1 and 2.2. We obtain

$$\begin{aligned}
 |T_2| &\leq \kappa_{37} \|\nabla(p-\tilde{p})\|_{L^\infty(J;L^2)} \|\nabla(\tilde{c}-c)\|_{L^\infty} \|\zeta\|_{L^\infty} \leq \kappa_{38} h^{s-2} \|\nabla(\tilde{c}-c)\| \|\zeta\| \\
 (3.33) \quad &\leq \kappa_{39} h^{s-2} h^{r-1} \|\zeta\| \leq \kappa_{40} \|\zeta\|^2 + \kappa_{41} h^{2r+2s-6}.
 \end{aligned}$$

Using the estimates (3.8), (3.11), (3.21), (3.22), (3.31), (3.32), and (3.33) in (3.7), we obtain

$$\begin{aligned}
 (3.34) \quad &\frac{\phi_*}{2} \|\zeta(t)\|^2 + \frac{3b_*}{4} \int_0^t \|\nabla \zeta(\tau)\|^2 d\tau \leq \frac{b_*}{4} \int_0^t \|\zeta(\tau)\|_1^2 d\tau + \kappa_{42} \int_0^t \|\zeta(\tau)\|^2 d\tau \\
 &+ \kappa_{43} \{h^{2r} + h^{2s} + h^{2r+2s-6}\}.
 \end{aligned}$$

We can next add $\frac{3b_*}{4} \int_0^t \|\zeta(\tau)\|^2 d\tau$ to both sides of (3.34) and use the definition of the H^1 norm to hide the first term on the right of (3.34) on the left side of the resulting inequality. Then an application of Gronwall's lemma yields

$$(3.35) \quad \|\zeta\|_{L^\infty(J;L^2)} + \|\nabla \zeta\|_{L^2(J;L^2)} \leq \kappa_{44} \{h^r + h^s + h^{r+s-3}\}.$$

The rest of the theorem follows as in Theorem 3.2 and Theorem 3.3 with the minor modification of using (3.35) instead of (3.14) in (3.16) and the assumption that $r + s \geq 5$.

4. A Priori Error Estimates for the Discrete Time Approximations

In this section, we develop a priori bounds for the errors $c^n - z^n$ and $v(p^n - w^n)$ at discrete time levels t^n where z^n and w^n are defined in (2.15)-(2.16). The results obtained will be similar to the corresponding estimates for the continuous case from Section 3. We assume that S , Q , and R and the restrictions on $\{M_h\}$ and $\{N_h\}$ of Section 2 hold.

Theorem 4.1. Let z^0 be determined such that

$$a) \quad \|z^0 - c_0\| \leq K_{45} h^r$$

(4.0) and

$$b) \quad -\hat{\epsilon} \leq z^0 \leq 1 + \hat{\epsilon}.$$

There exists a constant $K_{46} = K_{46}(\lambda, b_*, \phi_*, M; K_i, i \leq 6)$ and constants $t_0 > 0$ and $h_0 > 0$ such that if $\Delta t \leq t_0$ and $h \leq h_0$,

$$(4.1) \quad \sup_{t^n} \|z - c\|^2 + h^2 \sum_{n=0}^{N-1} \|v(z - c)^n\|^2 \Delta t + h^2 \sup_{t^n} \|v(w - p)\|^2 \leq K_{46} \{(\Delta t)^2 + h^{2r} + h^{2(s-1)}\}.$$

Proof. Assume that z^n satisfies the induction hypothesis,

$$(4.2) \quad -\hat{\epsilon} \leq z^n \leq 1 + \hat{\epsilon}$$

for $n = 0, 1, \dots, l-1$. Let $\zeta^n = z^n - \tilde{c}^n$ and $\eta^n = w^n - \tilde{p}^n$. Subtracting (2.6) from (2.16) we have, for $n = 0, 1, \dots, N-1$,

$$(4.3) \quad \begin{aligned} (a(z^{n+1}) \nabla \eta^{n+1}, \nabla y) &= ([a(c^{n+1}) - a(z^{n+1})] \nabla \tilde{p}^{n+1}, \nabla y) \\ &+ ([a(z^{n+1}) \nabla (z^{n+1}) - a(c^{n+1}) \nabla (c^{n+1})] \nabla z, \nabla y), \quad y \in N_h. \end{aligned}$$

As in Lemma 3.1, we let $y = \eta^{n+1} \in N_h$, use (4.2) and (Q) to obtain

$$(4.4) \quad \|\nabla \eta^{n+1}\| \leq K_{47} \{\|\zeta^{n+1}\| + h^r\}, \quad n = 0, 1, \dots, N-1.$$

Next, subtract (2.9) from (2.15) and use the notation from Section 2 to obtain, for $n = 0, 1, \dots, N-1$, and $j = 0$ or 1 ,

$$\begin{aligned}
(4.5) \quad & (\phi d_t \zeta^n, \chi) + (b(Z^n, \nabla W^n) \nabla \zeta^{n+1}, \nabla \chi) = (\phi [\frac{\partial c}{\partial t} (t^{n+1}) - d_t \tilde{c}^n], \chi) \\
& - \lambda((c - \tilde{c})^{n+1}, \chi) + ([b(c^{n+1}, \nabla p^{n+1}) - b(Z^n, \nabla W^n)] \nabla \tilde{c}^{n+1}, \nabla \chi) \\
& + ([u(c^{n+1}, \nabla p^{n+1}) - u(Z^n, \nabla W^n)] \cdot \nabla \tilde{c}^{n+1}, \chi) \\
& - (u(Z^n, \nabla W^n) \cdot \nabla \zeta^{n+1}, \chi), \quad \chi \in M_h.
\end{aligned}$$

For simplicity, let

$$(4.6) \quad \|f\|_*^2 = (\phi f, f).$$

Now letting $\chi = \zeta^{n+1} \in M_h$ and using (4.6), we estimate the left side of (4.5) by

$$\begin{aligned}
(4.7) \quad & \frac{1}{\Delta t} (\phi (\zeta^{n+1} - \zeta^n), \zeta^{n+1}) + (b(Z^n, \nabla W^n) \nabla \zeta^{n+1}, \nabla \zeta^{n+1}) \\
& \geq \frac{1}{2\Delta t} (\|\zeta^{n+1}\|_*^2 - \|\zeta^n\|_*^2) + b_* \|\nabla \zeta^{n+1}\|^2.
\end{aligned}$$

We next use Lemmas 2.1 and 2.2, (2.11), and (4.4) to see that

$$\begin{aligned}
(4.8) \quad & |([b(c^{n+1}, \nabla p^{n+1}) - b(Z^n, \nabla W^n)] \nabla \tilde{c}^{n+1}, \nabla \zeta^{n+1})| \\
& = |([b(c^{n+1}, \nabla p^{n+1}) - b(c^n, \nabla p^{n+1}) + b(c^n, \nabla p^{n+1}) - b(Z^n, \nabla p^{n+1}) \\
& \quad + b(Z^n, \nabla p^{n+1}) - b(Z^n, \nabla p^n) + b(Z^n, \nabla p^n) - b(Z^n, \nabla W^n)] \nabla \tilde{c}^{n+1}, \nabla \zeta^{n+1})| \\
& \leq \kappa_{50} \{\Delta t \|d_t c^n\| + \|(c - \tilde{c})^n\| + \|\zeta^n\| + \Delta t \|d_t \nabla p^n\| + \|\nabla(p - \tilde{p})^n\| \\
& \quad + \|\nabla \eta^n\|\} \cdot \|\nabla \zeta^{n+1}\| \\
& \leq \kappa_{51} \|\zeta^n\|^2 + \frac{b_*}{4} \|\nabla \zeta^{n+1}\|^2 + \kappa_{52} (\Delta t)^2 \{\|d_t c^n\|^2 + \|d_t \nabla p^n\|^2\} + \kappa_{53} \{h^{2r} + h^{2(s-1)}\}.
\end{aligned}$$

In a similar fashion, we obtain

$$\begin{aligned}
(4.9) \quad & |([u(c^{n+1}, \nabla p^{n+1}) - u(Z^n, \nabla W^n)] \cdot \nabla \tilde{c}^{n+1}, \zeta^{n+1})| \\
& \leq \frac{\phi_*}{16} \|\zeta^{n+1}\|^2 + \kappa_{54} \|\zeta^n\|^2 + \kappa_{55} (\Delta t)^2 \{\|d_t c^n\|^2 + \|d_t \nabla p^n\|^2\} \\
& \quad + \kappa_{56} \{h^{2r} + h^{2(s-1)}\}.
\end{aligned}$$

Next, from (2.3 f), we note that

$$(4.10) \quad |-(u(Z^n, \nabla W^n) \cdot \nabla \zeta^{n+1}, \zeta^{n+1})| \leq \kappa_{57} \|\nabla W^n\|_{L^\infty(\Omega)}^2 + \kappa_{58} \|\zeta^{n+1}\|^2 + \frac{b_*}{4} \|\nabla \zeta^{n+1}\|^2.$$

We then estimate the first two terms on the right side of (4.5) using Lemma 2.2 as follows:

$$\begin{aligned}
 & |(\phi[\frac{\partial c}{\partial t}(t^{n+1}) - d_t c^n + d_t c^n - d_t \tilde{c}^n], \zeta^{n+1}) - \lambda((c - \tilde{c})^{n+1}, \zeta^{n+1})| \\
 (4.11) \quad & \leq \frac{\phi_*}{16} \|\zeta^{n+1}\|^2 + \kappa_{59} (\|(c - \tilde{c})^{n+1}\|^2 + \|d_t(c - \tilde{c})^n\|^2) + \kappa_{60} \sigma_n^2 \\
 & \leq \frac{\phi_*}{16} \|\zeta^{n+1}\|^2 + \kappa_{61} h^{2r} + \kappa_{60} \sigma_n^2,
 \end{aligned}$$

where

$$(4.12) \quad \sigma_n^2 = \left(\int_{t^n}^{t^{n+1}} \left\| \frac{\partial^2 c}{\partial t^2}(\cdot, s) \right\| ds \right)^2 \leq \Delta t \int_{t^n}^{t^{n+1}} \left\| \frac{\partial^2 c}{\partial t^2}(\cdot, s) \right\|^2 ds$$

by Schwarz's inequality. We note that

$$(4.13) \quad \Delta t \sum_{n=0}^{N-1} \sigma_n^2 \leq (\Delta t)^2 \sum_{n=0}^{N-1} \int_{t^n}^{t^{n+1}} \left\| \frac{\partial^2 c}{\partial t^2}(\cdot, s) \right\|^2 ds \leq (\Delta t)^2 \left\| \frac{\partial^2 c}{\partial t^2} \right\|_{L^2(J; L^2)}^2.$$

We next multiply each of the estimates (4.7)-(4.11) by Δt and sum on n , $n = 0, \dots, \ell - 1$, to obtain from (4.5)

$$\begin{aligned}
 & \frac{1}{2} (\phi_* \|\zeta^\ell\|^2 - \phi_* \|\zeta^0\|^2) + b_* \sum_{n=0}^{\ell-1} \|\nabla \zeta^{n+1}\|^2 \Delta t \\
 (4.14) \quad & \leq \sum_{n=0}^{\ell-1} \frac{1}{2} (\|\zeta^{n+1}\|_*^2 - \|\zeta^n\|_*^2) + b_* \sum_{n=0}^{\ell-1} \|\nabla \zeta^{n+1}\|^2 \Delta t \\
 & \leq \kappa_{62} \sum_{n=0}^{\ell-1} \|\zeta^n\|^2 \Delta t + \frac{b_*}{2} \sum_{n=0}^{\ell-1} \|\nabla \zeta^{n+1}\|^2 \Delta t + \Delta t \frac{\phi_*}{8} \|\zeta^\ell\|^2 \\
 & \quad + \kappa_{57} \sum_{n=0}^{\ell-1} (\|\nabla w^n\|_{L^\infty(\Omega)} + \kappa_{58})^2 \|\zeta^{n+1}\|^2 \Delta t + \kappa_{63} \{(\Delta t)^2 + h^{2r} + h^{2(s-1)}\}.
 \end{aligned}$$

We see that an application of Gronwall's lemma would complete our argument if we did not have the next to the last term on the right of (4.14). We shall use an induction argument to treat this term. As an induction hypothesis assume that for h sufficiently small

$$(4.15) \quad \|\nabla w^n\|_{L^\infty(\Omega)} \leq 2\kappa_5, \quad \text{for } n = 0, 1, \dots, \ell - 1.$$

Using (2.2), Lemma 2.2, (2.11), (4.0), and (4.4) as in (3.16) we see that (4.15) clearly holds for $l = 1$. Using (4.15) we see that if

$$(4.16) \quad \Delta t \leq \phi_*(4K_{57}(2K_5 + K_{59})^2)^{-1} \equiv t_0,$$

then we have

$$(4.17) \quad \|z^l\|^2 + b_* \sum_{n=0}^{l-1} \|vz^{n+1}\|^2 \Delta t \leq K_{64} \sum_{n=0}^{l-1} \|z^n\|^2 \Delta t + K_{65} \{(\Delta t)^2 + h^{2r} + h^{2(s-1)}\}.$$

Then an application of Gronwall's lemma gives us an estimate which, with (2.2), Lemma 2.2 (2.11), and (4.4) shows as in (3.16), that the induction hypotheses (4.2) and (4.15) holds for $n = l$. Finally an application of the triangle inequality coupled with Lemmas 2.1 and 2.2 and (4.4) yields the desired result.

Just as in Section 3, if we make the assumption that $b = b(x, c)$ in (1.2) is independent of Vp , we can obtain optimal order L^2 rates of convergence. The proof will follow by combining the techniques of Theorem 3.4 with those of Theorem 4.1. We obtain the following result.

Theorem 4.2. Let the assumptions of Theorem 4.1 hold. Assume $b = b(x, c)$ in (1.2) is independent of Vp . There exist constants K_{66} , t_0 , and h_0 such that, if $\Delta t \leq t_0$ and $h \leq h_0$,

$$(4.18) \quad \sup_{t^n} \|z - c\|^2 + h^2 \sum_{n=0}^{N-1} \|v(z - c)^n\|^2 \Delta t + h^2 \sup_{t^n} \|v(w - p)\|^2 \leq K_{66} \{(\Delta t)^2 + h^{2r} + h^{2s} + h^{2r+2s-6}\}.$$

As we mentioned at the end of Section 2, if we replace (2.15) by (2.17) and extrapolate the coefficients, we can replace the $(\Delta t)^2$ in (4.17) and (4.18) by $(\Delta t)^4$ and get discretization errors in time which are $O((\Delta t)^2)$. In order to do this we must determine a starting procedure to obtain the approximation at time $t^1 = \Delta t$ since two levels must be known to determine the next level with this method. A predictor-corrector version of (2.15) will suffice as a starting procedure of sufficient accuracy. Since the proof techniques are similar to those presented in Theorem 4.1 and are

similar to those presented in [1], [2], [5], [8] and [10], we shall not present them here. We require slightly more smoothness on c and p for these results.

Theorem 4.3. If we have the added smoothness on c and p that

$$(4.19) \quad \left\| \frac{\partial^3 c}{\partial t^3} \right\|_{L^2(J; L^2)} + \left\| \frac{\partial^2 v p}{\partial t^2} \right\|_{L^2(J; L^2)} \leq K_{67}$$

and if z^0 and z^1 are determined to satisfy

$$(4.20) \quad a) \quad \|z^0 - c_0\| + \|z^1 - c(t^1)\| \leq K_{68} h^r,$$

and

$$b) \quad -\hat{\epsilon} \leq z^n \leq 1 + \hat{\epsilon}, \quad n = 0, 1,$$

then there exist constants K_{69} , t_0 , and h_0 such that, if $\Delta t \leq t_0$ and $h \leq h_0$,

$$(4.21) \quad \sup_{t^n} \|\tilde{z} - c\|^2 + h^2 \sum_{n=0}^{N-1} \|v(\tilde{z} - c)^n\|^2 \Delta t + h^2 \sup_{t^n} \|v(W - p)\|^2 \leq K_{69} \{(\Delta t)^4 + h^{2r} + h^{2s-2}\}.$$

Finally, in the case that $b = b(x, c)$, we can use the techniques of Theorem 4.2 to obtain the following result for $\tilde{z}$ defined in (2.17).

Theorem 4.4. Under the assumptions of Theorem 4.3, there exist constants K_{70} , t_0 , and h_0 such that, if $\Delta t \leq t_0$ and $h \leq h_0$, we have

$$\sup_{t^n} \|\tilde{z} - c\|^2 + h^2 \sum_{n=0}^{N-1} \|v(\tilde{z} - c)^n\|^2 \Delta t + h^2 \sup_{t^n} \|v(W - p)\|^2 \leq K_{70} \{(\Delta t)^4 + h^{2r} + h^{2s}\}.$$

REFERENCES

1. Douglas, J., Jr. and Dupont, T., "Galerkin methods for parabolic equations with non-linear boundary conditions," Numer. Math., 20 (1973), pp. 213-237.
2. Douglas, J., Jr., Dupont, T., and Ewing, R. E., "Incomplete iteration for time-stepping a nonlinear parabolic Galerkin method," SIAM J. Numer. Anal. (to appear).
3. Dupont, T., Fairweather, G., and Johnson, P. J., "Three-level Galerkin methods for parabolic equations," SIAM J. Numer. Anal., (1974), pp. 392-410.
4. Dupont, T., " L_2 error estimates for projection methods for parabolic equations in approximating domains," Mathematical Aspects of Finite Elements in Partial Differential Equations, Academic Press, Inc., New York, 1974, pp. 313-352.
5. Ewing, R. E., "Time-stepping Galerkin methods for nonlinear Sobolev partial differential equations," SIAM J. Numer. Anal. (to appear).
6. Peaceman, D. W., "Improved treatment of dispersion in numerical calculation of multidimensional miscible displacement," Soc. Pet. Engr. J., (Sept. 1966), pp. 213-216.
7. Peaceman, D. W., Fundamentals of Numerical Reservoir Simulation, Elsevier Publishing Company, 1977.
8. Rachford, H. H., Jr., "Two-level discrete-time Galerkin approximations for second order nonlinear parabolic partial differential equations," SIAM J. Numer. Anal., 10 (1973), pp. 1010-1026.
9. Settari, A., Price, H. S., and Dupont, T., "Development and application of variational methods for simulation of miscible displacement in porous media," Soc. Pet. Engr. J., (June 1977), pp. 228-246.
10. Wheeler, M. F., "A priori L^2 -error estimates for Galerkin approximations to parabolic partial differential equations," SIAM J. Numer. Anal., 10 (1973), pp. 723-759.

REE/MFW/scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1932	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) GALERKIN METHODS FOR MISCIBLE DISPLACEMENT PROBLEMS IN POROUS MEDIA		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
6. AUTHOR(s) Richard E. Ewing and Mary Fanett/Wheeler		7. PERFORMING ORG. REPORT NUMBER
8. CONTRACT OR GRANT NUMBER(s) DAAG29-78-G-0161 DAAG29-75-C-0024 MCS78-09525, MCS78-00913		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 7 (Numerical Analysis)
10. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below.
12. REPORT DATE March 1979		13. NUMBER OF PAGES 21
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) MRC-TSR-1932		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DAAG29-75-C-0024, DAAG29-78-G-0161		
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709 National Science Foundation Washington, D. C. 20550		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Galerkin methods Error estimates Fluid flow Numerical analysis 9 Technical summary rept.,		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A priori error estimates for Galerkin methods for numerical approximation of the coupled quasilinear system for $c = c(x, t)$ and $p = p(x, t)$ given by $\nabla \cdot [a(x, c) \{ \nabla p - \gamma(x, c) \nabla z \}] = 0,$ $\nabla \cdot [b(x, c, \nabla p) \nabla c] - u(x, c, \nabla p) \cdot \nabla c = \phi(x) \frac{\partial c}{\partial t},$		

next
Page

ell

20. ABSTRACT - Cont'd.

for $x \in \Omega$, $t \in (0, T]$, and appropriate Neumann boundary and initial conditions are considered. Equations of this type arise in models for the miscible displacement of one incompressible fluid by another in a porous medium. Estimates for both continuous time and fully-discrete time Galerkin methods are presented.