

AD-A068 948

PENNSYLVANIA STATE UNIV UNIVERSITY PARK DEPT OF CHEMISTRY F/G 7/3
FORMATION OF HYDRIDOCYCLOTRIPHOSPHAZENES VIA THE REACTIONS OF O--ETC(U)
APR 79 H R ALLCOCK, P J HARRIS N00014-75-C-0685

NL

UNCLASSIFIED

| OF |

AD
A068948



END
DATE
FILMED
7-79
DDC

AD A 068948

DDC FILE COPY

UNCLASSIFIED

LEVEL II

12

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
(6) FORMATION OF HYDRIDOCYCLOTRIPHOSPHAZENES VIA THE REACTIONS OF ORGANOCOPPER REAGENTS WITH HALOCYCLOTRIPHOSPHAZENES: THE REACTION MECHANISM		(7) Interim Technical Report
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
(10) Harry R./Allcock* and Paul J./Harris		(15) N00014-75-C-0685
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
Department of Chemistry, The Pennsylvania State University, University Park, Pennsylvania 16802		NR 356-577
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
Department of the Navy Office of Naval Research, Arlington, Va. 22217		(11) 26 Apr 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
(12) 4 pp.		39
16. DISTRIBUTION STATEMENT (of this Report)		18. SECURITY CLASS. (of this report)
Distribution unlimited; approved for publication		Unclassified
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE
DDC RECEIVED MAY 24 1979 D		
19. SUPPLEMENTARY NOTES		
Submitted for publication to the Journal of the American Chemical Society		
20. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Phosphazenes, hydridophosphazenes, alkylphosphazenes, model reactions, copper reagents, synthesis, mechanisms		
21. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
A new method is reported for the synthesis of cyclophosphazenes that contain alkyl and hydrido side groups. These syntheses are model reactions for the synthesis of a new class of high polymeric organophosphazenes.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

400 343

Jul 79

05 18 021

Office of Naval Research
Contract No. N00014-75-C-0685
Project No. NR 356-577

FORMATION OF HYDRIDOCYCLOTRIPHOSPHAZENES VIA THE REACTIONS
OF ORGANOCOPPER REAGENTS WITH HALOCYCLOTRIPHOSPHAZENES:
THE REACTION MECHANISM

by

Harry R. Allcock* and Paul J. Harris

Submitted for publication in the
Journal of the American Chemical Society

ADDITIONAL	
DTIC	White Section ☒
DDI	Defi Section ☐
UNCLASSIFIED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A	

Department of Chemistry
The Pennsylvania State University
University Park, Pennsylvania 16802

April 26, 1979

Reproduction in whole or in part is permitted for any
purpose of the United States Government

Approved for public release; Distribution unlimited

Formation of Hydridocyclotriphosphazenes via the Reactions of
Organocopper Reagents with Halocyclotriphosphazenes: The
Reaction Mechanism¹⁻³

Harry R. Allcock* and Paul J. Harris

Contribution from the Department of Chemistry, The Pennsylvania
State University, University Park, Pennsylvania 16802.
Received

Abstract: 1-Hydrido-1-alkyl-tetrachlorocyclotriphosphazenes, $N_3P_3Cl_4RH$ (II) (where $R = CH_3, C_2H_5, n-C_3H_7, i-C_3H_7, n-C_4H_9, i-C_4H_9, t-C_4H_9$), have been synthesized by the new reaction of hexachlorocyclotriphosphazene, $(NPCl_2)_3$, with alkyl Grignard reagents in the presence of $[(n-C_4H_9)_3PCuI]_4$, followed by treatment with 2-propanol. The structural characterization of these compounds is discussed together with a detailed study of the reaction mechanism that permits hydridophosphazene formation. These reactions proceed via the formation of metallophosphazene intermediates. The nature of these complexes is discussed.

An important need exists for the development of new synthetic methods for the preparation of cyclic or open-chain phosphazenes that contain alkyl or aryl groups bonded directly to the skeleton through phosphorus-carbon bonds. Cyclic alkyl- or aryl-phosphazenes are of interest for fundamental reactivity studies, as models for thermally stable high polymers, or as "monomers" for polymerization studies.⁴ The high polymers themselves are expected to constitute a new class of useful macromolecules.

A number of examples have been reported in the literature of reactions between halophosphazenes and Grignard or organolithium reagents.⁵⁻¹⁸ Such reactions lead to alkylation or arylation, or to cleavage of the phosphazene skeleton. In many cases the substitution and cleavage reactions occur concurrently.¹⁹ Our objective in this work was to find an alkylation or arylation system that would permit halogen atoms attached to a phosphazene ring or chain to be replaced in a controlled manner, without cleavage of the skeleton.


Organocopper reagents were chosen for this purpose because such species permit the selective replacement of halogen atoms in organic compounds by various alkyl or aryl residues.^{20,21} The wide scope and effectiveness of these reagents has been discussed elsewhere.²⁰⁻²⁹ Of all the methods available for the utilization of organocopper reagents, the "copper catalyzed" Grignard reaction is the easiest to use. Hence, this system was chosen for our study.

Although the synthesis of alkyl phosphazenes was the objective of this work, it was discovered that reactions of this type provided a new and unusual route for the synthesis of hydrido-phosphazene-species,³ compounds that have

been synthesized before by only one reaction pathway.³⁰⁻³³ Moreover, the synthesis described here is the only available method for the preparation of phosphazenes that contain both hydrido and halogeno groups. The use of hydridophosphazenes as synthetic intermediates is a subject of some consequence.

Results and Discussion

The Overall Reaction. When hexachlorocyclotriphosphazene, $(\text{NPCl}_2)_3$ was allowed to react with alkyl Grignard reagents in the presence of $[\text{n-Bu}_3\text{PCuI}]_4$,³⁴ followed by treatment with 2-propanol, cyclophosphazenes were formed that possessed an alkyl group and a hydrido unit linked to one phosphorus atom (II).



Compounds of structure II were prepared in which R is CH_3 , C_2H_5 , $\text{n-C}_3\text{H}_7$, $\text{i-C}_3\text{H}_7$, $\text{n-C}_4\text{H}_9$, $\text{i-C}_4\text{H}_9$, and $\text{t-C}_4\text{H}_9$. All these compounds are air- and moisture-sensitive, volatile products. When R is CH_3 , C_2H_5 , $\text{n-C}_3\text{H}_7$, $\text{i-C}_3\text{H}_7$, $\text{n-C}_4\text{H}_9$ or $\text{t-C}_4\text{H}_9$, they are white, crystalline compounds. The $\text{i-C}_4\text{H}_9$ product is a colorless oil.

Proof of Structure II. The structure of the hydrido-cyclotriphosphazenes (II) was determined unambiguously from ^{31}P and ^1H NMR spectra. The discussion of this structural proof will be limited to that of the methyl-derivative, $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$. The data for the other products are shown in Tables I and II.

Figure I shows the ^{31}P NMR spectrum of $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$.^{35,36} It is a simple AB_2 spin system,³⁷ with peaks at 17.6 ppm³⁸ (PCl_2 as a doublet with $J_{\text{PNP}} = 12$ Hz) and 13.8 ppm (PHCH_3 as a triplet with $J_{\text{PNP}} = 12$ Hz). The peak at 13.8 ppm was found to be significantly larger than expected, because of the nuclear Overhauser effect of the hydrogen atom bound to this phosphorus, and other relaxation effects.³⁹ In the proton undecoupled spectrum (Figure 2) the peak at 13.8 ppm was split into a doublet ($J_{\text{PH}} = 568$ Hz), with additional fine structure from the PNP and PCH couplings.

The phosphorus decoupled ^1H NMR spectrum⁴¹ of this compound (Figure 3) showed a CH_3 resonance at δ 1.75⁴² (a doublet with $J_{\text{HPCH}} = 2.9$ Hz⁴³) and the hydride resonance at δ 7.44 (a quartet with $J_{\text{HCPH}} = 2.9$ Hz). The phosphorus-undecoupled ^1H NMR spectrum (Figure 4) showed the CH_3 resonance split into a doublet of multiplets ($J_{\text{PCH}} = 16$ Hz, $J_{\text{PNPCH}} = 4$ Hz, $J_{\text{HPCH}} = 2.9$ Hz). In this spectrum the hydride resonance appeared as a doublet of triplets with additional fine structure ($J_{\text{PH}} = 568$ Hz, $J_{\text{PNPH}} = 11$ Hz, and $J_{\text{HCPH}} = 2.9$ Hz). A further confirmation of these coupling constants was obtained from ^1H homo-decoupled experiments.


Although the NMR spectra provided unambiguous evidence for structure II, further confirmation of the structure was obtained from infrared spectra,⁴⁴ mass spectra⁴⁵, and from microanalytical data. This information is summarized in Table II and in the Experimental section. Moreover, the structure of $\text{N}_3\text{-P}_3\text{Cl}_4(\text{i-C}_3\text{H}_7)\text{H}$ was confirmed by a single crystal X-ray structure determination. This will be discussed in a later paper.

The Reaction Mechanism

The formation of hydridocyclotriphosphazenes (II) from these reactions was a surprising result that raised a number of questions. (1) What is the source of the hydrogen atom that yields the hydridophosphazene structure? (2) Is copper an essential prerequisite for this reaction and, if so, how does the copper participate in the synthesis? (3) What is the reaction pathway that leads to the formation of II?

Source of Hydrogen Atom Bound to Phosphorus. Initially, a number of experiments were carried out to determine the effect of Grignard concentration of the formation of II. The sequence of operations involved the addition of successive amounts of methylmagnesium chloride to a solution of $(\text{NPCl}_2)_3$ (I) and $[\text{n-Bu}_3\text{PCuI}]_4$, followed by vapor phase chromatographic analysis of the reaction mixture.⁴⁶ Surprisingly, it was found that although the $(\text{NPCl}_2)_3$ was consumed in direct proportion to the amount of Grignard reagent added (up to 3-3.5 equivalents per $(\text{NPCl}_2)_3$ molecule), no trace of II was detected, even after 24 h of reaction. However, when 2-propanol was added to the reaction mixture (ostensibly to destroy the excess Grignard reagent), compound II ($\text{R} = \text{CH}_3$) could be detected by vapor phase chromatography.⁴⁶

This reaction sequence was then repeated with the use of $(\text{CH}_3)_2\text{CHOD}$. The product shown in III was isolated.




III

The incorporation of deuterium into III was demonstrated by the following results. A mass spectrum⁴⁵ showed a parent ion at m/e 292 as the base peak of the characteristic Cl_4 isotope pattern.⁴⁷ An infrared spectrum showed no band at 2400 cm^{-1} (P-H stretch) but showed instead a band at 1760 cm^{-1} that was assigned to the P-D stretch. The ratio ν_{PH}/ν_{PD} should be $\approx\sqrt{2}$ on the basis of a Hooke's law analysis. With compounds II ($R = CH_3$) and III the ratio is 1.363. The ^{31}P NMR spectrum³⁶ (in $CDCl_3$ solution with proton decoupling) showed a peak at 18.6 ppm (PCl_2 as a doublet with $J_{PNP} = 12\text{ Hz}$) and a peak at 11.4 ppm ($PDCH_3$ as a triplet, with $J_{PD} = 85\text{ Hz}$, of triplets with $J_{PNP} = 12\text{ Hz}$). The phosphorus-deuterium coupling constant was confirmed from the 2D NMR spectrum⁴⁸ (in C_6F_6 solution with proton decoupling) which showed the deuterium resonance as a doublet, with $J_{PD} = 85\text{ Hz}$, centered at 1.71 ppm downfield from external C_6D_6 . The observed difference between J_{PH} and J_{PD} is in good agreement with that found in other cases.^{33,49} Finally, the 1H NMR spectrum showed no P-H resonance, but gave the CH_3 resonance at δ 1.75 as a doublet of triplets ($J_{PCH} = 16\text{ Hz}$, $J_{PNPCH} = 4\text{ Hz}$), and this coupling pattern is in marked contrast to the multiplet structure observed for the methyl resonance of II ($R = CH_3$), as shown in Figure 4.

The incorporation of the deuterium atom into III showed unequivocally that the alcohol is the source of the hydrogen atom bound to the phosphorus in II. This led to the supposition that a non-volatile metallophosphazene intermediate (i.e. non-volatile in a vapor phase chromatography experiment) was present in solution before the addition of the alcohol. This prospect is considered in the following sections.

Involvement of Copper in the Reaction. The degree to which the formation of II was dependent on the concentration of $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ was explored by means of the following experiments. A series of reactions was performed in which the molar ratio of methylmagnesium chloride to $(\text{NPCl}_2)_3$ was maintained at 4:1 but in which the amount of $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ was increased. The results are shown in Figure 5. The maximum yields of II ($\text{R} = \text{CH}_3$) were obtained with copper atom to phosphazene ratios of more than 0.5:1.

When very low copper to phosphazene ratios ($<0.15:1$) were employed, it was found after the usual isolation procedure that the volatile products⁵⁰ consisted mainly of compounds IV and V rather than II ($\text{R} = \text{CH}_3$).



Compounds IV and V are normally formed from the reaction of methylmagnesium chloride with $(\text{NPCl}_2)_3$ in the absence of copper. In such reactions the remainder of the product consists of ring-cleaved phosphazenes.⁵¹ Thus, it appears that, at low copper concentrations, the trace of copper is removed by coordination to the phosphazene. The Grignard reagent then functions as if no copper were present, and reacts directly with the large amount of $(\text{NPCl}_2)_3$ in the system.

These results indicate that the copper does not play a catalytic role in the formation of II. On the contrary, it participates in the formation of a


complex that functions as a reaction intermediate. In this respect, the copper behaves in a different manner from that normally assumed for many Grignard-copper system reactions.

Nature of the Metallo-Phosphazene Intermediate. Two plausible possibilities exist for the mechanism of formation of a reactive copper-phosphazene intermediate. The intermediate could be generated by a direct interaction of $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ with $(\text{NPCl}_2)_3$, or the mechanism could involve the essential participation of the Grignard reagent.

Two reactions were carried out in order to distinguish between these possibilities. In the first, $(\text{NPCl}_2)_3$ and $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ were allowed to interact in stirred tetrahydrofuran. After 48 h, no reaction had taken place, as deduced by the absence of a change in the ^{31}P NMR spectrum of the reaction mixture. In the second reaction, increasing amounts of methylmagnesium chloride were added to a reaction mixture that contained $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ and $(\text{NPCl}_2)_3$ in a copper atom to phosphazene ratio of 1:2. At the same time, the ^{31}P NMR spectrum was monitored. The changes are shown in Figure 6a. 2-Propanol was then added to each mixture (in the NMR tube) and the ^{31}P NMR spectrum was again obtained. These results are shown in Figure 6b. The spectra show clearly that the metallo-phosphazene complex is not formed until the Grignard reagent is added. Complexation is complete after the addition of a 3:1 excess of Grignard to phosphazene. The spectra also illustrate how treatment of this intermediate with 2-propanol leads immediately to the formation of II ($\text{R} = \text{CH}_3$).

The metallo-phosphazene intermediate was not isolated from the reaction mixture. However, we believe that its structure is similar to the one shown in

VI. The evidence for this structure is as follows.




VI


First, the stoichiometry of two phosphazene rings per copper atom was established from the data shown in Figure 5. The same conclusion is derived from the ^{31}P NMR results just described, in which a 2:1 ratio of phosphazene to copper was employed.

Second, the "cuprate" nature of the intermediate can be inferred from its solubility in organic solvents (THF and diethyl ether) and from the fact that the metal must occupy a +1 oxidation state, [Copper +2 is reduced to Copper +1 in the presence of organometallic reagents²⁰].

Third, it seems reasonable to assume that the copper is bound to the phosphazene rings through the skeletal nitrogen atoms since these almost certainly represent the most electronegative sites in the molecule. This fact has been demonstrated elegantly by Schmidpeter and coworkers³² by their isolation of compound VIII from the reaction of VII with $n\text{-C}_4\text{H}_9\text{Li}$ (or potassium).



VII



VIII

These same authors showed that VIII could be coupled to another phosphazene ring by means of a reaction with a P-Cl bond.³² By contrast, species VI does not couple to a chlorophosphazene molecule, and this is compatible with a significant degree of covalent character in the phosphazene-copper bond.

Fourth, additional evidence that VI contains copper-nitrogen bonds was obtained from the ³¹P NMR spectrum (VI in Figure 6a). This spectrum shows three distinct phosphorus resonances, all broadened apparently due to coupling with the copper. This pattern is compatible with coordination to nitrogen rather than to tricoordinate phosphorus, where a simple AB₂ spin system would be expected.


Of course, the possibility of additional coordination of copper to the phosphazene ring through the other skeletal nitrogen atoms or through the tricoordinate phosphorus atom cannot be ruled out. Moreover, the possibility also exists that the other coordination sites at copper could be filled by n-Bu₃P molecules or by tetrahydrofuran, the reaction solvent.

Reactions with Methyl Copper. In view of the mechanistic implications discussed in the preceding sections, it was of interest to determine if alkyl copper reagents would react with $(\text{NPCl}_2)_3$ in a manner different from that found for $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$.

Methyl copper was obtained by the reaction of methylmagnesium chloride with $[\underline{n}\text{-Bu}_3\text{CuI}]_4$ at -80°C .²¹ A reaction with $(\text{NPCl}_2)_3$ in tetrahydrofuran at -80° to 25°C , followed by treatment with 2-propanol, yielded the cyclophosphazenes,⁵⁰ $\text{N}_3\text{P}_3\text{Cl}_6$ (I), $\text{N}_3\text{P}_3\text{Cl}_5\text{H}$, and $\text{N}_3\text{P}_3\text{Cl}_4\text{CH}_3\text{H}$ (II). The isolation of $\text{N}_3\text{P}_3\text{Cl}_5\text{H}$ from this reaction system was a valuable clue to the reaction mechanism, as discussed in the following section.


The Reaction Pathway. The exact reaction pathway that leads to the formation of the metallo-phosphazene and, eventually, to the hydridophosphazene remains partly unsolved. For example, it is not known whether radical or ionic mechanisms are involved in the initial steps. However, sufficient evidence has been obtained to allow the overall mechanism to be understood.

It is presumed that, although $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ is unreactive with $(\text{NPCl}_2)_3$, the reaction of $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ with the Grignard reagent yields an alkyl copper compound, R-Cu . The yellow precipitate observed in these reactions is believed to be the alkyl copper compound. A reaction is then assumed to occur between the alkyl copper reagent and $(\text{NPCl}_2)_3$, as shown in IX and X.




Significant quantities of the alkyl halides, R-Cl, were detected from the reaction mixture by the use of VPC/mass spectrometric analysis.⁵⁰ The coupled species, R-R, was also detected,⁵⁰ probably as a product from the reaction of R-Cl with the Grignard reagent or the organocopper compound.²⁰ In these terms, the formation of $N_3P_3Cl_5H$ by the interaction of $(NPCl_2)_3$ with methylcopper followed by protonation with 2-propanol provides evidence that a species of type X has a discrete existence after alkyl halide elimination occurs.

This metal-halogen exchange process is the key to the unique behavior of the organocopper reagent with chlorophosphazenes. Grignard reagents react with chlorophosphazenes to cause skeletal cleavage (XI), as well as alkyl-halogen exchange (XIII), because the alkyl group attacks at phosphorus (XII)^{19,51,52} rather than at chlorine. By contrast, the alkylcopper species attacks the chlorine atom. At the same time, the copper atom is better able to stabilize the negative charge on nitrogen than is magnesium.



The initial metal-halogen exchange reaction is believed to generate X, a species with a trivalent, tricoordinate phosphorus atom. The chlorine remaining on this phosphorus can then be replaced readily by an alkyl group (in the form of R^-) from either the Grignard or the organocopper reagent, to yield XIV.



The stoichiometry of the reaction requires that two phosphazene rings are associated with each copper atom. This could be accomplished by further alkylation of the copper in XIV to form a mixed "cuprate", which then undergoes a metal-halogen exchange reaction with a second phosphazene ring, and subsequent alkylation of the remaining "tricoordinate" P-Cl unit to yield VI. Thus, of the three equivalents of Grignard reagent per phosphazene ring required for the complete reaction, one equivalent is used for metallation, one for the methylation, and up to one equivalent for the reaction with the alkyl halide to yield R-R.

The final step in the reaction pathway is the interaction of the metallo-phosphazene complex VI with a 2-propanol to yield II. The copper-phosphazene bond may be cleaved to generate an N-H unit which then rearranges rapidly to form the P-H tautomer, II. Attempts to detect the presence of the N-H form, by the use of NMR spectroscopy, were unsuccessful, and it appears that cleavage of the copper-nitrogen bond could be a slower step than rearrangement of the proton from nitrogen to phosphorus. The ability of a cyclophosphazene system to undergo N-H to P-H tautomerism is well-documented.^{30-32,54}

Aryl Grignard reagents did not participate in the same type of reaction, and no aryl-hydrido-cyclophosphazenes have yet been prepared by the organo-copper route. Phenylmagnesium chloride reacted with $(\text{NPCl}_2)_3$ in the presence of $[\text{n-Bu}_3\text{PCuI}]_4$ to yield $\text{gem-N}_3\text{P}_3\text{Cl}_4\text{PH}_2$ and ring-cleavage products.^{19,53} Thus, in these reactions, phenylcopper appears to react in a way that is reminiscent of

alkyl Grignard reagents.⁵³ The differences in reactivity between alkyl- and aryl-copper reagents have been noted previously.²⁰

Compounds of structure II or VI are valuable not only for their mechanistic and structural features, but also because of their use as general synthetic intermediates. These aspects will be discussed in a later paper.

Experimental Section

Materials. Hexachlorocyclotriphosphazene (I) was supplied by Ethyl Corporation and was purified by sublimation, followed by 2 recrystallizations from *n*-hexane. The Grignard reagents were commercial products obtained from Aldrich or Alfa-Ventron. Tetrahydrofuran was distilled into the reaction flask under an atmosphere of dry nitrogen from a sodium-benzophenone ketal drying agent. The reagent, $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$, was prepared by standard methods³⁴ and was recrystallized from 2-propanol/ethanol before use. All reactions were carried out under an atmosphere of dry nitrogen.

General Synthetic Route to Hydrido-Phosphazenes. The syntheses of all the hydrido-phosphazenes were carried out in an identical manner. The following procedure is typical. Hexachlorocyclotriphosphazene (I) (5.0 g, 0.014 mol) and $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ (3.0 g, 0.019 mol) were stirred together in tetrahydrofuran (150 ml) at -80°C , and the Grignard reagent (10 ml of a 3 M solution in THF or ether) was added dropwise over a period of ≈ 30 min. The temperature was allowed to rise to $\approx -30^\circ\text{C}$, and a further 10 ml of the Grignard reagent was added. The reaction mixture was then stirred for 16 h and the temperature was allowed to rise to 25°C . The mixture was then cooled to 0°C and 2-propanol (10 ml) was added. Finally, removal of the solvent in vacuo and sublimation (or distillation) of the

residues gave the hydrido-phosphazenes (II) in up to 76% yield. NMR data for all the compounds are summarized in Table I. Yields, analytical data, mass spectral and infrared data are summarized in Table II.

Initial Monitoring of the Reaction. Hexachlorocyclotriphosphazene (5.0 g, 0.014 mol) and $[\underline{n}\text{-BuPCuI}]_4$ (3.0 g, 0.0019 mol) were stirred together in tetrahydrofuran (150 ml) at 0°C. An aliquot (5 μL) was withdrawn and was subjected to vapor phase chromatographic analysis. Methylmagnesium chloride (2 ml, 3.0 M solution of THF) was added and the reaction was stirred for 30 min. Stirring was discontinued and any solid matter was allowed to settle to the bottom of the flask. An aliquot was again withdrawn (5 μL) and was subjected to vapor phase chromatographic analysis. The above sequence was repeated until 20 ml of the Grignard reagent had been added to the reaction mixture. At this time the reaction mixture was stirred for a further 48 h, with samples being withdrawn every 12 h for analysis. Finally, $(\text{CH}_3)_2\text{CHOH}$ (10 ml) was added and, after stirring for 30 min, the final aliquot (5 μL) was removed for vapor phase chromatographic analysis.

Identification of the Source of Hydrogen. A reaction was carried out as described in the general synthesis section. However, $(\text{CH}_3)_2\text{CHOD}$ (10 ml) was used in the final step. The isolation and purification steps were the same as described previously. The compound, $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{D}$ (III) was obtained as white, air- and moisture-sensitive needles in 70% yield.

Variation in Copper Concentration. A series of reactions was carried out in which the amount of phosphazene and added Grignard reagent were maintained constant, but the quantity of copper in the reaction mixture was varied. The amount of $(\text{NPCl}_2)_3$ used in each case was 5.0 g, with 20 ml of a 3 M solution of

methylmagnesium chloride in THF. The reactions were carried out as described previously. The quantity of $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ used in each reaction is summarized in Table III. The results are shown graphically in Figure 5.⁵⁵

Attempted Reaction Between $(\text{NPCl}_2)_3$ and $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$. Hexachlorocyclotriphosphazene (5.0 g, 0.014 mol) was dissolved in THF (100 ml) at room temperature. A sample of this solution was removed and its ^{31}P NMR spectrum was obtained.³⁶ The spectrum consisted of a singlet at 19.8 ppm which was characteristic of unchanged $(\text{NPCl}_2)_3$.³⁸ A sample of $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ (3.0 g, 0.0019 mol) was also dissolved in THF (100 ml) and its ^{31}P NMR spectrum (broad band proton decoupled) was scanned.³⁶ The spectrum showed a broad singlet centered at -34 ppm.³⁸ These two solutions were mixed and the ^{31}P NMR spectrum of the resultant solution showed only peaks from $(\text{NPCl}_2)_3$ and $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$. No peak shift or other change occurred in the spectrum after 48 h of stirring. Thus, it was concluded that no reaction had occurred.

^{31}P NMR Monitoring of the Complete Reaction. To the mixture obtained from the above experiment was added methylmagnesium chloride in 2.5 ml aliquots (3.0 M solution in THF). After each addition a sample (3 ml) was removed and its ^{31}P NMR spectrum was scanned (Figure 6a). 2-Propanol (1.0 ml) was then added to the mixture in the NMR tube and the spectrum was scanned again (Figure 6b).

Reaction of $(\text{NPCl}_2)_3$ with Methylcopper. The reagent, $[\underline{n}\text{-Bu}_3\text{PCuI}]_4$ (11 g, 0.0069 mol), was dissolved in THF (150 ml) and was cooled to -80°C . Methylmagnesium chloride (10 ml, 3.0 M solution in THF) was then added dropwise. The rapid formation of methylcopper was evident from the deposition of a heavy

yellow precipitate. After the mixture had been stirred for 30 min, $(\text{NPCl}_2)_3$ (5.0 g, 0.014 mol), dissolved in THF (100 ml), was added rapidly to the suspension of methylcopper at -80°C . The reaction was stirred for a further 16 h, during which time the temperature was allowed to rise to 25°C . Finally, $(\text{CH}_3)_2\text{CHOH}$ (10 ml) was added to the mixture, which was then stirred for a further 30 min. Removal of the solvent under vacuum and sublimation of the residue gave a mixture of products. This was analyzed by gas chromatography/mass spectrometry and was found to consist of $(\text{NPCl}_2)_3$, $\text{N}_3\text{P}_3\text{Cl}_5\text{H}$, and $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$.

Isolation of Gaseous Side-Products. A reaction was carried out between $(\text{NPCl}_2)_3$, methylmagnesium chloride, and $[\text{n-Bu}_3\text{PCuI}]_4$ at 0°C in THF with the molar ratios as described earlier (general synthetic route). However, the reaction was carried out under a nitrogen atmosphere, but in a closed reaction vessel connected via an oil "bubbler" to a water-filled gas collection vessel. After addition of the Grignard reagent, the reaction mixture was stirred for 30 min, after which time gas evolution had almost ceased. A total of approximately 150 ml of gas was collected. Samples of the gas were withdrawn from the collection vessel and were subjected to gas chromatography/mass spectrometric analysis. Chloromethane, ethane, and a trace of methane were detected. In two control experiments, (a) $(\text{NPCl}_2)_3$ and MeMgCl were allowed to react,⁵¹ and (b) Methylmagnesium chloride was added to a solution of $[\text{n-Bu}_3\text{PCuI}]_4$ at 0°C . In neither case was chloromethane detected.

Use of Phenylmagnesium Chloride. Hexachlorocyclotriphosphazene (5.0 g, 0.014 mol) and $[\text{n-Bu}_3\text{PCuI}]_4$ (3.0 g, 0.0019 mol) were dissolved in THF (150 ml) and the mixture was cooled to -80°C . Phenylmagnesium chloride (20 ml, 3.0 M solution in THF) was added slowly and the reaction mixture was stirred for 16 h,

during which time the temperature was allowed to rise to 25°C. Finally, $(\text{CH}_3)_2\text{CHOH}$ (10 ml) was added to the mixture. Removal of the solvent and sublimation of volatile products from the residue gave $\text{N}_3\text{P}_3\text{Cl}_4\text{Ph}_2$ (identified by mass spectrometry) in very low yield.

Acknowledgments. This work was supported by the Office of Naval Research through Grant No. N00014-75-C-0685.

References and Notes

1. For a previous paper in this series see H. R. Allcock, P. P. Greigger, J. E. Gardner, and J. L. Schmutz, J. Am. Chem. Soc., **101**, 606 (1979).
2. See also T. L. Evans, T. J. Fuller, and H. R. Allcock, J. Am. Chem. Soc., **101**, 242 (1979).
3. A preliminary communication on this work appeared in J. Am. Chem. Soc., **100**, 6512 (1978).
4. R. J. Ritchie, P. J. Harris, and H. R. Allcock, Macromolecules, submitted.
5. H. R. Allcock and D. B. Patterson, Inorg. Chem., **16**, 197 (1977).
6. H. R. Calhoun, R. H. Lindstrom, R. T. Oakley, and N. L. Paddock, J. Chem. Soc., Chem. Commun., 343 (1975).
7. H. R. Calhoun, R. T. Oakley, N. L. Paddock, and J. Trotter, Can. J. Chem., **53**, 2413 (1975).
8. R. T. Oakley and N. L. Paddock, Can. J. Chem., **53**, 3038 (1975).
9. F. A. Cotton and A. Shaver, Inorg. Chem., **10**, 2362 (1971).
10. N. L. Paddock, T. N. Ranganathan, and S. M. Todd, Can. J. Chem., **49**, 164 (1971).
11. M. Biddlestone and R. A. Shaw, Phosphorus, **3**, 95 (1973).
12. R. A. Shaw, B. W. Fitzsimmons, and B. C. Smith, Chem. Rev., **62**, 247 (1962).
13. C. W. Allen, I. C. Paul, and T. Moeller, J. Am. Chem. Soc., **89**, 6361 (1967).
14. T. Moeller and F. Y. Tsang, Chem. & Ind., 361 (1962).
15. C. W. Allen and T. Moeller, Inorg. Chem., **7**, 2177 (1968).
16. C. W. Allen, F. Y. Tsang, and T. Moeller, Inorg. Chem., **7**, 2183 (1968).
17. T. N. Ranganathan, S. M. Todd, and N. L. Paddock, Inorg. Chem., **12**, 316 (1973).
18. C. W. Allen, J. Chem. Soc., Chem. Commun., 152 (1970).
19. H. R. Allcock and C. T.-W. Chu, Macromolecules, submitted.
20. G. H. Posner, Org. Synth., **22**, 253 (1975).

21. A. E. Jukes, Adv. Organomet. Chem., 12, 215 (1974).
22. J. F. Normant, Synthesis, 63 (1972).
23. G. H. Posner, Org. Synth., 19, 1 (1972).
24. G. Bahr and P. Burba, Methoden der Organischen Chemie, 4th ed., Vol. 13, Part 4, (E. Muller, Ed.), Georg Thieme Verlag, Stuttgart, 1970, p. 735.
25. M. Nilsson, Svensk Kem. Tidskr., 80, 192 (1968) (C.A. 69, 87034k, (1968)).
26. K. Wada and H. Hashimoto, Senryo to Yahukin, 12, 431 (1967) (C.A. 69, 87036n (1968)).
27. W. Carruthers, Chem. & Ind., 931 (1973).
28. I. Kuwajima, Yuki Gosei Kagaku Shi, 29, 616 (1971) (C.A. 75, 110354z (1971)); J. Synth. Org. Chem. Jap., 29, 616 (1971).
29. T. Kauffmann, Angew. Chem., Int. Ed. Engl., 13, 291 (1974).
30. A. Schmidpeter and H. Rossknecht, Chem. Ber., 107, 3146 (1974).
31. A. Schmidpeter and J. Ebeling, Angew. Chem., 80, 197 (1968), Angew. Chem. Int. Ed. Engl., 7, 209 (1968).
32. A. Schmidpeter, J. Hogel, and F. R. Ahmed, Chem. Ber., 109, 1911 (1976).
33. A. Schmidpeter, K. Blank, H. Eiletz, H. Smetana, and C. Weingand, Synth. React. Inorg. Metal-Org. Chem., 7, 1 (1977).
34. G. B. Kauffman and L. A. Teter, Inorg. Synth., 7, 9 (1963).
35. We thank Mr. Alan Freyer for his assistance in obtaining all the NMR spectra.
36. ³¹P NMR data were recorded on a JEOL JNM-PS-100 spectrometer operating at 40 MHz in the Fourier Transform mode. The data were processed on a Nicolet 1080 computer.
37. F. A. Bovey, "Nuclear Magnetic Resonance Spectroscopy", Academic Press, New York, 1969.
38. Positive ³¹P NMR chemical shifts are downfield from external phosphoric acid.

39. The flip angle employed in the F.T. data collection was 45° , and the pulse repetition rate was 4 sec. Measurements of the T1 spin-lattice relaxation time for the two types of phosphorus atoms in the compound by the 180° - τ - 90° method⁴⁰ gave a value of the T1 for the P(HCH₃) phosphorus atom as 4.19 ± 0.04 sec, and for the P(Cl₂) phosphorus atoms the value was 17.00 ± 0.09 sec. From this data it is clear that the P(Cl₂) phosphorus atoms will saturate much before the P(HCH₃) phosphorus atom under the pulse program employed in the ³¹P data collection, thus leading to an enhancement of the P(HCH₃) resonance signal.
40. R. L. Vold, J. S. Waugh, M. P. Klein, and D. E. Phelps, J. Chem. Phys., **48**, 3831 (1968).
41. ¹H NMR spectra were recorded on a JEOL JNM-PS-100 spectrometer operating at 100 MHz in the Fourier Transform mode. The data were processed on a Nicolet 1080 computer.
42. All ¹H chemical shifts are downfield from internal TMS and are quoted on the σ scale. For TMS $\sigma = 0$.
43. This coupling constant was given as 3-4 Hz in an earlier communication.³ However, a more accurate value for this coupling constant is now available due to our recent acquisition of ³¹P decoupling circuits on the ¹H NMR probe.
44. Infrared spectra were recorded on a Perkin-Elmer 580 infrared spectrometer and were run on KBr discs.
45. Mass spectra were recorded on an AEI MS-902 mass spectrometer operating at 70 eV ionizing potential.
46. Vapor phase chromatographic data were obtained using a Hewlett-Packard Model 5750 chromatographic instrument fitted with a 6' 10% OV17 column and a thermal conductivity detector. The injection port was at 250° , the helium carrier gas flow rate was 2.51 min^{-1} , and the oven temperature range was 50° to 250° at $20^\circ \text{ min}^{-1}$, following a 3 min post injection interval.

47. R. M. Silverstein, G. C. Bassler, and T. C. Morill, "Spectrometric Identification of Organic Compounds", 3rd. Ed., J. S. Wiley & Son, New York, 1974, p. 36.
48. ^2D NMR spectra were recorded on a JEOL JNM PS-100 spectrometer operating at 15 MHz in the Fourier Transform mode. The data were processed on a Nicolet 1080 computer.
49. A. A. Borisenko, N. M. Sergeyev, and Yu. A. Ustynyuk, Mol. Phys., **22**, 715 (1971).
50. As determined by gas chromatographic-mass spectral data, obtained with the use of a Finnigan 3200 instrument.
51. H. R. Allcock and R. J. Ritchie, to be published.
52. M. Biddlestone and R. A. Shaw, J. Chem. Soc. A, 178 (1969), *ibid*, 1760 (1970), *ibid*, 2715 (1971).
53. E. T. McBee, K. Okuhara, and C. J. Morton, Inorg. Chem., **4**, 1672 (1965).
54. A. Schmidpeter and H. Rossknecht, Angew. Chem., Int. Ed. Engl., **8**, 614 (1969).
55. At very low copper concentrations the main products of the reaction were shown by gas chromatography/mass spectrometry to consist of $\text{N}_3\text{P}_3\text{Cl}_5\text{CH}_3$, IV, and $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)_2$, V.

Figure 1. ^{31}P NMR spectrum of $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$ (proton decoupled).

Figure 2. Proton-coupled ^{31}P NMR spectrum of $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$.

Figure 3. ^1H NMR spectrum of $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$ (phosphorus decoupled).

Figure 4. Phosphorus-coupled ^1H NMR spectrum of $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$.

Figure 5. Curve showing the relationship between the percentage yield of $\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)\text{H}$ and the percentage of copper atoms to $(\text{NPCl}_2)_3$ molecules in the reaction mixture. For the circled points, see ref. 55.

Figure 6. Changes in the ^{31}P NMR spectrum following the addition of increasing amounts of CH_3MgCl to a 1:8 ratio mixture of $[\text{n-Bu}_3\text{PCuI}]_4$ and $(\text{NPCl}_2)_3$.
(a) Spectra of the reaction mixture before the addition of 2-propanol.
(b) Spectra after the addition of 2-propanol.

Table I

Hydridophosphazenes - NMR Data^{a,35}

Compound	³¹ P NMR ³⁶ ppm scale ³⁸		¹ H NMR ⁴¹ δ scale ⁴²		Alkyl	Coupling Constants ^b	
	P(R)H	PCl ₂	Hydride	-CH ₃		J _{PH}	J _{PCH}
N ₃ P ₃ Cl ₄ (CH ₄)H ^{c,d}	13.8 (t)	17.6 (d)	7.44 (d,t,m)	-CH ₃	1.75 (d,m)	= 568 Hz	
							= 16 Hz
							= 12 Hz
							= 11 Hz
							= 4 Hz
							= 2.9 Hz
N ₃ P ₃ Cl ₄ (C ₂ H ₅)H ^{c,d}	20.4 (t)	18.4 (d)	7.33 (d,t)	-CH ₂ -CH ₃	1.90 (m)	= 553 Hz	
							= 12 Hz
							= 8 Hz
							= 23 Hz
							= 7.5 Hz
							- unresolved
N ₃ P ₃ Cl ₄ (n-C ₃ H ₇)H ^{d,e}	17.1 (t)	19.0 (d)	7.33 (d)	-CH ₂ -CH ₂ -CH ₃	1.75 (m)	= 554 Hz	
				-CH ₂ CH ₂ CH ₃	1.09 (t)	= 12 Hz	
						= 8 Hz	
							other couplings unresolved

Table I (continued)

$\text{N}_3\text{P}_3\text{Cl}_4(\underline{1}\text{-C}_3\text{H}_7)\text{H}^{\text{c,d}}$	26.5 (t)	18.4 (d)	7.10 (d,t)	$-\underline{\text{CH}}(\underline{\text{CH}_3})_2$	1.98 (m)	$J_{\text{PH}} = 548 \text{ Hz}$
						$J_{\text{PNPH}} = 13 \text{ Hz}$
				$-\underline{\text{CH}}(\underline{\text{CH}_3})_2$	1.20 (d,d)	$J_{\text{PNP}} = 8 \text{ Hz}$
						$J_{\text{PCH}} = 7.5 \text{ Hz}$
						$J_{\text{PCCH}} = 23 \text{ Hz}$
						$J_{\text{HCCH}} = 7.0 \text{ Hz}$
$\text{N}_3\text{P}_3\text{Cl}_4(\underline{\text{m}}\text{-C}_4\text{H}_9)\text{H}^{\text{d,e}}$	17.9 (t)	19.4 (d)	7.33 (d,t)	$-(\underline{\text{CH}_2})_3\text{CH}_3$	0.96 (m)	$J_{\text{PH}} = 554 \text{ Hz}$
						$J_{\text{PNPH}} = 12 \text{ Hz}$
				$-(\underline{\text{CH}_2})_3\text{CH}_3$	1.60 (t)	$J_{\text{PNP}} = 4 \text{ Hz}$
						$J_{\text{HCCH}} = 18 \text{ Hz}$
						other couplings unresolved
$\text{N}_3\text{P}_3\text{Cl}_4(\underline{1}\text{-C}_4\text{H}_9)\text{H}^{\text{d,e}}$	15.7 (t)	19.0 (d)	7.36 (d,t)	$\underline{\text{CH}}_2\text{CH}(\underline{\text{CH}_3})_2$	2.40 (m)	$J_{\text{PH}} = 553 \text{ Hz}$
						$J_{\text{PNPH}} = 12 \text{ Hz}$
				$\text{CH}_2\text{CH}(\underline{\text{CH}_3})_2$	1.01 (d)	$J_{\text{PNP}} = 6 \text{ Hz}$
						$J_{\text{HCCH}} = 20 \text{ Hz}$
						other couplings unresolved

Table I (continued)

$N_3P_3Cl_4(t-C_4H_9)^{c,d}$	32.3 (t)	18.4 (d)	6.83 (d,t)	$-C(CH_3)_3$	1.17 (d)	$J_{PH} = 543 \text{ Hz}$
						$J_{PNPH} = 13 \text{ Hz}$
						J_{PNP} unresolved
						$J_{PCCH} = 20 \text{ Hz}$

a d = doublet, t = triplet, m = multiplet.

b Confirmed by homo- and hetero-nuclear decoupling experiments.

c ^{31}P NMR spectrum run in THF solution.

d 1H NMR spectrum run in $CDCl_3$ solution.

e ^{31}P NMR spectrum run in $CDCl_3$ solution

Table II

Compound	% Yield	M. Pt.	Infrared Data ^{44,a}		Mass Spectral Data ⁴⁵		Elemental Analysis	
			ν_{PH}	ν_{PN}	Found ^b	Calculated ^c	Found	Calculated
$\text{N}_3\text{P}_3\text{Cl}_4(\text{CH}_3)_4\text{H}$	76%	95°	2409 (m) 2399 (m)	1220 (s) 1180 (s) 1160 (s)	291	291	C 3.98 H 1.40 N 14.23 P 31.65 Cl 47.45	C 4.13 H 1.37 N 14.33 P 31.74 Cl 48.46
$\text{N}_3\text{P}_3\text{Cl}_4(\text{C}_2\text{H}_5)_5\text{H}$	60%	59°	2399 (m) 2395 (m)	1228 (s) 1185 (s) 1170 (sh)	305	305	C 7.82 H 2.00 N 13.71 P 30.39 Cl 46.17	C 7.82 H 1.95 N 13.68 P 30.29 Cl 46.25
$\text{N}_3\text{P}_3\text{Cl}_4(\text{n-C}_3\text{H}_7)_4\text{H}$	68%	62°	2418 (m) 2399 (m)	1215 (s) 1178 (s) 1160 (s)	319	319	C 11.43 H 2.57 N 12.99 P 28.68 Cl 44.32	C 11.21 H 2.49 N 13.08 P 28.93 Cl 44.24
$\text{N}_3\text{P}_3\text{Cl}_4(\text{i-C}_3\text{H}_7)_4\text{H}$	69%	65°	2395 (m) 2380 (m)	1210 (s) 1185 (s) 1170 (s)	319	319	C 11.34 H 2.50 N 13.08 P 28.75 Cl 44.10	C 11.21 H 2.49 N 13.08 P 28.93 Cl 44.24

Table II (continued)

$\text{M}_3\text{P}\text{Cl}_4(\text{m-C}_4\text{H}_9)\text{H}$	58%	43°	2420 (m)	1210 (s)	333	333	C	14.22	C	14.33
			2405 (m)	1185 (s)			H	3.00	H	2.98
				1160 (sh)			N	12.57	N	12.54
							P	27.87	P	27.76
							Cl	41.94	Cl	42.39

$\text{M}_3\text{P}\text{Cl}_4(\text{i-C}_4\text{H}_9)\text{H}^{\text{d}}$	36%	011	2401 (m)	1225 (s)	333	333			
			2395 (m)	1180 (s)					
				1165 (sh)					

$\text{M}_3\text{P}\text{Cl}_4(\text{t-C}_4\text{H}_9)\text{H}^{\text{d}}$	55%	39°	2375 (m)	1245 (s)	333	333			
			2365 (m)	1200 (s)					
				1185 (s)					

^a w = weak, m = medium, s = strong, br = broad, sh = shoulder.

^b As the base peak in a Cl_4 isotope pattern.

^c Calculated using $^{35}\text{Cl}_4$.



^d Elemental analysis could not be obtained on this compound due to its low thermal stability.

Table III



<u>[n-Bu₃PCuI]₄ in Reaction Mixture</u>		<u>Product Yield</u>	
<u>Wt.</u>	<u>% Cu/Phosphazene</u>	<u>Wt.</u>	<u>%^a</u>
0.28 g	5	0.50 g	12 ^b
0.56 g	10	0.63 g	15 ^b
0.84 g	15	0.95 g	23
1.12 g	20	1.25 g	30
1.40 g	25	1.60 g	38
1.68 g	30	1.93 g	46
1.96 g	35	2.40 g	58
2.25 g	40	2.55 g	61
2.52 g	45	2.93 g	70
2.80 g	50	3.06 g	74
3.08 g	55	3.04 g	74
4.20 g	75	3.12 g	76
5.50 g	100	3.15 g	76




^a Calculated for N₃P₃Cl₄(CH₃)H


^b See ref. 55




PPM










Equivalents of
Grignard/phosphazene
added.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz	1	U.S. Army Research Office P.O. Box 1211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Weapons Center China Lake, California 93555 Attn: Dr. A. B. Amster Chemistry Division	1
ONR Area Office One Hallidie Plaza, Suite 601 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Naval Civil Engineering Laboratory Port Hueneme, California 93401 Attn: Dr. R. W. Drisko	1
ONR Branch Office Building 114, Section D 666 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Professor K. E. Woehler Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Naval Ship Research and Development Center Annapolis, Maryland 21401 Attn: Dr. G. Bosmajian Applied Chemistry Division	1
		Naval Ocean Systems Center San Diego, California 91232 Attn: Dr. S. Yamamoto, Marine Sciences Division	1

TECHNICAL REPORT DISTRIBUTION LIST, 356B

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. T. C. Williams Union Carbide Corporation Chemical and Plastics Tarrytown Technical Center Tarrytown, New York	1	Douglas Aircraft Company 3855 Lakewood Boulevard Long Beach, California 90846 Attn: Technical Library CI 290/36-84 AUTO-Sutton	1
Dr. R. Soulen Contract Research Department Pennwalt Corporation 900 First Avenue King of Prussia, Pennsylvania 19406	1	NASA-Lewis Research Center 21000 Brookpark Road Cleveland, Ohio 44135 Attn: Dr. T. T. Serafini, MS 49-1	1
Dr. A. G. MacDiarmid University of Pennsylvania Department of Chemistry Philadelphia, Pennsylvania 19174	1	Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375	1
Dr. C. Pittman University of Alabama Department of Chemistry University, Alabama 35486	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. E. Allcock Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1	Dr. E. Fischer, Code 2853 Naval Ship Research and Development Center Annapolis Division Annapolis, Maryland 21402	1
Dr. M. Kenney Case-Western University Department of Chemistry Cleveland, Ohio 44106	1	Dr. Martin H. Kaufman, Head Materials Research Branch (Code 4542) Naval Weapons Center China Lake, California 93555	1
Dr. R. Lenz University of Massachusetts Department of Chemistry Amherst, Massachusetts 01002	1	Dr. J. Magill University of Pittsburg Metallurgical and Materials Engineering Pittsburg, Pennsylvania 22230	1
Dr. M. David Curtis University of Michigan Department of Chemistry Ann Arbor, Michigan 48105	1	Dr. C. Allen University of Vermont Department of Chemistry Burlington, Vermont 05401	1
Dr. M. Good University of New Orleans Department of Chemistry Lakefront New Orleans, Louisiana 70122	1	Dr. D. Bergbreiter Texas A&M University Department of Chemistry College Station, Texas 77843	1

472:GAN:710:ca:
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, 356B

No.
Copies

Professor R. Drago
Department of Chemistry
University of Illinois
Urbana, Illinois 61801

1

Dr. F. Brinkman
Chemical Stability & Corrosion
Division
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

1