

AD-A071 317

CENTER FOR NAVAL ANALYSES ALEXANDRIA VA
QUASI-RESIDUATED MAPPINGS AND BAER ASSEMBLIES. (U)
JUL 70 T S BLYTH, W C HARDY
CNA-PP-39

F/G 12/1

UNCLASSIFIED

NL

| OF |
AD
A071317



END
DATE
FILMED
8-79
DDC

DA071317

LEVEL

(13.5)

(12) 31p

(6) QUASI-RESIDUATED MAPPINGS AND BAER ASSEMBLIES

(10) by T. S. Blyth and W. C. Hardy

(14) CNA-PP-39

DDC RECEIVED JUL 18 1979 A

(9) Professional Paper No. 39

CENTER FOR NAVAL ANALYSES

(11) 74 July 1970

DDC FILE COPY.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

Accession for	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By <i>Acq. Unit 1379</i>	
Distribution/ <i>0652</i>	
Availability Codes	
Dist.	Avail and/or special
<i>A</i>	

077 270 *See*

79 07 16 082

The ideas expressed in this paper are those of the authors. The paper does not necessarily represent the views of either the Center for Naval Analyses, the United States Navy or any other sponsoring agency. It has been reproduced by CNA as a courtesy to the authors, under the CNA Professional Development Program.

Quasi-residuated Mappings and Baer Assemblies. By T.S. Blyth, Mathematical Institute, University of St. Andrews and W.C. Hardy, Centre for Naval Analyses, Arlington, Virginia.

SYNOPSIS

We consider, for a given ordered set E with minimum element 0 , the semigroup Q of 0 -preserving isotone mappings on E and examine necessary and sufficient conditions under which an element $f \in Q$ is such that the left [resp. right] annihilator of f in Q is a principal left [resp. right] ideal of Q generated by a particular type of idempotent. The results obtained lead us to introduce the concept of a Baer assembly which we use to extend to the case of a semilattice the Baer semigroup coördinatisation theory of lattices. We also derive a coördinatisation of particular types of semilattice.

in which point f belongs to set Q

1. Introduction.

In the publications [1] and [2] a profound connection is established between the theory of lattices and the theory of semigroups. We begin by describing a slight generalisation of this, a detailed account of which will be found in [3]. Let S be a semigroup and let K be a two-sided ideal of S . For each $x \in S$ define the left and right K -annihilator of x by

$$L_K(x) = \{y \in S; yx \in K\}, \quad R_K(x) = \{y \in S; xy \in K\}.$$

We say that the pair $\langle S; K \rangle$ forms a Baer semigroup if and only if, for each $x \in S$, $L_K(x)$ is a principal idempotent-generated left ideal of S and $R_K(x)$ is a principal idempotent-generated right ideal of S . If $\langle S; K \rangle$ is a Baer semigroup then the sets $\mathcal{L}(S) = \{L_K(x); x \in S\}$ and $\mathcal{R}(S) = \{R_K(x); x \in S\}$ form dually isomorphic

lattices. To proceed in the opposite direction, let E and F be bounded ordered sets; *i.e.*, ordered sets which have a minimum element and a maximum element. A mapping $f : E \rightarrow F$ is said to be *residuated* if and only if it is isotone (order-preserving) and such that there exists a (necessarily unique) mapping $f^+ : F \rightarrow E$ which is isotone and such that $f^+ \circ f \geq \text{id}_E$ and $f \circ f^+ \leq \text{id}_F$. When such a mapping f^+ exists, it is given by the prescription

$$f^+(y) = \max\{x \in E; f(x) \leq y\}.$$

It can be shown that $f : E \rightarrow F$ is residuated if and only if the pre-image under f of each principal order ideal of F (*i.e.*, a subset of the form $[+, x] = \{y \in F; y \leq x\}$) is a principal order ideal of E . The set $\text{Res}(E)$ of residuated mappings from E to itself forms a semigroup under composition of mappings and an investigation into just when $\langle \text{Res}(E); \{0\} \rangle$ forms a Baer semigroup yields the fundamental result: *if E is a bounded ordered set then the following are equivalent:*

- (1) E is a lattice;
- (2) $\langle \text{Res}(E); \{0\} \rangle$ is a Baer semigroup;
- (3) there is a Baer semigroup $\langle S; K \rangle$ such that $E = \mathcal{R}(S)$.

When situation (3) holds we say that $\langle S; K \rangle$ *coördinatises* E .

In this paper we shall extend this result to the case of a *semilattice*. This we shall accomplish by examining annihilators in a much larger semigroup, namely the semigroup of isotone mappings which admit 0 as a fixed point. We obtain our coördination theorem using the concept of a Baer assembly and we apply

this to obtain coordinatisations of particular semilattices.

2. Quasi-residuated mappings.

Let E and F be ordered sets. We shall say that a mapping $f : E \rightarrow F$ is *quasi-residuated* if and only if it is isotone and such that there exists a mapping $h : F \rightarrow E$ such that $f \circ h \leq \text{id}_E$. It can be shown that f is quasi-residuated if and only if it is isotone and such that $(\forall y \in F) \{x \in E; f(x) \leq y\} \neq \emptyset$. In the case where E has a minimum element 0 it is easily seen that $f : E \rightarrow E$ is quasi-residuated if and only if it is isotone and such that $f(0) = 0$.

In what follows we shall assume that E has a minimum element 0 and we shall denote by Q the set of quasi-residuated mappings on E . It is clear that Q forms a semigroup under composition of mappings and that this semigroup admits a zero element, namely the zero mapping on E which we shall also denote by 0 without confusion. For each $f \in Q$ we shall write

$$L(f) = \{g \in Q; g \circ f = 0\}, \quad R(f) = \{g \in Q; f \circ g = 0\}.$$

Clearly $L(f)$ is a left ideal of Q and $R(f)$ is a right ideal of Q . Let us now examine under what conditions these annihilators are idempotent-generated principal ideals.

Definitions. We shall say that $f \in Q$ has a *principal kernel* if and only if $\{x \in E; f(x) = 0\}$ admits a maximum element which we shall denote by $f^+(0)$; and that $f \in Q$ has a *bounded image* if and only if $\{f(x); x \in E\}$ admits a maximum element which we shall denote

by $f(\pi)$. [Note that if E also has a maximum element π then by the isotone property every $f \in Q$ has a bounded image.]

THEOREM 1 Given $f \in Q$, the following conditions are equivalent:

- (1) f has a principal kernel;
- (2) $R(f)$ is a principal right ideal generated by an idempotent which has a bounded image.

Proof. (1) \Rightarrow (2) : Suppose first that f has a principal kernel. For each $y \in E$ define a mapping $\theta_y : E \rightarrow E$ by

$$\theta_y(x) = \begin{cases} x & \text{if } x \leq y; \\ y & \text{if } x \not\leq y. \end{cases}$$

Clearly each θ_y is isotone and idempotent. Moreover, each $\theta_y \in Q$ and has a bounded image. Now by the definition of $f^+(0)$ we have

$$(f \circ \theta_{f^+(0)})(x) = \begin{cases} f(x) = 0 & \text{if } x \leq f^+(0); \\ f[f^+(0)] = 0 & \text{if } x \not\leq f^+(0), \end{cases}$$

so that $f \circ \theta_{f^+(0)}$ is the zero map. Hence $\theta_{f^+(0)} \in R(f)$ and consequently $\theta_{f^+(0)} \circ Q \subseteq R(f)$. On the other hand, if $g \in Q$ then

$$\begin{aligned} g \in R(f) &\Rightarrow (\forall x \in E) f[g(x)] = 0 \\ &\Rightarrow (\forall x \in E) g(x) \leq f^+(0) \\ &\Rightarrow g = \theta_{f^+(0)} \circ g \\ &\Rightarrow g \in \theta_{f^+(0)} \circ Q \end{aligned}$$

and so we deduce that $R(f) = \theta_{f^+(0)} \circ Q$ which gives (2).

(2) \Rightarrow (1) : If now (2) holds, say $R(f) = \zeta \circ Q$ where $\zeta = \zeta \circ \zeta$, let $\zeta(\pi)$ denote the greatest element of $\text{Im } \zeta$. Since, for $x \in E$,

$$(f \circ \theta_x)(y) = f[\theta_x(y)] = \begin{cases} f(y) & \text{if } y \leq x; \\ f(x) & \text{if } y \not\leq x, \end{cases}$$

we see, using the fact that f is isotone, that

$$\theta_x \in R(f) \iff f(x) = 0.$$

we thus have

$$\begin{aligned} f(x) = 0 &\Rightarrow \theta_x \in R(f) = \zeta \circ Q \\ &\Rightarrow \theta_x = \zeta \circ \theta_x \\ &\Rightarrow x = \theta_x(x) = \zeta[\theta_x(x)] \leq \zeta(\pi). \end{aligned}$$

Since clearly $\zeta \in R(f)$ we have $f[\zeta(\pi)] = 0$. It thus follows that f has a principal kernel with $f^+(0) = \zeta(\pi)$.

THEOREM 2 Given $f \in Q$, the following conditions are equivalent:

- (1) f has a bounded image;
- (2) $L(f)$ is a principal left ideal generated by an idempotent which has a principal kernel.

Proof. (1) \Rightarrow (2) : Suppose that f has a bounded image. For each $y \in E$ define a mapping $\psi_y : E \rightarrow E$ by

$$\psi_y(x) = \begin{cases} 0 & \text{if } x \leq y; \\ x & \text{if } x \not\leq y. \end{cases}$$

Clearly each ψ_y is isotone and idempotent. Moreover, each $\psi_y \in Q$ and has a principal kernel. Since by definition $f(\pi)$ is the greatest element of $\text{Im } f$, we see easily that $\psi_{f(\pi)} \circ f$ is the zero map and so $\psi_{f(\pi)} \in L(f)$ whence $Q \circ \psi_{f(\pi)} \subseteq L(f)$. On the other hand, if $g \in Q$ then

$$\begin{aligned} g \in L(f) &\Rightarrow (\forall x \in E) \quad g[f(x)] = 0 \\ &\Rightarrow g[f(\pi)] = 0 \\ &\Rightarrow \{x \leq f(\pi) \Rightarrow g(x) = 0 = g(0)\} \\ &\Rightarrow g = g \circ \psi_{f(\pi)} \in Q \circ \psi_{f(\pi)} \end{aligned}$$

and so we deduce that $L(f) = Q \circ \psi_{f(\pi)}$ which gives (2).

(2) \Rightarrow (1) : Suppose that (2) holds, say $L(f) = Q \circ \xi$ where $\xi = \xi \circ \xi$ and $\xi^+(0)$ exists. Since $\xi \circ f$ is the zero map, we have

$$(*) \quad (\forall x \in E) \quad f(x) \leq \xi^+(0).$$

The property (1) will follow from this if we can show that $\xi^+(0)$ is an element of $\text{Im } f$. This we do as follows :

(a) suppose that f is the zero map : then clearly $Q = L(f) = Q \circ \xi$ from which it follows that there exists $h \in Q$ such that $\text{id}_E = h \circ \xi$. Hence we have $\xi(x) = 0 \Rightarrow x = 0$ and consequently $\xi^+(0) = 0 = f(0) \in \text{Im } f$.

(b) suppose that f is not the zero map : then (*) shows that $\xi^+(0) \neq 0$ so we can define a mapping $t : E \rightarrow E$ by setting

$$(**) \quad t(x) = \begin{cases} 0 & \text{if } (\exists z \in \text{Im } f) \ x \leq z ; \\ \xi^+(0) & \text{otherwise.} \end{cases}$$

To see that t is isotone, let $x \leq y$; then

(i) if $(\exists z \in \text{Im } f) \ x \leq z$ and $(\exists z^* \in \text{Im } f) \ y \leq z^*$ then we have $t(x) = 0 = t(y)$;

(ii) if $(\exists z \in \text{Im } f) \ x \leq z$ and $(\forall z^* \in \text{Im } f) \ y \not\leq z^*$ then we have $t(x) = 0 < \xi^+(0) = t(y)$;

(iii) the case where $(\forall z \in \text{Im } f) \ x \not\leq z$ and $(\exists z^* \in \text{Im } f) \ y \leq z^*$ cannot arise since by hypothesis $x \leq y$;

(iv) if $(\forall z \in \text{Im } f) \ x \not\leq z$ and $(\forall z^* \in \text{Im } f) \ y \not\leq z^*$ then we have $t(x) = \xi^+(0) = t(y)$.

Thus in all possible cases $t(x) \leq t(y)$ and so t is isotone. Moreover, $t(0) = 0$ and so $t \in Q$. Noting that if $x \in \text{Im } f$ then $t(x) = 0$, we see that $t \in L(f) = Q \circ \xi$ and so $t = t \circ \xi$. Thus

$$t[\xi^+(0)] = (t \circ \xi)[\xi^+(0)] = t(0) = 0.$$

It follows from (**) that there exists $x^* \in E$ with $\xi^+(0) \leq f(x^*)$ and from (*) that $f(x^*) \leq \xi^+(0)$. Hence $\xi^+(0) = f(x^*) \in \text{Im } f$.

THEOREM 3 Let $f \in Q$ have a principal kernel. Then for each idempotent $\zeta \in Q$ such that $R(f) = \zeta \circ Q$ the greatest element of $\text{Im } \zeta$ exists and is $f^+(0)$.

Proof. Consider the mapping $\theta_{f^+(0)}$ as defined in Theorem 1. Clearly $\theta_{f^+(0)} \in R(f)$ and so, if $R(f) = \zeta \circ Q$ with $\zeta = \zeta \circ \zeta$, we have $\theta_{f^+(0)} = \zeta \circ \theta_{f^+(0)}$. Consequently

$$f^+(0) = \theta_{f^+(0)}[f^+(0)] = (\zeta \circ \theta_{f^+(0)})[f^+(0)] = \zeta[f^+(0)].$$

Now $f \circ \zeta = 0$ and so $(\forall x \in E) \zeta(x) \leq f^+(0)$. It follows from this that $f^+(0)$ is the maximum element of $\text{Im } \zeta$.

THEOREM 4 Let $f \in Q$ have a bounded image. Then for each idempotent $\xi \in Q$ such that $L(f) = Q \circ \xi$ the element $\xi^+(0)$ exists and is $f(\pi)$.

Proof. Consider the mapping $\psi_{f(\pi)}$ as defined in Theorem 2. Clearly $\psi_{f(\pi)} \in L(f)$ and so, if $L(f) = Q \circ \xi$ where $\xi = \xi \circ \xi$, we have $\psi_{f(\pi)} = \psi_{f(\pi)} \circ \xi$. Consequently

$$\xi(x) = 0 \Rightarrow \psi_{f(\pi)}(x) = 0 \Rightarrow x \leq f(\pi).$$

But clearly if $x \leq f(\pi)$ then $\xi(x) \leq \xi[f(\pi)] = 0$. It therefore follows that $\xi^+(0)$ exists and is none other than $f(\pi)$.

The results of Theorems 3 and 4 give rise to the following two results which we shall be able to express in a useful equivalent way in the next section.

THEOREM 5 If E is an ordered set with minimum element 0 define

$$\mathcal{R} = \{R(f); f \in Q \text{ and } f \text{ has a principal kernel}\}$$

and order \mathcal{R} by set inclusion. Then the ordered sets E, \mathcal{R} are isomorphic.

Proof. By virtue of Theorem 3 we can define a mapping $\delta : \mathcal{R} \rightarrow E$ by the prescription $\delta[R(f)] = f^+(0)$ which can be written in the equivalent form $\delta[\zeta \circ Q] = \zeta(\pi)$ where ζ is an idempotent such that $R(f) = \zeta \circ Q$. Now $\zeta \circ Q \subseteq \eta \circ Q$ implies that $\zeta = \eta \circ \zeta$ which gives $\text{Im } \zeta \subseteq \text{Im } \eta$ and hence $\zeta(\pi) \leq \eta(\pi)$. On the other hand, if $\zeta(\pi) \leq \eta(\pi)$ and if $\eta \circ Q = R(g)$ where $g \in Q$ has a principal kernel then by Theorem 3 we have $\eta(\pi) = g^+(0)$ and consequently $\zeta(\pi) \leq g^+(0)$. It follows that $g \circ \zeta = 0$ so that $\zeta \in R(g) = \eta \circ Q$ and hence $\zeta \circ Q \subseteq \eta \circ Q$. We thus have

$$\delta(\zeta \circ Q) \leq \delta(\eta \circ Q) \iff \zeta(\pi) \leq \eta(\pi) \iff \zeta \circ Q \subseteq \eta \circ Q.$$

To show that δ is an order isomorphism, it therefore suffices to show that it is surjective. For this purpose, we note that for each $x \in E$ the mapping ψ_x as defined in Theorem 2 has a principal kernel and that $\psi_x^+(0) = x$. Thus $\delta[R(\psi_x)] = x$ and so δ is surjective.

THEOREM 6 If E is an ordered set with minimum element 0 define

$$\mathcal{L} = \{L(f); f \in Q \text{ and } f \text{ has a bounded image}\}$$

and order \mathcal{L} by set inclusion. Then the ordered sets E, \mathcal{L} are dually isomorphic.

Proof. By virtue of Theorem 4 we can define a mapping $\gamma : \mathcal{L} \rightarrow E$

by the prescription $\gamma[L(f)] = f(\pi)$ which can be written in the equivalent form $\gamma(Q \circ \xi) = \xi^+(0)$ where ξ is an idempotent such that $L(f) = Q \circ \xi$. Now $Q \circ \xi \subseteq Q \circ \eta$ implies that $\xi = \xi \circ \eta$ from which we deduce that $\eta(x) = 0 \Rightarrow \xi(x) = 0$ and so $\eta^+(0) \leq \xi^+(0)$. On the other hand, if $\eta^+(0) \leq \xi^+(0)$ and if $Q \circ \eta = L(g)$ where $g \in Q$ has a bounded image then by Theorem 4 we have $\eta^+(0) = g(\pi)$ and hence $\xi \circ g = 0$. Thus $\xi \in L(g) = Q \circ \eta$ and so $Q \circ \xi \subseteq Q \circ \eta$. We thus have

$$\gamma(Q \circ \xi) \leq \gamma(Q \circ \eta) \iff \xi^+(0) \leq \eta^+(0) \iff Q \circ \xi \subseteq Q \circ \eta.$$

To show that γ is a dual isomorphism it therefore suffices to show that it is surjective. For this purpose, we note that for each $x \in E$ the mapping θ_x as defined in Theorem 1 has a bounded image and that $\theta_x(\pi) = x$. Thus $\gamma[L(\theta_x)] = x$ and so γ is surjective.

3. Baer assemblies.

Definition. Let A be a semigroup, let B and C be distinguished subsets of A and let K be a two-sided ideal of A . We shall say that $\langle A; B, C; K \rangle$ forms a *Baer assembly* if and only if the following properties are satisfied :

$$\left\{ \begin{array}{l} (\forall x \in B) (\exists e = e^2 \in C) \quad L_K(x) = Ae ; \\ (\forall y \in C) (\exists f = f^2 \in B) \quad R_K(y) = fA . \end{array} \right.$$

By way of example, we mention that any Baer semigroup $\langle S; K \rangle$ may be regarded as a Baer assembly $\langle S; S, S; K \rangle$. Also, if E is an ordered set with a minimum element 0 , if Q denotes the semigroup of quasi-residuated mappings on E and if B, C are the subsets of Q con-

sisting respectively of those mappings which have bounded images, principal kernels then Theorems 1 and 2 show that $\langle Q; B, B; \{0\} \rangle$ is a Baer assembly.

We shall say that a Baer assembly $\langle A; B, C; K \rangle$ is *normal* if and only if $K \subseteq B \cap C$. Any Baer semigroup is clearly a normal Baer assembly. Likewise, if E is an ordered set with a minimum element and a maximum element and if Q, B, C are defined as above then we have $0 \in B \cap C$ and $B = Q$ so that $\langle Q; Q, C; \{0\} \rangle$ is a normal Baer assembly.

THEOREM 7 *If $\langle A; B, C; K \rangle$ is a normal Baer assembly then*

- (1) *A has an identity element which belongs to $B \cap C$;*
- (2) *K is a principal ideal generated by a central idempotent.*

Proof. (1) By hypothesis each $k \in K$ satisfies $k \in B \cap C$ and so there exist idempotents $e \in C, f \in B$ such that $A = L_K(k) = Ae$ and $A = R_K(k) = fA$. These equalities show that e is a right identity for A and that f is a left identity for A . It follows that $e = f$ and is a two-sided identity for A ; we denote this as usual by 1 . Since $e \in C$ and $f \in B$ we have $1 \in B \cap C$.

(2) Since $1 \in B \cap C$ there exist idempotents g, h such that $K = L_K(1) = Ag$ and $K = R_K(1) = hA$. These equalities show that g is a right identity for K and that h is a left identity for K . Hence $g = h$ is an identity for K which we shall denote by k° . We thus have $K = Ak^\circ = k^\circ A$. Now for each $x \in A$ we have $xk^\circ \in K$ and so $xk^\circ = k^\circ xk^\circ$. Likewise $k^\circ x \in K$ and so $k^\circ x = k^\circ xk^\circ$. It is immediate that k° is an idempotent which belongs to the centre of A .

In the next result we list further fundamental properties of normal Baer assemblies, by way of preparation for which we make the following remarks. If A is an arbitrary semigroup then it is well known that the power set $P(A)$, equipped with the multiplication defined by

$$\begin{cases} XY = \{xy; x \in X, y \in Y\}; \\ X\emptyset = \emptyset = \emptyset X, \end{cases}$$

is a residuated semigroup in which the residuals are given by

$$X \cdot Y = \{z \in A; (\forall y \in Y) zy \in X\}, \quad X \cdot Y = \{z \in A; (\forall y \in Y) yz \in X\}.$$

Suppose now that K is a non-empty subset of A and consider the mappings $\hat{L}_K, \hat{R}_K : P(A) \rightarrow P(A)$ given by the prescriptions

$$\hat{L}_K(X) = K \cdot X, \quad \hat{R}_K(X) = K \cdot X.$$

We have :

THEOREM 8 *If A is a semigroup and K is a non-empty subset of A then*

$$(1) \hat{L}_K \circ \hat{R}_K \circ \hat{L}_K = \hat{L}_K \text{ and } \hat{R}_K \circ \hat{L}_K \circ \hat{R}_K = \hat{R}_K.$$

If moreover $\langle A; B, C; K \rangle$ is a normal Baer assembly and if we define

$$\mathcal{L}(B) = \{L_K(x); x \in B\} \text{ and } \mathcal{R}(C) = \{R_K(x); x \in C\} \text{ then}$$

$$(2) \text{ if } e = e^2 \in B \text{ then } eA \in \mathcal{R}(C) \Leftrightarrow eA = (\hat{R}_K \circ \hat{L}_K)(eA) \text{ and if } f = f^2 \in C \text{ then } Af \in \mathcal{L}(B) \Leftrightarrow Af = (\hat{L}_K \circ \hat{R}_K)(Af);$$

(3) *the restriction \hat{L}_K^\rightarrow of \hat{L}_K to $\mathcal{R}(C)$ is a dual order isomorphism of $\mathcal{R}(C)$ onto $\mathcal{L}(B)$ whose inverse is \hat{R}_K^\rightarrow , the restriction of \hat{R}_K to $\mathcal{L}(B)$.*

Proof. (1) Since

$$\hat{R}_K(X) = K \cdot X \supseteq Y \Leftrightarrow XY \subseteq K \Leftrightarrow X \subseteq K \cdot Y = \hat{L}_K(Y),$$

we see that, when considered as a mapping from $P(A)$ to its dual, \hat{R}_K is residuated with residual given by $(\hat{R}_K)^+ = \hat{L}_K$. The equalities in (1) are then an immediate consequence of $\hat{L}_K \circ \hat{R}_K \geq \text{id}_{P(A)}$ and $\hat{R}_K \circ \hat{L}_K \geq \text{id}_{P(A)}$ [which express the fact that the pair (\hat{R}_K, \hat{L}_K) sets up a Galois connection on $P(A)$].

(2) Given $e = e^2 \in B$, suppose first that $eA \in \mathcal{R}(C)$; say $eA = R_K(x)$ where $x \in C$. Noting that $R_K(x) = \hat{R}_K\{x\}$ we have, by (1),

$$(\hat{R}_K \circ \hat{L}_K)(eA) = (\hat{R}_K \circ \hat{L}_K \circ \hat{R}_K)\{x\} = \hat{R}_K\{x\} = R_K(x) = eA.$$

Conversely, suppose that $e = e^2 \in B$ with $eA = (\hat{R}_K \circ \hat{L}_K)(eA)$. If we let $L_K(e) = Ae^*$ where e^* is an idempotent in C then, since A has an identity element by Theorem 7, $\hat{L}_K(eA) = L_K(e)$ and so

$$eA = (\hat{R}_K \circ \hat{L}_K)(eA) = \hat{R}_K[L_K(e)] = \hat{R}_K(Ae^*) = R_K(e^*) \in \mathcal{R}(C).$$

(3) It follows from the above that $R_K^{\rightarrow} \circ L_K^{\rightarrow} = \text{id}_{\mathcal{R}(C)}$ and that $L_K^{\rightarrow} \circ R_K^{\rightarrow} = \text{id}_{\mathcal{L}(B)}$. Since L_K^{\rightarrow} and R_K^{\rightarrow} are each antitone, the result follows.

Remark. In the proof of (2) above we used the fact that, since A has an identity element, $\hat{L}_K(eA) = L_K(e)$. In what follows we shall make use of this fact several times without reference.

Definition. If E is an ordered set then we shall say that E is *coordinatised* by a normal Baer assembly $\langle A; B, C; K \rangle$ if and only if $E = \mathcal{R}(C)$ or, equivalently [by Theorem 8(3)], $E \stackrel{d}{\cong} \mathcal{L}(B)$ where $\stackrel{d}{\cong}$ denotes dual order isomorphism.

With this terminology we see that the results of §2 can be succinctly summarised as follows: if E is an ordered set with minimum element 0, if Q denotes the semigroup of quasi-residuated

mappings on E and if B, C denote respectively the subsets of Q consisting of those mappings which have bounded images, principal kernels then $\langle Q; B, C; \{0\} \rangle$ is a normal Baer assembly which coördinatises E .

Having observed this fundamental result, we can now explicitly state the ultimate goal of the theory we are developing : to isolate those Baer assemblies which coördinatised ordered sets of a given type. In the following sections we begin the pursuit of this goal by restricting our attention to semilattices. During the course of this, the Baer semigroup coördinatisation of lattices will emerge in a rather nice way.

4. Coördinatisation of \wedge -semilattices.

THEOREM 9 *An ordered set E is an \wedge -semilattice with maximum element π if and only if, for each $x \in E$, $\theta_x \in \text{Res}(E)$. In this case the residual map θ_x^+ is given by*

$$\theta_x^+(y) = \begin{cases} \pi & \text{if } y \geq x; \\ y \wedge x & \text{if } y \not\geq x. \end{cases}$$

Proof. By definition, we have

$$\theta_x(y) = \begin{cases} y & \text{if } y \leq x; \\ x & \text{if } y \not\leq x. \end{cases}$$

Now if $x \leq z$ or $x \geq z$ then clearly $x \wedge z$ exists. Suppose then that $x \not\leq z$ and $x \not\geq z$. Given $y \in E$, we have

$$\theta_x(y) \leq z \Rightarrow \theta_x(y) \neq x \Rightarrow \theta_x(y) = y \Rightarrow \begin{cases} y \leq x; \\ y \leq z, \end{cases}$$

and conversely if $y \leq x, z$ then $\theta_x(y) = y \leq z$. Thus

$$\{y ; \theta_x(y) \leq z\} = [+ , x] \wedge [+ , z],$$

and so $x \wedge z$ exists if and only if $\max\{y ; \theta_x(y) \leq z\}$ exists. If we observe that $\text{Im } \theta_x = [+ , x]$, we see also that $\max\{y ; \theta_x(y) \leq x\}$ exists if and only if π exists. We conclude that E is an \wedge -semilattice with maximum element π if and only if each θ_x is residuated. Finally, it is readily verified that the mapping θ_x^+ as given above is isotone and such that $\theta_x^+ \circ \theta_x \geq \text{id}_E$ and $\theta_x \circ \theta_x^+ \leq \text{id}_E$, so that θ_x^+ is indeed the residual of θ_x .

This result, together with the following, which examines the close relationship between residuated mappings and mappings which have principal kernels, allows us to isolate a Baer assembly which allows a coördinatisation of \wedge -semilattices.

THEOREM 10 *Let E be an ordered set with minimum element 0 and let Q denote the semigroup of quasi-residuated mappings on E . Given $g \in Q$, the following conditions are equivalent :*

(1) *if $f \in Q$ has a principal kernel then $f \circ g$ has a principal kernel;*

(2) $g \in \text{Res}(E)$.

Proof. (1) \Rightarrow (2) : By hypothesis $g(0) = 0$ and so, given any $y \in E$, we have $\{x \in E ; g(x) \leq y\} \neq \emptyset$. Now the mapping ψ_y as defined in Theorem 2 has a principal kernel and so, applying (1), we obtain

$$g(x) \leq y = \psi_y^+(0) \Leftrightarrow (\psi_y \circ g)(x) = 0 \Leftrightarrow x \leq (\psi_y \circ g)^+(0).$$

It follows that g is residuated with g^+ given by the formula

$$g^+(y) = (\psi_y \circ g)^+(0).$$

(2) \Rightarrow (1) : If g is residuated and f has a principal kernel then we have

$$(f \circ g)(x) = 0 \Leftrightarrow g(x) \leq f^+(0) \Leftrightarrow x \leq g^+[f^+(0)],$$

from which it follows that $(f \circ g)^+(0)$ exists and is $g^+[f^+(0)]$.

Corollary If C denotes the subset of Q consisting of those mappings with principal kernels then, in $P(Q)$, $\text{Res}(E) = C \cdot C$.

Proof. Using the above, we have

$$g \in \text{Res}(E) \Leftrightarrow C \circ g \subseteq C \Leftrightarrow g \in C \cdot C.$$

Definition. By a right Baer assembly we shall mean a normal Baer assembly $\langle A; B, D; K \rangle$ in which

- (1) B is a subsemigroup of A ;
- (2) $B \subseteq D$;
- (3) in $P(A)$, $B \subseteq C \cdot C$ [i.e., $CB \subseteq C$].

THEOREM 11 Let E be an ordered set with minimum element 0 .

Let Q denote the semigroup of quasi-residuated mappings on E and let C be the subset of Q consisting of those mappings which have principal kernels. Then the following conditions are equivalent :

- (1) E is an \wedge -semilattice with a maximum element π ;
- (2) $\langle Q; \text{Res}(E), C; \{0\} \rangle$ is a right Baer assembly;
- (3) there is a right Baer assembly $\langle A; B, D; K \rangle$ with $E = \mathcal{R}(D)$.

Proof. (1) \Rightarrow (2) : Immediate from Theorems 9 and 10.

(2) \Rightarrow (3) : If (2) holds then (3) follows by taking $A = Q$, $B = \text{Res}(E)$, $D = C$, $K = \{0\}$ and applying Theorem 5.

(3) \Rightarrow (1) : Let $e, f \in \mathcal{R}(D)$, say $eA = R_K(s)$ and $fA = R_K(t)$,

e and f being idempotents in B . Since $te \in DB \subseteq D(D \cdot D) \subseteq D$ we have $R_K(te) = gA$ where $g = g^2 \in B$. Thus, from $(te)(eg) = teg \in K$ we deduce that $eg \in R_K(te) = gA$ so that $eg = geg$ and hence eg is an idempotent in B . Now

$$\begin{aligned} x \in egA \Rightarrow x = egx \Rightarrow & \begin{cases} tx = tegx \in K \Rightarrow x \in R_K(t), \\ sx = segx \in seA \subseteq K \Rightarrow x \in R_K(s) \end{cases} \\ & \Rightarrow x \in R_K(s) \wedge R_K(t); \end{aligned}$$

and conversely

$$\begin{aligned} x \in R_K(s) \wedge R_K(t) \Rightarrow x \in eA \Rightarrow x = ex \\ & \Rightarrow tex = tx \in K \\ & \Rightarrow x \in R_K(te) = gA \\ & \Rightarrow x = gx \\ & \Rightarrow x = ex = egx \in egA. \end{aligned}$$

We have thus shown that

$$eA \wedge fA = R_K(s) \wedge R_K(t) = egA.$$

We now show that $egA \in \mathcal{R}(D)$ whence it follows that $\mathcal{R}(D)$ is an \wedge -semilattice. From $egA \subseteq eA, fA$ we obtain, by Theorem 8(1), (2),

$$\hat{R}_K[L_K(eg)] \subseteq \hat{R}_K[L_K(e)] \wedge \hat{R}_K[L_K(f)] = eA \wedge fA = egA$$

and, since $eg \in \hat{R}_K[L_K(eg)]$, the reverse inclusion holds. Hence we have $\hat{R}_K[L_K(eg)] = egA$ and so, by Theorem 8(2), $egA \in \mathcal{R}(D)$. Now, by Theorem 7, A has an identity element and for any $k \in K$ we have $R_K(k) = A = 1A$; hence A is the maximum element of $\mathcal{R}(D)$. Since $K = R_K(1)$ is clearly the minimum element of $\mathcal{R}(D)$, we see that $\mathcal{R}(D)$ is a bounded \wedge -semilattice, whence so also is E .

In the next section we shall develop an analogue of Theorem 11 for the case of a \vee -semilattice.

5. Coördinatisation of \mathcal{U} -semilattices.

Let E be a \mathcal{U} -semilattice with minimum element 0 , let Q be the semigroup of quasi-residuated mappings on E and let C be the subset of Q consisting of those mappings with principal kernels. Let us now consider the subset T of Q consisting of all the quasi-residuated \mathcal{U} -homomorphisms on E . Clearly T is a subsemigroup of Q which contains the zero map. Since each residuated mapping on a \mathcal{U} -semilattice is necessarily a \mathcal{U} -homomorphism, we have $\text{Res}(E) \subseteq T$. Now each mapping θ_x as defined in Theorem 1 is readily seen to be an element of T and so we deduce from Theorem 1 that, in the semigroup T ,

$$(\forall f \in T \cap C) \quad R(f) = \theta_{f^+(0)} \circ T.$$

Now it is readily verified that in general the mapping ψ_y as defined in Theorem 2 is not an element of $\text{Res}(E)$. But if we consider for each $y \in E$ the mapping $\hat{\psi}_y : E \rightarrow E$ defined by

$$\hat{\psi}_y(x) = \begin{cases} 0 & \text{if } x \leq y; \\ x \vee y & \text{if } x \not\leq y, \end{cases}$$

then each $\hat{\psi}_y$ is idempotent. Moreover, as can readily be verified, each $\hat{\psi}_y \in \text{Res}(E)$ with $\hat{\psi}_y^+$ given by

$$\hat{\psi}_y^+(z) = \begin{cases} z & \text{if } z \geq y; \\ y & \text{if } z \not\geq y. \end{cases}$$

Finally, we observe that for each $f \in T$ which has a bounded image we have, in the semigroup T ,

$$\begin{aligned}
 g \in L(f) &\Leftrightarrow (\forall x \in E) \quad g[f(x)] = 0 \\
 &\Leftrightarrow g[f(\pi)] = 0 \\
 &\Leftrightarrow (\forall x \in E) \quad (g \circ \hat{\psi}_{f(\pi)})(x) = \begin{cases} g(0) = g(x) & \text{if } x \leq f(\pi); \\ g[x \vee f(\pi)] = g(x) & \text{if } x \not\leq f(\pi), \end{cases} \\
 &\Leftrightarrow g \circ \hat{\psi}_{f(\pi)} = g \\
 &\Leftrightarrow g \in T \circ \hat{\psi}_{f(\pi)}.
 \end{aligned}$$

These considerations lead us to the following result in which $\stackrel{d}{\cong}$ denotes dual order isomorphism and the term *left Baer assembly* is defined from the previous notion using left/right duality.

THEOREM 12 *Let E be an ordered set with minimum element 0. Then the following conditions are equivalent :*

- (1) *E is a \vee -semilattice with a maximum element π ;*
- (2) *there is a left Baer assembly $\langle A; B, D; K \rangle$ with $E \stackrel{d}{\cong} \mathcal{L}(B)$.*

Proof. (1) \Rightarrow (2): The existence of a maximum element implies that the zero map is residuated and that every isotone mapping on E has a bounded image. Since $\text{Res}(E) \subseteq T \wedge C$ the previous considerations show that $\langle T; T, \text{Res}(E); \{0\} \rangle$ is a left Baer assembly. [Note that since T is a semigroup containing id_E we have $T = T \cdot T$ in $P(T)$.] That $E \stackrel{d}{\cong} \mathcal{L}(T)$ follows from the fact that Theorems 2,4,6 carry over *in toto* from Q to the subsemigroup T (with ψ_y replaced by $\hat{\psi}_y$).

(2) \Rightarrow (1): This is analogous to the proof of (3) \Rightarrow (1) in Theorem 11.

Remark. Observe that when E is a bounded lattice E can be simultaneously coördinated by the left Baer assembly $\langle T; T, \text{Res}(E); \{0\} \rangle$ and the right Baer assembly $\langle Q; \text{Res}(E), C; \{0\} \rangle$. The latter can be

replaced by $\langle T; \text{Res}(E), \text{Res}(E); \{0\} \rangle$, which is both a right and a left Baer assembly, and an application of Theorems 11 and 12 produce the result : *an ordered set E is a bounded lattice if and only if it can be coördinated by a two-sided Baer assembly.* This result recaptures the Baer semigroup coördination of lattices since the coördination shows that whenever E is a bounded lattice $\text{Res}(E)$ is a Baer semigroup which coördinates E and if a two-sided Baer assembly is of the form $\langle A; B, B; K \rangle$ then $\langle B; K \rangle$ is necessarily a Baer semigroup.

6. Coördination of Glivenko and Glivenko-Brouwer semilattices.

By a *Glivenko semilattice* (or pseudo-complemented semilattice) we shall mean an \wedge -semilattice E with minimum element 0 in which each translation $t_x : y \rightarrow x \wedge y$ has a principal kernel. By a *Glivenko-Brouwer semilattice* (implicative semilattice, or relatively pseudo-complemented semilattice) we mean an \wedge -semilattice E with minimum element 0 in which each translation is a residuated mapping. By a *Glivenko* [resp. *Glivenko-Brouwer*] *assembly* we shall mean a right Baer assembly $\langle A; B, D; K \rangle$ in which there is an abelian idempotent subsemigroup \hat{D} of D [resp. \hat{B} of B] such that, for each $x \in D$, there exists $e \in \hat{D}$ [resp. $e \in \hat{B}$] with $R_K(x) = eA$ and, for each $e \in \hat{D}$ [resp. $e \in \hat{B}$], $eA \in \mathcal{R}(D)$.

Just as each Glivenko-Brouwer semilattice is in particular a Glivenko semilattice, it follows from the fact that $B \subseteq D$ that each Glivenko-Brouwer assembly is in particular a Glivenko assembly. The

nomenclature we have chosen for these assemblies is justified in the following coördinatisation theorem.

THEOREM 13 *Let E be an ordered set with minimum element 0 . Then the following conditions are equivalent :*

- (1) E is a Glivenko [resp. Glivenko-Brouwer] semilattice;
- (2) E can be coördinatised by a Glivenko [resp. Glivenko-Brouwer] assembly.

Proof. (1) \Rightarrow (2): If E is a Glivenko [resp. Glivenko-Brouwer] semilattice then each translation t_x has a principal kernel [resp. is residuated] and $\{t_x; x \in E\}$ is an abelian idempotent semigroup. Q and C being as before, given $f \in C$ and $g \in Q$ we have, since E has a maximum element $\pi = t_0^+(0)$,

$$\begin{aligned} g \in R(f) &\Leftrightarrow f[g(\pi)] = 0 \\ &\Leftrightarrow (\forall x \in E) \quad g(x) \leq f^+(0) \\ &\Leftrightarrow (\forall x \in E) \quad g(x) \wedge f^+(0) = g(x) \\ &\Leftrightarrow g = t_{f^+(0)} \circ g \\ &\Leftrightarrow g \in t_{f^+(0)} \circ Q. \end{aligned}$$

It follows that $R(f) = t_{f^+(0)} \circ Q$ where $t_{f^+(0)} \in C$ [resp. $t_{f^+(0)} \in \text{Res}(E)$], an immediate consequence of which is that $\langle Q; \text{Res}(E), C; \{0\} \rangle$ forms a Glivenko [resp. Glivenko-Brouwer] assembly which coördinatises E .

(2) \Rightarrow (1): Suppose that $\langle A; B, D; K \rangle$ is a Glivenko [resp. Glivenko-Brouwer] assembly which coordinatises E . If $e, g \in \hat{D}$ [resp. $e, g \in \hat{B}$] then from $eg = ge$ we deduce that $egA \subseteq eA \wedge gA$. But if $h \in \hat{D}$ [resp. $h \in \hat{B}$] is such that $hA \subseteq eA$ and $hA \subseteq gA$ then we have $h = eh =$

$geh = egh$ and so $hA \subseteq eA$. Hence in $\mathcal{R}(D)$ we have $eA \wedge gA = egA$.

In the case of a Glivenko assembly, if $e \in \hat{D} \subseteq D$ then $R_K(e) \in \mathcal{R}(D)$ so that from

$$eA \wedge fA = efA \subseteq K \iff ef \in K \iff f \in R_K(e)$$

we see that the pseudo-residual (pseudo-complement) $(eA)^*$ exists and is none other than $R_K(e)$. Hence E is a Glivenko semilattice.

In the case of a Glivenko-Brouwer assembly, if $e, f, g \in \hat{B}$ and $f^* \in D$ is such that $L_K(f) = Af^*$ then $f^*e \in DB \subseteq D(D \cdot D) \subseteq D$ and so $R_K(f^*e) \in \mathcal{R}(D)$. We then have

$$\begin{aligned} eA \wedge gA = egA \subseteq fA &\Rightarrow eg = egf = feg \\ &\Rightarrow f^*eg = f^*feg \in K \\ &\Rightarrow g \in R_K(f^*e) \\ &\Rightarrow gA \subseteq R_K(f^*e), \end{aligned}$$

and conversely

$$\begin{aligned} gA \subseteq R_K(f^*e) &\Rightarrow f^*eg \in K \\ &\Rightarrow eg \in R_K(f^*) = \hat{R}_K(Af^*) = \hat{R}_K[L_K(f)] = fA \\ &\Rightarrow eA \wedge gA = egA \subseteq fA. \end{aligned}$$

It follows that $t_{eA}^+(fA)$ exists in $\mathcal{R}(D)$ and is $R_K(f^*e)$. Thus $\mathcal{R}(D)$ is a Glivenko-Brouwer semilattice whence so also is E . [Note that in this case we have the same formula for pseudo-residuals as in the Glivenko case; for, taking f to be the central idempotent k° of K [Theorem 7(2)] we can take $f^* = 1$ [Theorem 7(1)] to obtain $t_{eA}^+(k^\circ A) = R_K(e)$.]

REFERENCES

- [1] M.F. JANOWITZ : Baer semigroups, *Duke Math.J.*, 32,1965,83-96.
- [2] M.F. JANOWITZ : A semigroup approach to lattices, *Canad.J. Math.*, 18,1966,1212-1223.
- [3] T.S. BLYTH and M.F. JANOWITZ : *Residuation Theory*, Pergamon Press (to appear).

List of CNA Professional Papers*

- PP 1
Brown, George F. and Lloyd, Richmond M., "Static Models of Bank Credit Expansion," 27 pp., 23 Sep 1969, (Published in the Journal of Financial and Quantitative Analysis, Jun 1971) AD 703 925
- PP 2
Lando, Mordechai E., "The Sex-Differential in Canadian Unemployment Data," 5 pp., 9 Jan 1970, AD 699 512
- PP 3
Brown, George F.; Corcoran, Timothy M. and Lloyd, Richmond M., "A Dynamic Inventory Model with Delivery Lag and Repair," 16 pp., 1 Aug 1969, AD 699 513
- PP 4
Kadane, Joseph B., "A Moment Problem for Order Statistics," 14 pp., 13 Jan 1970, (Published in the Annals of Mathematical Statistics, Apr 1971) AD 699 514
- PP 5
Kadane, Joseph B., "Optimal Whereabouts Search," 28 pp., Oct 1969, (Published in the Journal of the Operations Research Society of America, Vol. XIX, 1971) AD 699 515
- PP 6 - Classified
- PP 7
Friedheim, Robert L., "The Continental Shelf Issue at the United Nations: A Quantitative Content Analysis," 25 pp., 7 Jan 1970, (To be published in "Pacem in Maribus," edited by Elaine H. Burnell and Piers von Simson, Center for the Study of Democratic Instructions) (See also PP 28) AD 699 516
- PP 8
Rose, Marshall and White, Alex, "A Comparison of the Importance of Economic Versus Non-Economic Factors Affecting the Residential Housing Market During the Two Decades Subsequent to World War II," 128 pp., 15 Jan 1970, AD 699 517
- PP 9
Rose, Marshall, "A Thesis Concerning the Existence of Excess Capacity at Naval Shipyards Prior to the Escalation of Hostilities in Southeast Asia in 1964," 67 pp., 9 Jan 1970, AD 699 518
- PP 10 - Classified
- PP 11
O'Neill, David M., "The Effect of Discrimination on Earnings: Evidence from Military Test Score Results," 19 pp., 3 Feb 1970, (Published in the Journal of Human Resources, Summer 1970) AD 703 926
- PP 12
Brown, George F. and Lloyd, Richmond M., "Dynamic Models of Bank Credit Expansion Under Certainty," 29 pp., 3 Feb 1970, AD 703 931
- PP 13
Overholt, John L., "Analysis Data Inputs and Sensitivity Tests in War Games," 30 pp., Mar 1971, AD 722 858
- PP 14
Rose, Marshall, "Determination of the Optimal Investment in End Products and Repair Resources," 38 pp., 18 Feb 1970, (Published in the Annual Meeting of the American Association of Cost Engineers Proceedings, Jun 1971, Montreal, Canada) AD 702 450

*CNA Professional Papers with an AD number may be obtained from the National Technical Information Service, U.S. Department of Commerce, Springfield, Virginia 22151. Other papers are available from the author at the Center for Naval Analyses, 1401 Wilson Boulevard, Arlington, Virginia 22209.

CNA Professional Papers – Cont'd.

PP 15

Rose, Marshall, "Computing the Expected-End Product Service Time Using Extreme Value Properties of Sampling Distribution," 29 pp., 18 Feb 1970, (Published in Operations Research, Mar-Apr 1971) AD 702 451

PP 16

Rose, Marshall, "Study of Reparable Item Re-supply Activities," 35 pp., 18 Feb 1970, AD 702 452

PP 17

Brown, Lee (Lt., USN) and Rose, Marshall, "An Incremental Production for the End-Item Repair Process," 17 pp., 3 Mar 1970, (Published in Annual Conference of the American Institute of Industrial Engineers Transactions, May 1970, Cleveland, Ohio) AD 702 453

PP 18

Rose, Marshall, "Inventory and the Theory of the Firm," 14 pp., 18 Feb 1970, AD 702 454

PP 19

Rose, Marshall, "A Decomposed Network Computation for End-Product Repair Curves," 24 pp., 18 Feb 1970, AD 702 455

PP 20

Brown, George F.; Corcoran, Timothy M. and Lloyd, Richmond M., "Inventory Models with a Type of Dependent Demand and Forecasting, with an Application to Repair," 4 pp., 10 Feb 1970, (Published in Management Science: Theory Section, Mar 1971) AD 702 456

PP 21

Silverman, Lester P., "Resource Allocation in a Sequential Flow Process," 21 pp., 5 Mar 1970, AD 702 457

PP 22

Gorlin, Jacques, "Israeli Reprisal Policy and the Limits of U.S. Influence," 27 pp., 23 Mar 1970, AD 703 534

PP 23

Rose, Marshall, "An Aircraft Rework Cost-Benefit Model," 13 pp., 12 Mar 1970, (Published in the 5th Annual DoD Cost Research Symposium Proceedings, Mar 1970) AD 702 514

PP 24

Lloyd, Richmond and Sutton, S. Scott, "An Application of Network Analysis to the Determination of Minimum Cost Aircraft Pipeline Factors," 51 pp., 31 Mar 1970, (Presented at NATO Conference on Problems in the Organization and Introduction of Large Logistic Support Systems, May 1970, Luxembourg) AD 703 536

PP 25

Saperstone, Stephen, "An Approach to Semi-Markov Processes," 38 pp., 23 Mar 1970, AD 703 537

PP 26

Brown, George F. and Corcoran, Timothy M., "The Reliability of a Complex System with Spares, Repair, and Cannibalization," 45 pp., 23 Mar 1970, AD 703 538

PP 27

Fain, Janice B.; Fain, William W.; Feldman, Leon and Simon, Susan, "Validation of Combat Models Against Historical Data," 18 pp., 14 Apr 1970, (Published in 9th Symposium of the National Gaming Council Proceedings, Apr 1970) AD 704 744

PP 28

Friedheim, Robert L. and Kadane, Joseph B., "Quantitative Content Analysis of the United Nations Seabed Debates: Methodology and a Continental Shelf Case Study," 32 pp., 24 Mar 1970, (Published in International Organization, Vol. XXIV, No. 3, 1970) AD 703 539

PP 29

Saperstone, Stephen H., "Controllability of Linear Oscillatory Systems Using Positive Controls," 27 pp., Apr 1970, AD 704 745

- PP 30
DeVany, Arthur S., "The Effect of Social Policy on the Social and Private Value of a Child," 20 pp., 27 Apr 1970, AD 704 746
- PP 31
DeVany, Arthur S., "Time in the Budget of the Consumer," 51 pp., 27 Apr 1970, AD 704 747
- PP 32
Overholt, John L., "Fitting Korean War Data by Statistical Method," 11 pp., 5 May 1970, (Presented at the 9th Symposium of the National Gaming Council, Apr 1970, Washington, D.C.) AD 705 349
- PP 33
DeVany, Arthur S., "A Theory of Household Demand and Labor Supply," 23 pp., 5 May 1970, AD 705 350
- PP 34
Kadane, Joseph B. and Fisher, Franklin M.*, "The Covariance Matrix of the Limited Information Estimator and the Identification Test: Comment," 6 pp., 14 May 1970, (To be published in *Econometrica*) AD 706 310
*Department of Economics, Massachusetts Institute of Technology
- PP 35
Lando, Mordechai E., "Full Employment and the New Economics—A Comment," 4 pp., 14 May 1970, (Published in the *Scottish Journal of Political Economy*, Vol. XVII, Feb 1969) AD 706 420
- PP 36
DeVany, Arthur S., "Time in the Budget of the Consumer: The Theory of Consumer Demand and Labor Supply Under a Time Constraint," 151 pp., 15 Jun 1970, AD 708 348
- PP 37
Kadane, Joseph B., "Testing a Subset of the Over-identifying Restrictions," 7 pp., 19 Jun 1970, (To be published in *Econometrica*) AD 708 349
- PP 38
Saperstone, Stephen H., "The Eigenvectors of a Real Symmetric Matrix are Asymptotically Stable for Some Differential Equation," 19 pp., Jul 1970, AD 708 502
- PP 39
Hardy, W. C. and Blyth, T. S.*, "Quasi-Residuated Mappings and Baer Assemblies," 22 pp., 14 Jul 1970, (To be published by the Royal Society of Edinburgh)
*Mathematical Institute, University of St. Andrew
- PP 40
Silverman, Lester P. and Forst, Brian E., "Evaluating Changes in the Health Care Delivery System: An Application to Intensive Care Monitoring," 19 pp., Jul 1970, AD 710 631
- PP 41
Piersall, Charles H. (LCdr), "An Analysis of Crisis Decision-Making," 55 pp., Aug 1970, (To be published in the *American Political Science Review*) AD 719 705
- PP 42
Sullivan, John A., "Measured Mental Ability, Service School Achievement and Job Performance," 22 pp., 31 Jul 1970, AD 720 359
- PP 43
Forst, Brian E., "Estimating Utility Functions Using Preferences Revealed under Uncertainty," 13 pp., Jun 1971, (Presented at the 39th National Meeting of the Operations Research Society of America, 5 May 1971) AD 726 472
- PP 44
Schick, Jack M., "Conflict and Integration in the Near East: Regionalism and the Study of Crises," 43 pp., Oct 1970, (Presented at the 66th Annual Meeting of the American Political Science Association, Sep 1970)
- PP 45
Brown, George F. and Lloyd, Richmond M., "Fixed Shortage Costs and the Classical Inventory Model," 13 pp., Jul 1970, AD 713 057

CNA Professional Papers – Cont'd.

- PP 46
Hardy, William C. and Blyth, T. S.*, "A Coordination of Lattices by One-Sided Baer Assemblies," 21 pp., Jul 1970, (To be published by the Royal Society of Edinburgh)
*Mathematical Institute, University of St. Andrew
- PP 47
Silverman, Lester, P., "Resource Allocation in a Sequential Flow Process with an Application to the Naval Resupply System," 18 pp., Oct 1970, (Presented at the 11th American Meeting of the Institute of Management Sciences, Oct 1970; Presented at the 26th Meeting of the Military Operations Research Society, Nov 1970) AD 713 028
- PP 48
Gray, Burton C., "Writeup for B34TCNA—A Step-Wise Multiple Regression Program," 15 pp., Oct 1970, AD 713 029
- PP 49
Friedheim, Robert L., "International Organizations and the Uses of the Ocean," 88 pp., Oct 1970, (To be published in Volume of Essays on International Administration, Oxford University Press AD 714 387
- PP 50
Friedheim, Robert L. and Kadane, Joseph B., "Ocean Science in the United Nations Political Arena," 45 pp., Jun 1971, AD 731 865
- PP 51
Saperstone, Stephen H., "Global Controllability of Linear Systems with Positive Controls," 29 pp., Nov 1970, AD 714 650
- PP 52
Forst, Brian E., "A Decision-Theoretic Approach to Medical Diagnosis and Treatment," 14 pp., Nov 1970, (Presented at the Fall 1970 11th American Meeting of the Institute of Management Sciences, Oct 1970, Los Angeles, California) AD 714 651
- PP 53
Kadane, Joseph B., "On Division of the Question," 12 pp., Nov 1970, (Published in Public Choice, Fall 1971) AD 714 652
- PP 54
Kadane, Joseph B., "How to Burgle If You Must: A Decision Problem," 13 pp., May 1971, AD 723 850
- PP 55
Brown, George F., "Optimal Management of Bank Reserves," 35 pp., Aug 1970, AD 715 569
- PP 56
Horowitz, Stanley A., "Economic Principles of Liability and Financial Responsibility for Oil Pollution," 26 pp., Mar 1971, AD 722 376
- PP 57
Lando, Mordechai E., "A Comparison of the Military and Civilian Health Systems," 20 pp., Dec 1970, AD 716 897
- PP 58
Kadane, Joseph B., "Chronological Ordering of Archeological Deposits by the Minimum Path Length Method," 16 pp., Jun 1971, AD 726 475
- PP 59
Dyckman, Zachary Y., "An Analysis of Negro Employment in the Building Trades," 309 pp., Jan 1971
- PP 60
Lando, Mordechai E., "Health Services in the All Volunteer Armed Force," 33 pp., Jan 1971, (Published in Studies Prepared for the President's Commission on an All-Volunteer Force, Government Printing Office, Nov 1970) AD 716 899
- PP 61
Robinson, Jack, "Classification Management Training and Operations, An Approach," 14 pp., Jul 1971, (Presented at the 7th Annual Seminar, National Classification Management Society, Washington, D.C., 13-16 Jul 1971) AD 727 719

- PP 62
Brown, George F. and Schwartz, Arnold N., "The Cost of Squadron Operation: A Theoretical and Empirical Investigation," 10 pp., Jan 1971 (Published in the Transactions of the 1971 American Association of Cost Engineers International Meeting, Jun 1971) AD 722 377
- PP 63
Lockman, Robert F., "Analyses of Selection and Performance Measures for CNA Support Personnel," 45 pp., Feb 1971, AD 720 360
- PP 64
Utgoff, Victor A. and Kashyap, R.L.*, "On Behavior Strategy Solutions in Two-Person Zero-Sum Finite Extended Games with Imperfect Information," 37 pp., Feb 1971, (Accepted for publication in the SIAM Journal on Applied Mathematics) AD 720 361
*School of Electrical Engineering, Purdue University
- PP 65
O'Neill, David M.; Gray, Burton C. and Horowitz, Stanley, "Educational Equality and Expenditure Equalization Orders: The Case of Hobson V. Hansen," 43 pp., Feb 1971, AD 720 362
- PP 66
Schwartz, Arnold N.; Sheler, James A. (LCdr) and Cooper, Carl R. (Cdr), "Dynamic Programming Approach to the Optimization of Naval Aircraft Rework and Replacement Policies," 39 pp., Mar 1971, (To be published in the Naval Research Logistics Quarterly) AD 720 363
- PP 67
Kuzmack, Richard A., "Measures of the Potential Loss from Oil Pollution," 16 pp., Mar 1971, (Published as Chapter 13 in Legal, Economic, and Technical Aspects of Liability and Financial Responsibility as Related to Oil Pollution, The George Washington University, Dec 1970) AD 722 378
- PP 68
Blechman, Barry M. and Holt, James, T., "Cost/Effectiveness Analysis of Foreign Policy Alternatives: Need, Approach, and Prospects," 41 pp., Mar 1971, (Presented at the 1971 Western Regional Meeting of the International Studies Association, Mar 1971) AD 722 379
- PP 69
Rogers, Warren F. (Cdr), "Exact Null Distributions of Rank Test Statistics," 47 pp., Mar 1971, AD 722 380
- PP 70
Rogers, Warren F. (Cdr), "On A Theorem of Weyl," 17 pp., Mar 1971, AD 722 381
- PP 71
Lloyd, Richmond M., "Dynamic Programming Models of Short Term Bank Reserve Management," 233 pp., Mar 1971, AD 727 724
- PP 72
Kadane, Joseph B. and Iversen, Gudmund R.*, "Estimation of Multinomial Process When Only the Sum and the Number Governed by Each Process is Observed," 13 pp., Apr 1971, AD 722 382
*University of Michigan
- PP 73
Victor A. Utgoff and Kashyap, R.L.*, "On Behavior Strategy Solutions in Two-Person Zero-Sum Finite Extended Games with Imperfect Information, Part II: Determination of a Minimally Complex Behavior Strategy Solution in a Medical Decision Process," 22 pp., May 1971, (Accept for publication in the SIAM Journal on Applied Mathematics) AD 723 851
*School of Electrical Engineering, Purdue University
- PP 74
Brown, Jr. George F.; Silverman, Lester P. and Perlman, Bernard L. (AWF3), "Optimal Positioning of Inventory Stock in a Multi-Echelon System," 37 pp., May 1971, (Presented at the 39th Annual Meeting of the Operations Research Society of America, May 1971) AD 723 852
- PP 75
Stoloff, Peter H., "The Navy Personal Response Program: Review, Evaluation and Recommendations," 22 pp., Jul 1971, AD 727 725

CNA Professional Papers - Cont'd.

PP 76

Canes, Michael E., "Measurement and Selection of Defense," 21 pp., Aug 1971, AD 731 868

PP 77

McConnell, James M., "The Soviet Navy in the Indian Ocean," 16 pp., Aug 1971, AD 731 869

PP 78

Blechman, Barry M., "A Quantitative Description of Arab-Israeli Interactions, 1949-1969: Data Sets and Processor," 43 pp., Sep 1971, AD 731 870

PP 79

Wilson, Desmond P. and Brown, Nicholas (Cdr), "Warfare at Sea: Threat of the Seventies," 14 pp., Nov 1971, AD 734 856

PP 80

Weinland, Robert G., "The Changing Mission Structure of the Soviet Navy," 15 pp., Nov 1971, AD 734 077

PP 81

Forst, Brian E., "The Grisly Analytics of Death, Disability, and Disbursements." 20 pp., Nov 1971, (Presented at the 40th National Meeting of the Operations Research Society of America, 28 Oct 71) AD 732 555

PP 82

Forst, Brian E., "A Doctor's Introduction to Decision Analysis." 22 pp., Nov 1971, (Presented at the Engineering Foundation Conference on Quantitative Decision Making for the Delivery of Ambulatory Care, 22 Jul 1971) AD 732 556

PP 83

Weiher, Rodney and Horowitz, Stanley A., "The Relative Costs of Formal and On-the-Job Training for Navy Enlisted Occupations," 44 pp., Nov 1971, AD 734 857

PP 84

Weiher, Rodney and Horowitz, Stanley A., "A Production Function for Trained Recruits," 27 pp., Nov 1971, AD 734 858

PP 85

Brown, Jr. George F., "Comparison of Forecast Accuracy When the Disturbances Are Small: Directly Estimated Reduced Forms vs. K-Class Induced Reduced Forms," 17 pp., Dec 1971, AD 736 355

PP 86

Harrison, Robert A., "Multivariate Regression Analysis and Slaughter Livestock," 33 pp., Dec 1971, AD 736 356

PP 87

Harkins, James A., "Computer Software: A Major Weapon System Component." 9 pp., Jan 1972, (Presented at the Washington Chapter ACM 10th Annual Symposium, 24 Jun 71, National Bureau of Standards, Gaithersburg, Md.) AD 736 357

PP 88

Lockman, Robert F., "An Evaluation of the Applicant Interview Form for CNA Support Jobs," 19 pp., Mar 1972

PP 89

Barney, Gerald O., "System Dynamics and the Analysis of Social Policy," 27 pp., Apr 1972, (Presented at the XIX International Meeting of the Institute of Management Sciences, Apr 4-8, 1972, Houston, Tex.)

PP 90

Heider, Dr. Charles H., "An N-Step, 2-Variable Search Algorithm for the Component Placement Problem," 58 pp., Apr 1972

PP 91

Piersall, Jr. Charles H. and Borgstrom, Robert E., "Cost Analysis of Optional Methods of Shipboard Domestic Waste Disposal," 23 pp., Apr 1972, (To be presented at the Annual Northeast Regional Anti-pollution Conference at the University of Rhode Island, Jul 1972)

PP 92

Forst, Brian E., "Decision Analysis and Medical Malpractice," 30 pp., May 1972, (Presented at the 41st National Meeting of the Operations Research Society of American, New Orleans, La., 27 Apr 1972)

PP 93

Zedlewski, Edwin W., "Estimation and Inference in
Binary Response Regressions," 52 pp., May 1972