

AD-A072 519

HARVARD UNIV CAMBRIDGE MA CENTER ON DECISION AND CON--ETC F/G 12/1  
A NONSTANDARD THEORY OF GAMES. PART II. ON NON-ATOMIC REPRESENT--ETC(U)  
JUN 79 A A LEWIS

N00014-77-C-0533

UNCLASSIFIED

TR-7


NL

| OF |

AD  
A072519



END  
DATE  
FILMED  
9-79  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

**LEVEL**


CENTER ON DECISION AND CONFLICT

IN

COMPLEX ORGANIZATIONS

*P*  
*B.S.*

DA072519



DDC  
REF ID: A61116  
AUG 9 1978  
REGISTRY  
C

DDC FILE COPY



HARVARD UNIVERSITY

79 08 07 039

DDC  
RECEIVED  
AUG 9 1979  
C

6 A NONSTANDARD THEORY OF GAMES, PART II  
ON NON-ATOMIC REPRESENTATIONS  
OF \*FINITE GAMES.

AO 72518

may wish to keep

10 Alain A./Lewis\*

11 Jun 79

12 17p.

14 TR-7

9 Technical Report No. 7

Prepared under Contract No. N0014-77-C-0533  
Project No. NR 277-240  
for the Office of Naval Research

This document has been approved for public release  
and sale; its distribution is unlimited.

Reproduction in whole or part is permitted for any  
purpose of the United States Government.

Harvard University  
Littauer #308  
Cambridge, Mass. 02138

\*Presently at The Rand Corporation, Information Sciences  
Division, Santa Monica, California.

411 099

Introduction

The idea of using Loeb's construction to obtain measurable representations of \*Finite games we owe to Professor Don Brown of Yale University and remark that Professor Salim Rashid of Dartmouth College first obtained results analogous to Theorem I.13 for the case of the core of a nonstandard exchange economy in his dissertation at Yale written under Professor Brown (Reference 10). An additional framework of application for Loeb's construction has been obtained by Rashid along with Professor Robert Anderson of Princeton University in a nonstandard characterization of weak convergence (Reference 11). Also, the work of Professor H. Jerome Keisler in "Hyperfinite Model Theory," Logic Colloquium 1978 (North Holland Press, 1979), and "An Infinitesimal Approach to Stochastic Analysis," Preliminary Paper, University of Wisconsin, 1978, employs Loeb's construction to elegantly derive results concerning random social phenomena and the mathematics of stochastic processes.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

### I. Loeb Space Representations

We demonstrate that a \*Finite cooperative game,  $\Gamma^* = \langle F^*, A(F^*), v^* \rangle$ , when viewed as a construction on a non-standard \*Finite measure space of participants  $\phi^* = \langle F^*, A(F^*), u_{F^*} \rangle$ , where  $u_{F^*}$  is \*Finitely additive, has a standard non-atomic representation,  $\psi(\Gamma^*)$ , on  $\psi = \langle X(F^*), X(A(F^*)), m \rangle$ , where  $X(A(F^*))$  is the smallest  $\sigma$ -algebra containing  $A(F^*)$ . Further, the set of  $u_{F^*}$ -measurable payoff configurations  $(x^*, F^*)$  of  $\Gamma^*$  for  $u_{F^*}(S) = \frac{\|F^* \cap S\|}{\|F^*\|}$  and  $S \in A(F^*)$ , are such that  $st(x^*)$  is  $m$ -measurable in  $\psi(\Gamma^*)$ , and such that

$$\left( \frac{1}{\|F^*\|} \sum_{j \in F^*} x^*(j) = \int_{X(F^*)} st(x^*) dm \right)_{\text{Mod } M_1}$$

Definition I.1: Consider a nonstandard \*Finite measure space,  $\phi^* = \langle A(F^*), u_{F^*} \rangle$  in the sense of [1], [2], or [3], where  $F^* = [0, \omega]$ ,  $\omega \in N^* - N$ ,  $A(F^*)$  is the internal algebra of sets in  $F^*$ , and  $u_{F^*}(S) = \frac{\|F^* \cap S\|}{\|F^*\|}$  for  $S \in A(F^*)$ .

A \*Finite cooperative game on  $\phi^*$  is a triplet  $\Gamma^* = \langle F^*, A(F^*), v^* \rangle$  for  $v^*$  superadditive on  $A(F^*)$  and  $v^*(F^*)$   $Q$ -bounded in  $R_+^*$  with  $v^*(\emptyset) = 0$ .

The set of payoff configurations for  $\Gamma^*$  is denoted as  $(x^*, F^*)$ , where  $x^* : F^* \rightarrow R_+^*$ , and for every player  $j \in F^*$ ,  $x^*(j) \in M_0^+$  such that

$$\left( \sum_{j \in F^*} x^*(j) = v^*(F^*) \right)_{\text{Mod } M_1}$$

The Q-boundedness of  $v^*(F^*)$  allows normalization by  $\frac{1}{\|F^*\|}$ , so that the set of payoffs for  $F^*$  then satisfies the condition that

$$\left( \frac{1}{\|F^*\|} \sum_{j \in F^*} x^*(j) = \tilde{v}^*(F^*) \right)_{\text{Mod } M_1}$$

where  $\tilde{v}^*(F^*) = \frac{1}{\|F^*\|} v^*(F^*)$ . Alternatively, the set of payoff configurations can be characterized as:

$$(x^*, F^*) = \left\{ x^* \in (R_+^*)^{F^*} : (\forall j \in F^*) x^*(j) \in M_0^+ \dots \left( \int_{F^*} x^* du_{F^*} = \tilde{v}^*(F^*) \right)_{\text{Mod } M_1} \right\}$$

by definition 4.1 of [1].

**Definition I.2:** A function  $f : F^* \rightarrow R^*$  is said to be  $u_{F^*}$ -measurable if:

$$(u_{F^*}(S) = 0)_{\text{Mod } M_1} \Rightarrow \left( \int_S |f| du_{F^*} = 0 \right)_{\text{Mod } M_1}$$

► **Lemma I.3:** The set of payoff configurations  $(x^*, F^*)$  are  $u_{F^*}$ -measurable.

**Proof:** For  $S \in A(F^*)$ ,  $u_{F^*}(S) = \int_{F^*} x_S du_{F^*}$ . Since  $\bar{x}^* \in (x^*, F^*)$  means that  $(\forall j \in F^*) x^*(j) \in M_0^+$ , it suffices to consider  $\sup_{j \in S} \bar{x}^*(j) = \bar{x}_S^* \in M_0^+$ .

Then,  $(u_{F^*}(S) = 0)_{\text{Mod } M_1} \Leftrightarrow \left( \frac{\|F^* \cap S\|}{\|F^*\|} = 0 \right)_{\text{Mod } M_1}$  and

$$\frac{1}{\|F^*\|} \sum_{j \in S} \bar{x}^*(j) \leq \frac{1}{\|F^*\|} (\|F^* \cap S\| \cdot \bar{x}_S^*).$$

However,  $x_S^* \in M_0^+$  implies  $\left( (u_{F^*}(S) \cdot x_S^*) = 0 \right)_{\text{Mod } M_1}$ . Then

$$\left( (\bar{x}_S^* \cdot \int_{F^*} \chi_S du_{F^*}) = 0 \right)_{\text{Mod } M_1} \text{ and the result follows.}$$

Q.E.D.

The following lemmatta will be employed.

► Lemma I.4: Let  $g: X \rightarrow Y^*$  for  $X \subseteq R$  and  $Y^*$  internal in  $R^*$ . Then there exists an internal mapping  $h: X^* \rightarrow Y^*$  such that  $(h|_X) = g$ .

Proof: Robinson [4], Theorem 2.11.1.

► Lemma I.5: If  $Q \subseteq N^*$ , and  $Q$  is internal, then  $Q$  has a first element.

Proof: Robinson [4], Theorem 3.1.7.

Consider the algebra of coalitions of the game,  $\Gamma^*$ ,  $A(F^*)$ , derived from the measure space,  $\phi = \langle F^*, A(F^*), u_{F^*} \rangle$ . Then viewing  $A(F^*)$  as a standardly indexed algebra of sets,  $u_{F^*}$ , induces, by way of  $st(u_{F^*})$ , a finitely additive set valued function on  $A(F^*)$  with values in  $R_+$ .

The following lemma is used fundamentally in the representation scheme.

► Lemma I.6: Let  $\{F_j\}_{j \in \mathbb{N}}$  be a disjoint family of sets in  $A(F^*)$ , then any  $F_0 \subset \bigcup_{j \in \mathbb{N}} F_j$  is such that  $F_0 \subset \bigcup_{j=1}^K F_j$  for  $K_0 \in \mathbb{N}$ . The external union of disjoint internal sets is not in  $A(F^*)$ .

Proof: (Loeb [6], Proposition 1)

If we consider the sequence  $\{F_j\}_{j \in \mathbb{N}}$  as a mapping from  $\mathbb{N}$  into  $A(F^*)$ , then by Lemma I.4,  $\{F_j\}_{j \in \mathbb{N}}$  can be extended to the internal sequence  $\{F_j\}_{j \in \mathbb{N}^*}$ .

The following set is internal,  $\{K \in \mathbb{N}^* : F_0 \subset \bigcup_{j=1}^K F_j\} = \bar{K}$ . Then by Lemma I.5, there is a first element to  $\bar{K}$ ,  $\bar{K}_0$ . If  $K_0 \in \mathbb{N}^* - \mathbb{N}$ , then  $F_0 \not\subset \bigcup_{j=1}^{K_0-1} F_j$ . By Lemma I.5, once more,  $F_0 \not\subset \bigcup_{j \in \mathbb{N}} F_j$  by implication. This is contradictory, however. Therefore  $K_0 \in \mathbb{N}$ .

Q.E.D.

Denote by  $X(A^*F^*)$ , the smallest  $\sigma$ -algebra of sets in  $F^*$ , which contains the internal algebra  $A(F^*)$ . Then  $X(A(F^*))$  is an external algebra by the above lemma.

The principal achievement of Loeb [6] is to show, on the basis of Lemma I.6,  $u_{F^*}$  generates an outer measure on  $st(u_{F^*})$  which can be extended on  $X(A(F^*))$  to a standard countably additive measure space,  $\langle F^*, X(A(F^*)), m \rangle$ . The following is a statement of Loeb's theorem.

► Theorem I.7: The measure  $u_{F^*}$  extended to  $st(u_{F^*})$  again extends on  $\langle X(F^*), X(A(F^*)), m \rangle$  to a  $\sigma$ -additive measure  $m$ . For  $X \in X(A(F^*))$ ,  $m(X) = \inf_{S \in A(F^*), X \supset S} st(u_{F^*}(S))$ . If  $st(u_{F^*}(F^*)) < \infty$

and  $m(X)$  is  $\sigma$ -finite, then the extension is unique and for  $X \in X(A(F^*))$ ,  $m(X) = \sup_{S \in A(F^*) X \supset S} st(u_{F^*}(S))$  and for some  $S \in A(F^*)$ ,  $m((S \Delta X)) = 0$ .

Proof: Loeb [6], Theorem 1, or Royden [8], Chapter 12.

► Theorem I.8: Let  $\Gamma^* = \langle F^*, A(F^*), v^* \rangle$  be a \*Finite cooperative game defined on the \*Finite measure space  $\phi^* = \langle F^*, A(F^*), u_{F^*} \rangle$  and let  $\tilde{v}^*(F^*) \in M_0^+$ . Then  $\Gamma^*$  has a standard non-atomic representation,  $\psi(\Gamma^*)$  on  $\psi = \langle X(F^*), X(A(F^*)), m \rangle$ , the Loeb space of  $\phi^*$ .

Proof: Let each player  $j \in F^*$  be mapped via the identity mapping to himself in  $X(F^*)$  and denote by  $j_x$  and  $S_x$  the images of  $j$  and  $S \in A(F^*)$  in  $X(A(F^*))$ , respectively. Since  $\tilde{v}^*(F^*) \in M_0^+$  and  $v^*(\cdot)$  is superadditive,  $\tilde{v}^*(S) \in M_0^+$  for  $S \in A(F^*)$ . Let  $\tilde{v}_x^*(\cdot)$  be such that  $\tilde{v}_x^*(S_x) = st(\tilde{v}^*(S))$ . Then  $\psi(\Gamma^*) = \langle X(F^*), X(A(F^*)), \tilde{v}_x^* \rangle$  is a non-atomic representation of  $\Gamma^*$ .

Q.E.D.

► Theorem I.9: The set of payoff configurations  $(x^*, F^*)$  for  $\Gamma^*$  are such that  $\tilde{x}^* \in (x^*, F^*)$  implies that  $st(\tilde{x}^*)$  is  $m$ -measurable in  $\psi(\Gamma^*)$ .

Proof: By definition,  $\bar{x}^*(j) \in M_0^+$  for all  $j \in F^*$ . Thus  $\text{st}(\bar{x}^*(j)) < p$  for  $p$  standard and real. In particular,  $\sup_{j \in F^*} \text{st}(\bar{x}^*(j)) < p$ . By Proposition 13, p. 228, of Royden [8], it is sufficient to show that  $\{j_x \in X(F^*) : \text{st}(\bar{x}^*(j)) < p\}$  is measurable in  $\psi$ . However,  $\{j_x \in X(F^*) : \text{st}(\bar{x}^*(j)) < p\} = \bigcup_{K=1}^{\infty} \{j \in F^* : \bar{x}^*(j) < p - 1/K\}$  for  $K \in \mathbb{N}$ . The expression on the right is a member of  $X(A(F^*))$ .

Q.E.D.

► Theorem I.10: The set of payoff configurations  $(x^*, F^*)$  for  $\Gamma^*$  are such that  $\bar{x}^* \in (x^*, F^*)$  implies

$$\left( \int_{X(F^*)} \text{st}(\bar{x}^*) dm = \int_{F^*} \bar{x}^* du_{F^*} \right) \text{Mod } M_1$$

Proof: (Loeb [6], Theorem 3) Consider the strategic equivalence of  $\Gamma^*$  defined as  $\bar{v}^*(S) + \|S\| \cdot K$  for  $K$  standard and positive. Then  $(x^*, F^*)$  translates to  $(R_+^* + K)^{F^*}$  and each  $\bar{x}^*(j) \in M_0^+ + K$  and the following set is non-empty and denumerable:

$$D = \{r \in R : m(\text{st}\bar{x}^{*-1}(r)) > 0\}$$

Let  $T = m(F^*) + 1$  and select  $\epsilon > 0$  in  $R$  to be fixed. Let  $y_t$  be in  $R$  such that  $\sup_{j \in F^*} \text{st}(\bar{x}^*(j)) < y_t$  and construct the partition  $0 = y_0 < y_1 < \dots < y_t$  such that  $y_i \notin D$  for  $i = 1, \dots, t$  and  $y_i - y_{i-1} < \epsilon/3T$ .

Allow the following to be defined:

$$\underline{S}_{u_{F^*}} = \sum_{i=1}^t y_{i-1} u_{F^*}(\bar{x}^{*-1}[y_{i-1}, y_i])$$

$$\bar{S}_{u_{F^*}} = \sum_{i=1}^t y_i u_{F^*}(\bar{x}^{*-1}[y_{i-1}, y_i])$$

$$\underline{S}_m = \sum_{i=1}^t y_{i-1} m(\text{st}\bar{x}^{*-1}[y_{i-1}, y_i])$$

$$\bar{S}_m = \sum_{i=1}^t y_i m(\text{st}\bar{x}^{*-1}[y_{i-1}, y_i])$$

Then,

$$\underline{S}_{u_{F^*}} \leq \int_{F^*} \bar{x}^* du_{F^*} \leq \bar{S}_{u_{F^*}}$$

$$\underline{S}_m \leq \int_{X(F^*)} \text{st}(\bar{x}^*) dm \leq \bar{S}_m$$

and

$$\bar{S}_{u_{F^*}} - \underline{S}_{u_{F^*}} \leq \frac{e}{3T} \sum_{i=1}^t u_{F^*}(\bar{x}^{*-1}[y_{i-1}, y_i]) < e/3$$

$$\bar{S}_m - \underline{S}_m \leq \frac{e}{3T} \sum_{i=1}^t m(\text{st}\bar{x}^{*-1}[y_{i-1}, y_i]) < e/3$$

For  $i$ ,  $1 \leq i \leq t$ , since  $y_i \notin D$ , we have that

$$\text{st}\bar{x}^{*-1}(y_{i-1}, y_i) \leq \bar{x}^{*-1}(y_{i-1}, y_i) \leq \bar{x}^{*-1}[y_{i-1}, y_i] \leq \text{st}\bar{x}^{*-1}[y_{i-1}, y_i].$$

$$\text{Then, } m(\text{st}\bar{x}^{*-1}[y_{i-1}, y_i]) = m(\text{st}\bar{x}^{*-1}[y_{i-1}, y_i]) =$$

$$m(\text{st}\bar{x}^{*-1}(y_{i-1}, y_i)) \leq m(\bar{x}^{*-1}(y_{i-1}, y_i)) \text{ and}$$

$$u_{F^*}(\bar{x}^{*-1}(y_{i-1}, y_i)) \leq u_{F^*}(\bar{x}^{*-1}[y_{i-1}, y_i]) \leq u_{F^*}\bar{x}^{*-1}[y_{i-1}, y_i]$$

$$\text{and } m(\bar{x}^{*-1}[y_{i-1}, y_i]) \leq m(\text{st}\bar{x}^{*-1}[y_{i-1}, y_i]).$$

$$\text{Now } \left[ m(\bar{x}^{*-1}(y_{i-1}, y_i)) = u_{F^*}(\bar{x}^{*-1}(y_{i-1}, y_i)) \right]_{\text{Mod } M_1} \text{ and}$$

$$\left[ u_{F^*}\bar{x}^{*-1}[y_{i-1}, y_i] = m(\bar{x}^{*-1}[y_{i-1}, y_i]) \right]_{\text{Mod } M_1} \text{ by definition of}$$

$m$  as  $st(u_{F^*})$ , from which one obtains that

$$\left( u_{F^*}(\bar{x}^{*-1}[y_{i-1}, y_i]) = m(st\bar{x}^{*-1}[y_{i-1}, y_i]) \right)_{\text{Mod } M_1} \text{ and thus}$$

$$\left( \underline{S}_{u_{F^*}} = \underline{S}_m \right)_{\text{Mod } M_1} \text{ and finally, that}$$

$$\left( \int_{F^*} \bar{x}^* du_{F^*} = \int_{X(F^*)} s \in (\bar{x}^*) dm \right)_{\text{Mod } M_1}$$

Q.E.D.

► Corollary I.11: For  $\bar{x}^* \in (x^*, F^*)$  for  $\Gamma^*$ , a nonstandard  
\*Finite cooperative game with  $\tilde{v}^*(F^*) \in M_0^+$ ,

$$st \left[ \frac{1}{\|F^*\|} \sum_{j \in F^*} \bar{x}^*(j) \right] = \int_{X(F^*)} st(\bar{x}^*) dm$$

for  $st(\bar{x}^*) \in (x_x^*, X(F^*))$  of  $\psi(\Gamma^*)$ .

Proof: Theorem I.10 and Definition I.2.

Q.E.D.

► Corollary I.12: If  $\bar{x}^*$  is a payoff configuration of  $\Gamma^*$ ,  
then  $st(\bar{x}^*)$  is a payoff configuration of  $\psi(\Gamma^*)$ , if  
 $\tilde{v}^*(F^*) \in M_0^+$ .

Proof: Since  $\bar{x}^* \in (\bar{x}^*, F^*)$ ,  $(\forall j \in F^*) \bar{x}^*(j) \in (R_+^* \cap M_0^+)$ , and  
clearly  $st\bar{x}^*(j) \in R_+$ . By Corollary I.11, one has  
 $st(\tilde{v}^*(F^*)) = st \left[ \frac{1}{\|F^*\|} \sum_{j \in F^*} \bar{x}^*(j) \right] = \int_{X(F^*)} st(\bar{x}^*) dm$  and  
therefore  $\int_{X(F^*)} st(\bar{x}^*) dm = \tilde{v}_x^*(F_x^*)$ .

Q.E.D.

It is now possible to characterize solution concepts for the non-atomic representations of \*Finite cooperative games of the form  $\Gamma^*$ , based on the solution concepts developed in Part I of the series. As the characterization of other solution concepts is similar, we give that of the Quasi-Kernel as typical.

Recall from Part I that  $QK^*(\Gamma^*)$  is defined as the set of payoffs in  $(x^*, F^*)$  such that  $\left\{ \tilde{S}_{ij}^*(\tilde{x}^*) = \tilde{S}_{ji}^*(\tilde{x}^*) \right\} \text{Mod } M_1$  a.e. in  $F^*$ , where  $\tilde{S}_{ij}^*(\tilde{x}^*) = \max_{S \in T_{ij}^*} [v^*(S) - \sum_{j \in S} \tilde{x}^*(j)]$ . By Corollary I.12, the payoff configurations for  $\psi(\Gamma^*)$  are the standard parts of payoff configurations for  $\Gamma^*$ . Then, in an analogous fashion, let  $QK^*(\psi(\Gamma^*))$  be the set of payoff configurations in  $(x^*, F^*)$  for which

$\left\{ \tilde{S}_{i_x j_x}^*(st(\tilde{x}^*)) = \tilde{S}_{j_x i_x}^*(st(\tilde{x}^*)) \right\}$  a.e. in  $X(F^*)$ , where

$$\tilde{S}_{i_x j_x}^*(st(\tilde{x}^*)) = \max_{S_x \in T_{i_x j_x}^*} [\tilde{v}_x^*(S_x) - \int_{S_x} st(\tilde{x}^*) dm] \text{ and}$$

$$T_{i_x j_x}^* = \{S_x \in X(A(F^*)) : i_x \in S_x \dots j_x \notin S_x\}. \text{ The major difference between } QK^*(\Gamma^*) \text{ and } QK^*(\psi(\Gamma^*)) \text{ is that the latter}$$

solution concept is in terms of averages and precise quality between  $\tilde{S}_{i_x j_x}^*$  and  $\tilde{S}_{j_x i_x}^*$  except on sets having null measure in  $\psi(\Gamma^*)$ . The following variety of theorem is made accessible by the above characterization.

► **Theorem I.13:** Let  $\Gamma^*$  be a \*Finite cooperative game such that  $\tilde{v}^*(F^*) \in M_0^+$ , then for  $\tilde{x}^* \in (\tilde{x}^*, F^*)$   $st(\tilde{x}^*) \in QK^*(\psi(\Gamma^*))$  if and only if  $\tilde{x}^* \in QK^*(\Gamma^*)$ .

Proof: Let  $\bar{x}^* \in (x^*, F^*)$  such that  $\bar{x}^* \in QK^*(\Gamma^*)$ . Then we wish to show that the following holds for  $\psi(\Gamma^*)$ :

$$\tilde{S}_{i_x j_x}^* (\text{st}(\bar{x}^*)) = \tilde{S}_{j_x i_x}^* (\text{st}(\bar{x}^*)) \text{ a.e. in } X(F^*)$$

$$\text{If } \bar{x}^* \in QK^*(\Gamma^*), \text{ then } \left( \tilde{S}_{ij}^*(\bar{x}^*) = \tilde{S}_{ji}^*(\bar{x}^*) \right)_{\text{Mod } M_1} \text{ a.e. in } F^*.$$

Then by definition, one has

$$\left( \left| \max_{S \in T_{ij}^*} [v^*(S) - \sum_{j \in S} \bar{x}^*(j)] - \max_{S \in T_{ji}^*} [v^*(S) - \sum_{j \in S} \bar{x}^*(j)] \right| = 0 \right)_{\text{Mod } M_1} \text{ a.e. in } F^*$$

Then

$$\left( \left| \max_{S \in T_{ij}^*} \left[ \frac{1}{\|F^*\|} [v^*(S) - \sum_{j \in S} \bar{x}^*(j)] \right] - \max_{S \in T_{ji}^*} \left[ \frac{1}{\|F^*\|} [v^*(S) - \sum_{j \in S} \bar{x}^*(j)] \right] \right| = 0 \right)_{\text{Mod } M_1} \text{ a.e. in } F^*$$

which is simply,

$$\left( \left| \max_{S \in T_{ij}^*} (\tilde{v}^*(S) - \int_S \bar{x}^* du_{F^*}) - \max_{S \in T_{ji}^*} (\tilde{v}^*(S) - \int_S \bar{x}^* du_{F^*}) \right| = 0 \right)_{\text{Mod } M_1} \text{ a.e. in } F^*$$

Then it follows that

$$\text{st} \left( \left| \max_{S \in T_{ij}^*} (\tilde{v}^*(S) - \int_S \bar{x}^* du_{F^*}) - \max_{S \in T_{ji}^*} (\tilde{v}^*(S) - \int_S \bar{x}^* du_{F^*}) \right| \right) = 0 \text{ a.e. in } F^*$$

which implies that

$$\text{st} \left( \max_{S \in T_{ij}^*} (\tilde{v}^*(S) - \int_S \bar{x}^* du_{F^*}) \right) = \text{st} \left( \max_{S \in T_{ji}^*} (\tilde{v}^*(S) - \int_S \bar{x}^* du_{F^*}) \right) \text{ a.e. in } F^*$$

and, in addition, that

$$\max_{S \in T_{ij}^*} \left[ \text{st}(\tilde{v}^*(S)) - \text{st} \left( \int_S \bar{x}^* du_{F^*} \right) \right] = \max_{S \in T_{ji}^*} \left[ \text{st}(\tilde{v}^*(S)) - \text{st} \left( \int_S \bar{x}^* du_{F^*} \right) \right] \text{ a.e. in } F^*$$

Since players in  $F^*$  map identically to themselves as players in  $X(F^*)$ , and since negligible sets in  $F^*$  are null

in  $X(F^*)$ , by the definitions of Theorem I.8 and the result of Corollary I.11, one obtains from the last expression that

$$\max_{S_x \in T_{i_x j_x}^*} \left( \bar{v}_x^*(S_x) - \int_{S_x} st(\bar{x}^*) dm \right) = \max_{S_x \in T_{j_x i_x}^*} \left( \bar{v}_x^*(S_x) - \int_{S_x} st(\bar{x}^*) dm \right) \text{ a.e. in } X(F^*)$$

and therefore that  $\bar{S}_{i_x j_x}^*(st(\bar{x}^*)) = \bar{S}_{j_x i_x}^*(st(\bar{x}^*))$  a.e. in  $X(F^*)$ .

The rest is clear.

The above reasoning is completely reversible to obtain the converse.

Q.E.D.

References

- [1] A. R. Bernstein and F. Wattenberg, "Nonstandard Measure Theory," in Applications of Nonstandard Analysis to Analysis and Probability Theory, W.A.J. Luxemburg, ed., Holt, Rhinehart and Winston, 1969.
- [2] C. Ward Henson, "On the Nonstandard Representation of Measures," Trans. AMS, Vol. 172, pp. 437-446, 1972.
- [3] P. A. Loeb, "A Nonstandard Representation of Measurable Spaces  $L_{\infty}$  and  $L_{\infty}^*$ ," Contributions to Nonstandard Analysis, North Holland, 1972.
- [4] A. Robinson, Nonstandard Analysis, North Holland, 1966.
- [5] R. Anderson, \*Finite Probability, Doctoral dissertation, Yale University, 1977.
- [6] P. A. Loeb, "Conversion from Nonstandard to Standard Measure Spaces," Trans. AMS, Vol. 211, 1975.
- [7] W.A.J. Luxemburg, "Applications of Ultrapowers to Analysis," Bull. AMS, 1962.
- [8] H. Royden, Real Analysis, Macmillan, 1968.
- [9] C. Ward Henson, "Unbounded Loeb Measures," University of Illinois paper, May 1977.
- [10] S. Rashid, Economics with Infinitely Many Traders, Doctoral dissertation, Yale University, 1976.
- [11] R. M. Anderson and S. Rashid, "A Nonstandard Characterization of Weak Convergence," Proceedings AMS, Vol. 69, No. 2, May 1978.
- [11] Alain A. Lewis, "A Nonstandard Theory of Games, Part I: On the Existence of the Quasi-Kernel and Related Solution Concepts for \*Finite Cooperative Games," Discussion Paper, Center on Decision and Conflict in Complex Organizations, Harvard University, 1979.
- [12] Alain A. Lewis, "A Nonstandard Theory of Games, Part II: On Non-Atomic Representations of \*Finite Games," Discussion Paper, Center on Decision and Conflict in Complex Organizations, Harvard University, 1979.

- [14] Alain A. Lewis, "A Nonstandard Theory of Games, Part III: Noncooperative \*Finite Games," Discussion Paper, Center on Decision and Conflict in Complex Organizations, Harvard University, 1979.
- [15] Alain A. Lewis, "A Nonstandard Theory of Games, Part IV: Equilibrium Points for \*Finite Games," Discussion Paper, Center on Decision and Conflict in Complex Organizations, Harvard University, 1979.

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Center on Decision and Conflict in Complex Organizations		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP	
3. REPORT TITLE A Nonstandard Theory of Games, Part II: On Non-Atomic Representations of *Finite Games			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report No. 7			
5. AUTHOR(S) (First name, middle initial, last name) Alain A. Lewis *			
6. REPORT DATE June, 1979	7a. TOTAL NO. OF PAGES 14	7b. NO. OF REFS 15	
8a. CONTRACT OR GRANT NO. N00014-77-C-0533	9a. ORIGINATOR'S REPORT NUMBER(S) Technical Report No. 7		
b. PROJECT NO. NR-277-240	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
c.			
d.			
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its publication is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Logistics and Mathematics Statistics Branch, Department of the Navy, Office of Naval Research, Wash. D.C.	
13. ABSTRACT The techniques of Nonstandard Measure Theory, as developed by Professor Peter A. Loeb of the University of Illinois, are employed to demonstrate that the *Finite games treated in Part I of this series, and accordingly the solution concepts defined in that context, have standard non-atomic representations on the Loeb Space generated by the internal algebra of coalitions.  *Presently at The Rand Corporation, Information Sciences Division, Santa Monica, California.			