

AD-A072 689

DAVID W TAYLOR NAVAL SHIP RESEARCH AND DEVELOPMENT CE--ETC F/G 12/1
ON THE SOLUTION OF A CERTAIN LINEAR LEAST SQUARES FIT PROBLEM.(U)
JUN 79 D A GIGNAC

UNCLASSIFIED

DTNSRDC/CMLD-79/07


NL

1 OF 1
AD
A072689




END
DATE
FILMED

9 - 79
DDC



LEVEL II



**DAVID W. TAYLOR NAVAL SHIP
RESEARCH AND DEVELOPMENT CENTER**

Bethesda, Md. 20084


DTNSRDC/CMLD-79/07

DDC FILE COPY
ON THE SOLUTION OF A CERTAIN LINEAR LEAST SQUARES FIT PROBLEM
AD A 072689

ON THE SOLUTION OF A CERTAIN LINEAR LEAST
SQUARES FIT PROBLEM

Donald A. Gignac

APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.




COMPUTATION, MATHEMATICS, AND LOGISTICS DEPARTMENT
DEPARTMENTAL REPORT

June 1979

DTNSRDC/CMLD-79/07

79 08 13 085

MAJOR DTNSRDC ORGANIZATIONAL COMPONENTS



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER DTNSRDC/CMLD-79/07	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON THE SOLUTION OF A CERTAIN LINEAR LEAST SQUARES FIT PROBLEM		5. TYPE OF REPORT & PERIOD COVERED Interim July 1978 - April 1979
7. AUTHOR(s) Donald A. Gignac		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS David W. Taylor Naval Ship Research and Development Center Bethesda, Maryland 20084		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS SR 0140301, ZF 53532001 Work Units: 1-1808-009 1-1808-010
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE JUNE 1979
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		13. NUMBER OF PAGES 26
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
18. SUPPLEMENTARY NOTES		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Least Squares Fits Matrix Analysis Linear Equations Holography Error Bounds Method of Least Squares Singular Value Decomposition Error Analysis		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report investigates the difficulties encountered with respect to a certain linear least squares fit problem which arises in the field of holographic interferometry. It was found that the addition of another variable to the equations of this linear least squares fit problem explained these difficulties and provided consistent results. This report also demonstrates the usefulness of Stewart's recently published perturbation bounds for the purpose of error bounding.		

DD FORM 1473 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE S/N 0102-014-6601

406 847 UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

TABLE OF CONTENTS

	Page
ABSTRACT.....	1
ADMINISTRATIVE INFORMATION.....	1
INTRODUCTION.....	1
STEWART'S PERTURBATION BOUNDS.....	2
THE TESTS.....	5
OBSERVATIONS AND CONCLUSIONS.....	12
PROGRAMS.....	13
ACKNOWLEDGMENTS.....	21
REFERENCES.....	21

LIST OF TABLES

1 - Inequality 1 Results.....	6
2 - Inequalities 2 and 3 Results.....	7
3 - Inequalities 1, 2, and 3 Results (Joint Variation).....	8
4 - Results Illustrative of the Effect of the Fourth Parameter in the Modeling of the Problem.....	9
5 - Upper Bounds to Inequalities 2 and 3 Bounds.....	10

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced Justification	<input type="checkbox"/>
By _____	
Distribution/ _____	
Availability Codes	
Dist.	Avail and/or special
A	

ABSTRACT

This report investigates the difficulties encountered with respect to a certain linear least squares fit problem which arises in the field of holographic interferometry. It was found that the addition of another variable to the equations of this linear least squares fit problem explained these difficulties and provided consistent results. This report also demonstrates the usefulness of Stewart's recently published perturbation bounds for the purpose of error bounding.

ADMINISTRATIVE INFORMATION

This work was performed under NAVSEA Mathematical Sciences Program, "Numerical Methods for Naval Vehicles," Program Element 61153N, Task Area SR 0140301, Task 15321, DTNSRDC Work Unit 1-1808-010. The NAVSEA cognizant program manager is Ms. B. Orleans, SEA 03R.

INTRODUCTION

Recently Dr. Surendra K. Dhir (Head, Numerical Structural Mechanics Branch, Code 1844) called the author's attention to difficulties arising from a linear least squares fit problem involving holographic displacement measurements.^{1*} In the problem described, it is desired to fit a polynomial of the form $w = ax + by + cz$ by a set of experimentally obtained data points (w_i, a_i, b_i, c_i) , $i=1, \dots, n$; but under certain circumstances the polynomial would have to be set up in a differenced form, i.e.,

$$w_{i+1} - w_i = (a_{i+1} - a_i)x + (b_{i+1} - b_i)y + (c_{i+1} - c_i)z \quad i=1, \dots, n-1$$

Although the polynomial can be fitted to this "differenced" data, the relationship between this "differenced" problem and the original problem is neither apparent nor simple.

One possible approach is to regard the two least squares fit problems as perturbations of one another (after adding a redundant equation to the

* A complete listing of references is given on page 21.

reduction involves no loss of generality since any $m \times n$ linear least square fit problem $Ax=b$ and any of its perturbations $(A+E)(x+h)=b+k$ can be reduced to this required form by the respective transformations

$$(U^T A V)(V^T x) = U^T b$$

$$(U^T A V + U^T E V)(V^T x + V^T h) = U^T b + U^T k$$

For the purpose of our problem we shall also assume that the columns of A are linearly independent and that the first n rows of A are also linearly independent.

Let the linear least squares fit problem $A(x+h)=b+k$ result from a perturbation of the right-hand side of $Ax=b$. Define κ as $\|A\|_2 \|A^\dagger\|_2^*$ and η as $\|b_1\|_2 / \|A\|_2 \|x\|_2$ where b_1 is the sub-vector of the first n components of b . Define k_1 similarly. Then Stewart provides the perturbation bound

$$\|h\|_2 / \|x\|_2 \leq \kappa \eta \|k_1\|_2 / \|b_1\|_2 \quad (\text{Inequality 1})$$

Next let the linear least squares fit problem $(A+E)x=b$ result from a perturbation of the coefficient matrix of $Ax=b$. Define $\bar{\kappa}$ as $\|A\|_2 \|B_{11}^{-1}\|_2$ where B_{11}^{-1} is the inverse of the n^{th} order principal submatrix of $A+E$. Let E_{11} be the n^{th} order principal submatrix of E and let E_{21} be the $(m-n) \times n$ submatrix of E below E_{11} . Define η and b_1 as before.

* The reader is no doubt familiar with the concept of the norm of a vector x , written $\|x\|$, and defined by

$$\|x\| = \sqrt{\sum_{i=1}^n x_i^2}$$

where the x_i are the components of the vector x . Since there are several norms defined for vectors, let us be more precise and refer to the above norm as $\|x\|_2$. For A , an $m \times n$ real matrix, $\|A\|_2$ is defined to be σ_1 , the largest singular value of A . It can be shown that $\|x\|_2 = \|Vx\|_2$ and $\|A\|_2 = \|U^T A V\|_2$ where U and V are unitary matrices. Accordingly this norm (along with others) is said to be unitarily invariant. For more information on norms and their properties the reader is referred to Stewart.³

Let b_2 be the $m-n$ subvector of b below b_1 . Then Stewart obtains the perturbation bound

$$\frac{\|h\|_2}{\|x\|_2} \leq \bar{\kappa} \frac{\|E_{11}\|_2}{\|A\|_2} + \frac{\|E_{21}\|_2}{\bar{\kappa}^2 \|A\|_2} \left(\eta \frac{\|b_2\|_2}{\|b_1\|_2} + \frac{\|E_{21}\|_2}{\|A\|_2} \right) \quad (\text{Inequality 2})$$

Define the real valued matrix function

$$\psi(F) = \frac{\|F\|_2}{(1 + \|F\|_2^2)^{1/2}}$$

Stewart bounds the last term of Inequality 2 and obtains another perturbation bound

$$\frac{\|h\|_2}{\|x\|_2} \leq \bar{\kappa} \frac{\|E_{11}\|_2}{\|A\|_2} + \bar{\kappa} \eta \frac{\|b_2\|_2}{\|b_1\|_2} \psi \left(\frac{\bar{\kappa} \|E_{21}\|_2}{\|A\|_2} \right) \quad (\text{Inequality 3})$$

Clearly these three inequalities can be used to provide "worst possible case" bounds for a given least square fit problem $Ax=b$. In the case of Inequality 1 one needs only to provide some upper bound for $\|k_1\|_2$ to obtain such a bound for $\|h\|_2 / \|x\|_2$ when only the right-hand side b is perturbed. As regards Inequalities 2 and 3 one must provide upper bounds for $\|E_{11}\|$ and $\|E_{21}\|$ to obtain similar worst possible case bounds for $\|h\|_2 / \|x\|_2$ when only A is perturbed. The reader is reminded that

$$\|(A_{11} + E_{11})^{-1}\|_2 \leq \frac{\|A_{11}^{-1}\|_2}{1 - \|\bar{A}_{11}\|_2 \|E_{11}\|_2}$$

(see Wilkinson,⁴ p. 92) if $\|\bar{A}_{11}^{-1}\|_2 \|E_{11}\|_2 < 1$. In Inequality 3 the function ψ may be bounded from above by 1.0.

THE TESTS

We shall refer to the 7 x 3 linear least squares fit problem $Ax = b$ where

$$A = \begin{matrix} -.21458051E+00 & -.59223911E+00 & -.18537080E+01 \\ -.35869262E+00 & -.54449323E+00 & -.18176699E+01 \\ -.22761852E+00 & -.47379546E+00 & -.18959881E+01 \\ -.30491734E+00 & -.45507956E+00 & -.18743727E+01 \\ -.37405904E+00 & -.43714104E+00 & -.18487759E+01 \\ -.23612950E+00 & -.35108814E+00 & -.19235878E+01 \\ -.38382191E+00 & -.32889374E+00 & -.18685387E+01 \end{matrix}$$

and

$$b = \begin{matrix} .36055440E-03 \\ .35650322E-03 \\ .35852881E-03 \\ .35650322E-03 \\ .35447764E-03 \\ .35650322E-03 \\ .35245205E-03 \end{matrix}$$

as the "original" problem. It is estimated that the elements of the columns of A and b are subject to uncertainties of 5%, 11%, 2%, and 5%, respectively.

We define the "differenced" 7 x 3 linear least squares fit problem thus: for $i = 1, \dots, 6$ subtract the i^{th} equation from the $i+1^{\text{st}}$ equation of the original problem; the seventh equation of the differenced problem is the difference of the first and seventh equations of the original problem. We define the "augmented" 7 x 4 linear least squares fit problem merely by adding a fourth column of 1.0's to the coefficient matrix of the original problem and another unknown x_4 to the solution vector. Now solve this augmented problem for the value of x_4 , take the original 7 x 3 linear least squares fit problem $Ax = b$ and subtract the value of x_4 from the right-hand side b of the original problem. We shall refer to the resulting 7 x 3 linear least squares fit problem as the "modified" problem.

These linear least square fit problems were perturbed ten times each. The ratio of the norm of the solution perturbation to the solution norm was computed along with their bounds provided by Inequalities 1 through 3. These results are compared in Tables 1-5 for perturbations* of either or

* The perturbations in Table 5 were reduced by a factor of 10 to meet the condition $\|A_{11}^{-1}\| \|E_{11}\| < 1$ for the approximation of $\|(A_{11} + E_{11})^{-1}\|_2$.

TABLE 1 - INEQUALITY 1 RESULTS

Perturbation	Original Problem			Differenced Problem			Augmented Problem		
	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 1 Bound	Worst Possible Bound 1	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 1 Bound	Worst Possible Bound 1	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 1 Bound	Worst Possible Bound 1
1	0.162	0.425	1.426	0.409	0.555	1.714	3.482	8.377	25.538
2	0.307	0.349	1.437	0.300	0.496	1.728	3.550	7.198	25.741
3	0.202	0.280	1.438	0.347	0.729	1.738	2.884	5.785	25.760
4	0.333	0.371	1.449	0.239	0.852	1.751	2.951	7.285	25.962
5	0.305	0.417	1.450	0.286	1.116	1.761	2.288	7.811	25.984
6	0.193	0.238	1.451	0.214	0.241	1.725	2.227	4.803	26.001
7	0.175	0.251	1.442	0.377	0.497	1.732	2.655	5.230	25.841
8	0.132	0.184	1.443	0.117	0.609	1.742	5.152	6.115	25.856
9	0.181	0.310	1.434	0.407	0.926	1.741	3.092	6.355	25.695
10	0.310	0.339	1.446	0.300	1.032	1.754	3.160	6.844	25.897

TABLE 2 - INEQUALITIES 2 AND 3 RESULTS

Perturbation	Original Problem		Differenced Problem		Modified Problem				
	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound
1	0.090	0.420	0.201	0.326	1.016	0.769	0.098	0.401	0.184
2	0.181	0.534	0.300	0.424	1.520	0.999	0.172	0.514	0.282
3	0.068	0.281	0.143	0.412	1.382	1.023	0.057	0.266	0.129
4	0.152	0.352	0.192	0.511	1.745	1.245	0.085	0.336	0.178
5	0.190	0.578	0.274	0.580	2.424	1.523	0.167	0.556	0.255
6	0.071	0.411	0.175	0.406	2.043	1.589	0.021	0.392	0.159
7	0.091	0.408	0.213	0.673	2.659	1.896	0.061	0.391	0.197
8	0.248	0.787	0.383	0.563	3.048	1.988	0.167	0.759	0.360
9	0.061	0.428	0.127	0.543	4.166	2.562	0.105	0.406	0.107
10	0.155	0.500	0.208	0.514	3.220	2.261	0.086	0.478	0.189

TABLE 3 - INEQUALITIES 1, 2, AND 3 RESULTS (JOINT VARIATION)

Perturbation	Original Problem		Differenced Problem	
	$\frac{\ h\ _2}{\ x\ _2}$	Inequalities 1+2 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequalities 1+2 Bound
1	0.327	0.845	0.716	1.571
2	0.549	0.883	0.672	2.016
3	0.260	0.561	0.695	2.111
4	0.482	0.723	0.661	2.597
5	0.417	0.995	0.841	3.540
6	0.236	0.649	0.604	2.284
7	0.159	0.659	1.019	3.156
8	0.247	0.971	0.608	3.657
9	0.263	0.738	0.652	5.092
10	0.489	0.839	0.662	4.252

TABLE 4 - RESULTS ILLUSTRATIVE OF THE EFFECT OF THE FOURTH PARAMETER IN THE MODELING OF THE PROBLEM

Perturbation	Augmented Problem-- All Four Columns of a Perturbed			Augmented Problem-- First Three Columns of a Perturbed			Modified Problem-- Joint Variation			
	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 1 Bound	Inequality 2 Bound	Inequality 3 Bound
1	1.033	55.454	3.802	0.234	854.805	12.382	6.489	164.686	0.401	0.184
2	0.974	1086.539	16.866	0.120	2270.770	30.324	7.305	166.113	0.514	0.282
3	0.907	17.133	1.961	0.066	94.114	3.193	6.357	166.241	0.266	0.129
4	0.976	25.356	2.402	0.076	38.646	2.852	7.100	167.669	0.336	0.178
5	1.006	14.791	1.795	0.189	7.000	6.996	6.879	167.804	0.556	0.255
6	0.839	32.004	2.228	0.048	72.742	3.438	6.331	167.940	0.392	0.159
7	0.914	8.517	1.816	0.119	2473.818	22.534	6.128	166.783	0.391	0.197
8	1.012	1191.792	16.300	0.168	140.175	6.585	6.329	166.917	0.760	0.360
9	1.006	36.107	1.741	0.063	44.546	2.479	6.353	165.765	0.406	0.107
10	1.002	142.086	3.737	0.306	341.516	6.784	7.113	167.192	0.478	0.189

TABLE 5 - UPPER BOUNDS TO INEQUALITIES 2 AND 3 BOUNDS

Perturbation	Original Problem			Differenced Problem			Modified Problem		
	Worst Possible Inequality 2 Bound = 0.2513 Worst Possible Inequality 3 Bound = 0.2123			Worst Possible Inequality 2 Bound = 3.5122 Worst Possible Inequality 3 Bound = 2.4646			Worst Possible Inequality 2 Bound = 0.2418 Worst Possible Inequality 3 Bound = 0.2028		
	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound	$\frac{\ h\ _2}{\ x\ _2}$	Inequality 2 Bound	Inequality 3 Bound
1	0.0082	0.0219	0.0197	0.0285	0.0740	0.0721	0.0094	0.0201	0.0180
2	0.0166	0.0313	0.0291	0.0384	0.1056	0.1014	0.0181	0.0294	0.0273
3	0.0069	0.0160	0.0146	0.0382	0.0963	0.0937	0.0044	0.0145	0.0132
4	0.0146	0.0210	0.0195	0.0480	0.1242	0.1204	0.0084	0.0195	0.0179
5	0.0189	0.0305	0.0276	0.0520	0.1665	0.1593	0.0176	0.0284	0.0254
6	0.0040	0.0217	0.0191	0.0379	0.1596	0.1561	0.0016	0.0197	0.0170
7	0.0096	0.0240	0.0219	0.0610	0.1908	0.1851	0.0046	0.0221	0.0201
8	0.0218	0.0390	0.0356	0.0590	0.2360	0.2261	0.0180	0.0367	0.0333
9	0.0061	0.0155	0.0127	0.0558	0.2514	0.2412	0.0064	0.0134	0.0106
10	0.0148	0.0242	0.0213	0.0499	0.2645	0.2553	0.0094	0.0220	0.0191

both the right- and left-hand sides of these problems. We shall comment on these results in the next section. These results were obtained by single precision computation on the CDC 6400 at DTNSRDC. Subroutines from the IMSL Library⁵ were used to compute singular value decompositions, solve linear least squares fit problems, and generate random numbers.

Let

$$A' = \begin{pmatrix} -.21458043E+00 & -.61617945E+00 & -.18521478E+01 \\ -.36091016E+00 & -.52242982E+00 & -.18192539E+01 \\ -.21953501E+00 & -.52034141E+00 & -.18609241E+01 \\ -.31149031E+00 & -.41563715E+00 & -.18808611E+01 \\ -.36469258E+00 & -.43905926E+00 & -.18250381E+01 \\ -.23855953E+00 & -.32482739E+00 & -.19453948E+01 \\ -.38297324E+00 & -.33146348E+00 & -.18640814E+01 \end{pmatrix}$$

and

$$b' = \begin{pmatrix} .36055496E-03 \\ .34712449E-03 \\ .35893074E-03 \\ .34162634E-03 \\ .35680069E-03 \\ .34089129E-03 \\ .35576825E-03 \end{pmatrix}$$

be the perturbations of the A and b matrices defined previously. The solution of the original problem $Ax = b$ is

$$x = \begin{pmatrix} -.41300149E-04 \\ -.66551182E-04 \\ -.16793035E-03 \end{pmatrix}$$

The solution x' of the problem $Ax' = b$ (only b perturbed) is

$$x' = \begin{pmatrix} -.67766679E-04 \\ -.77162085E-04 \\ -.15851240E-03 \end{pmatrix}$$

The solution x'' of $A'x'' = b$ (only coefficient matrix perturbed) is

$$x'' = \begin{pmatrix} -.56582677E-04 \\ -.72542926E-04 \\ -.16455871E-03 \end{pmatrix}$$

The solution x''' of $A'x''' = b'$ (both b and coefficient matrix perturbed) is

$$x''' = \begin{pmatrix} -.88120583E-04 \\ -.10061679E-03 \\ -.15007406E-03 \end{pmatrix}$$

These results should prove useful as benchmark results for those who wish to run the problems published in this report.

OBSERVATIONS AND CONCLUSIONS

Let us begin by explaining the discrepancy between the solutions of the original and the differenced linear least squares fit problems. A system of m linear equations in n unknowns $Ax = b$ is said to be consistent if there exists a solution vector x which satisfies each equation of this system exactly. It is obvious from the rules of algebraic manipulation that any solution of a consistent system is likewise a solution of its differenced system. If $Ax = b$ is inconsistent, we define the linear least squares fit "solution" to be that real vector x which minimizes the norm of the residual vector $r = Ax - b$. Thus it is merely a notational convenience to write an inconsistent linear least squares fit problem as $Ax = b$, since equality obtains only when we include the residual vector r thus

$$Ax = b + r$$

If we construct the differenced system by subtracting the equations of this last system from one another, we can see that the residual vector r' for x with respect to the differenced system is

$$r' = \begin{pmatrix} r_1 - r_2 \\ r_2 - r_3 \\ \vdots \\ r_n - r_1 \end{pmatrix}$$

In general r' is not the residual vector of minimum norm for the differenced system and hence x is not the least squares fit solution for the differenced system.

Next let us consider the modeling of the original 7×3 problem. If we include an extra variable k in each equation of the original system (thus obtaining the 7×4 augmented system), we find that the first three unknowns of the augmented system least squares fit solution do not agree at all with the least squares fit solution of the original system. Rather,

they coincide with the least squares fit solution of the 7×3 system whose coefficient matrix is A but whose right-hand side is diminished by the value of k (modified system). Clearly this fourth unknown k is crucial.

Finally, this investigation has demonstrated the value of using Stewart's perturbation bounds in the error analysis of the linear least squares fit problem. In particular the Inequality 1 estimates seem to bound $\|h\|_2/\|x\|_2$ quite nicely when only b is perturbed. In fact these Inequality 1 bounds are roughly a third of the worst possible error estimate afforded by Inequality 1. As might be expected the Inequalities 2 and 3 estimates do not bound $\|h\|_2/\|x\|_2$ when A is perturbed as closely as the Inequality 1 estimates do for perturbed b . However at worst the Inequalities 2 and 3 estimates appear to be not greater than eight times $\|h\|_2/\|x\|_2$. Usually the Inequality 3 bounds are "sharper" (closer) than the Inequality 2 bounds, although this is not always the case. These Inequalities 2 and 3 bounds are roughly a tenth of the worst possible error estimates provided by Inequalities 2 and 3, respectively.

So far all results obtained indicate that the errors from a joint perturbation of A and b are roughly the sum of the respective errors and hence may be bounded by the sum of the bounds provided by Inequalities 1 and 2 or 1 and 3.

PROGRAMS

The following programs were used to obtain some of the results published in this report. The reader is reminded of the special row and column rank conditions holding for this problem. In particular, the procedure for obtaining an $m \times m$ unitary matrix U' from the $m \times m$ unitary matrix U output from the LSVLR subroutine works only under these conditions. (The last $m-n$ columns of the $m \times m$ identity matrix I are annexed to the $m \times m$ matrix U . U' is obtained by orthogonalizing these m columns using Gram-Schmidt algorithm. Of course, this procedure can be generalized by modifying it with suitable before and after permutations.)

```

PROGRAM SKDPROB(OUTPUT,TAPE6=OUTPUT)
DIMENSION A(7),B(7),C(7),AA(7,3),BB(7),R(7),XKAREA(99),X(7),Y(7),
L Z(7),XN(7),S(7),CC(7),CCOS(3,3),Q(7,3),H(7),V(7,7),G(7,3)
DIMENSION TKK(3,3),BQ(3)
DIMENSION T(7)
DIMENSION QQ(7,3),QB(7),QC(7,7),QD(3,3),QE(3)
DIMENSION QU(7,7),XKK(7),XK(7)
DIMENSION QF(7)
DIMENSION QP(7,3)
DIMENSION E11(7,3),GG(7,3),GQ(7),GU(7,3),GV(3,3),G4(7,3),EL(7,3),
L QR(7,3),ET(7,3)
DIMENSION E21(4,3)
DIMENSION CF(7)

DO 34 JQ=1,10
IF (J2.EQ.1) K2=JQ

XX=-5.165+6.1
YY=-2.459-0.7
ZZ=2.53-19.4
X0=Y0=Z0=0.0
R0=SQRT((XX-X0)*(XX-X0)+(YY-Y0)*(YY-Y0)+(ZZ-Z0)*(ZZ-Z0))
X(1)=X(3)=X(6)=4.25
X(2)=X(5)=X(7)=5.7
X(4)=5.45
Y(1)=Y(2)=1.875
Y(3)=Y(4)=Y(5)=1.25
Y(6)=Y(7)=-1.25
Z(1)=Z(3)=Z(6)=-5.12
Z(2)=Z(5)=Z(7)=-5.2
Z(4)=-5.65
DO 1 K=1,7
R(K)=SQRT((XX-X(K))*(XX-X(K))+(YY-Y(K))*(YY-Y(K))+(ZZ-Z(K))*
L (ZZ-Z(K)))
A(K)=(XX-X0)/R)+(XX-X(K))/R(K)
B(K)=(YY-Y0)/R)+(YY-Y(K))/R(K)
1 C(K)=(ZZ-Z0)/R)+(ZZ-Z(K))/R(K)
XN(1)=17.8
XN(2)=XN(4)=XN(5)=17.6
XN(3)=17.7
XN(6)=17.5
XN(7)=17.4
XLA4BDA=5145.0*3.937 E-03
DO 47 I=1,7
Q2(I,1)=A(I)
Q2(I,2)=B(I)
Q2(I,3)=C(I)
QB(I)=XLA4BDA*XN(I)
CF(I)=QB(I)
47 QF(I)=QB(I)
B4R4=SQRT(QB(1)**2+QB(2)**2+QB(3)**2+QB(4)**2+QB(5)**2+
1 QB(6)**2+QB(7)**2)
DO 48 J=1,3
DO 48 I=1,7
QR(I,J)=Q2(I,J)
48 QP(I,J)=Q2(I,J)

```

```

M=7
N=3
ISM=1
WRITE(6,99)((Q2(I,J),J=1,3),I=1,7)
99 FORMAT(1X,2H2Q,3E16.8)
CALL LSVALR(QQ,M,N,M,N,ISM,WKAREA,QE,QC,20)
ANRM=AMAX1(QE(1),QE(2),QE(3))
XAPS[2]=AMAX1(1./QE(1),1./QE(2),1./QE(3))
WRITE(6,62)((QC(I,J),J=1,3),I=1,7)
62 FORMAT(1X,2H2C,3E16.7)
CALL VCOMP(QC,20,M,N)
WRITE(6,63)((QJ(I,J),J=1,7),I=1,7)
63 FORMAT(1X,2H2U,7E16.7)
DO 15 I=1,7
BB(I)=0.0
DO 15 K=1,7
15 BB(I)=BB(I)+2U(K,I)*QB(K)
BINRM=SQRT(BB(1)**2+BB(2)**2+BB(3)**2)
BINRM=SQRT(BB(1)**2+BB(2)**2+BB(3)**2)
B2NM=SQRT(BB(4)**2+BB(5)**2+BB(6)**2+BB(7)**2)
NB=1
IDGT=8
CALL LLSQAR(QP,23,M,N,NB,M,M,IDGT,WKAREA,IER)
WRITE(6,45)(I,2B(I),I=1,3)
45 FORMAT(1X,2H2B,I10,E16.8)
XSQD=QB(1)**2+2B(2)**2+2B(3)**2
XNRM=SQRT(XSQD)
HOLD=XNRM
KQ=JQ
CALL GGUB(KQ,7,T)
DO 30 I=1,7
30 A(I)=A(I)+(-1)**I*T(I)*.047*A(I)
CALL GGUB(2*KQ,7,T)
DO 31 I=1,7
31 B(I)=B(I)-(-1)**I*T(I)*.113*B(I)
CALL GGUB(3*KQ,7,T)
DO 32 I=1,7
32 C(I)=C(I)+(-1)**I*T(I)*.023*C(I)
CALL GGUB(4*KQ,7,T)
DO 33 I=1,7
33 XN(I)=XN(I)-(-1)**I*T(I)*.05*XN(I)
WRITE(6,212)(T(I),XN(I),I=1,7)
212 FORMAT(1X,1HT,E15.8,24XN,E16.8)
XLAMBDA=5145.0*3.937E-09
DO 2 K=1,7
BB(K)=XLAMBDA*XN(K)
XKK(K)=BB(K)-C(K)
CC(K)=BB(K)
AA(K,1)=A(K)
AA(K,2)=B(K)
2 AA(K,3)=C(K)
DO 50 LL=1,3
XK(LL)=.0
DO 50 K=1,7
60 XK(LL)=XK(LL)+2C(K,LL)*XKK(K)
XK2=XK(1)**2+XK(2)**2+XK(3)**2
XK2=SQRT(XK2)

```

```

RHS=(XAPSI2*XK2)/XNRM
WRITE(5,65)RHS
65 FORMAT(1X,4H R45, E16.8)
DO 83 I=1,7
DO 83 J=1,3
GG(I,J)=AA(I,J)
83 Q(I,J)=AA(I,J)
WRITE(5,50)
50 FORMAT(1H1/1X,9HAA AND 83)
WRITE(5,51)(AA(K,1),AA(K,2),AA(K,3),BB(K),K=1,7)
51 FORMAT(1X,3E16.8, E20.8)

DO 11 I=1,7
DO 11 J=1,3
11 ET(I,J)=QR(I,J)
DO 68 I=1,7
DO 68 J=1,3
EL(I,J)=0.0
GH(I,J)=0.0
DO 68 K=1,3
EL(I,J)=EL(I,J)+ET(I,K)*QD(K,J)
68 GH(I,J)=GH(I,J)+GG(I,K)*QD(K,J)
DO 69 I=1,7
DO 69 J=1,3
ET(I,J)=0.0
GG(I,J)=0.0
DO 69 K=1,7
ET(I,J)=ET(I,J)+QU(K,I)*EL(K,J)
69 GG(I,J)=GG(I,J)+QU(K,I)*GH(K,J)
WRITE(5,25)((ET(I,J),J=1,3),I=1,7)
26 FORMAT(1X,2HET,3E16.8)
WRITE(5,27)((GG(I,J),J=1,3),I=1,7)
27 FORMAT(1X,2HGG,3E15.8)
DO 12 I=1,7
DO 12 J=1,3
12 ET(I,J)=GG(I,J)-ET(I,J)
WRITE(5,26)((ET(I,J),J=1,3),I=1,7)
DO 71 I=1,3
DO 71 J=1,3
71 E11(I,J)=ET(I,J)
DO 72 I=4,7
DO 72 J=1,3
72 E21(I-3,J)=ET(I,J)
CALL LSVALR(GG,N,N,M,4,ISW,WKAREA,GQ,GU,GV)
WRITE(5,28)(GQ(I),I=1,3)
28 FORMAT(1X,2HS,4E16.8)
BINRM=AMAX1(1.0/GQ(1),1.0/GQ(2),1.0/GQ(3))
CALL LSVALR(E11,N,N,M,N,ISW,WKAREA,GQ,GU,GV)
WRITE(5,29)(GQ(I),I=1,3)
E11NRM=AMAX1(GQ(1),GQ(2),GQ(3))
M=4
CALL LSVALR(E21,4,N,M,4,ISW,WKAREA,GQ,GU,GV)
WRITE(5,28)(GQ(I),I=1,3)
M=7
E21NRM=AMAX1(GQ(1),GQ(2),GQ(3))
ABCDEF=BINRM*E21NRM
C4I=ABCDEF/SQRT(1.0+ABCDEF**2)

```

```

R4S=BINRM*E11NRM+(B2NM*BINRM*CHI)/XNRM
XRHS=BINRM*E11NRM+(BINRM**2*B2NM*E21NRM)/XNRM+BINRM**2*E21NRM**2
WRITE(6,64)R4S,XRHS
64 FORMAT(1X,4H R4S,E16.8,1X,5H XRHS,E16.8)
WRITE(6,22)RHS,BINRM,E11NRM,BNRM,CHI,XNRM,E21NRM
22 FORMAT(1X,1HW,7E16.8)
XKAPBAR=ANRM*BINRM
XNU=BINRM/(ANRM*4OLD)
WRITE(6,17)XKAPBAR,XNU
17 FORMAT(1X,8HKA'PABAR,E16.8,3H NU,E16.8)

```

```

C
DO 86 J=1,3
H(J)=0.
DO 78 I=1,7
78 H(J)=H(J)+AA(I,J)*AA(I,J)
86 H(J)=SQRT(H(J))
DO 80 J=1,2
ILIM=J+1
DO 80 I=ILIM,3
SUM=J.0
DO 79 K=1,7
79 SUM=SUM+AA(K,J)*AA(K,I)
80 CCOS(I,J)=SUM/(H(I)*H(J))

```

```

C
M=7
NAA=3
NBB=1
IAA=IBB=7
IDGT=8

```

```

C
CALL LLSQAR(AA,BB,M,NAA,NBB,IAA,IBB,IDGT,WKAREA,IER)

```

```

C
WRITE(6,200)((AA(K,J),J=1,3),K=1,7)
200 FORMAT(1HQ/(1X,84PSUEDINV,3E15.7))
WRITE(6,100)IER
100 FORMAT(1HQ/1X,5H IER =,I10)
WRITE(6,101)(K,BB(K),K=1,3)
101 FORMAT(1X,3HUVW,I9,E16.8)
SUM1=BB(1)*BB(1)+BB(2)*BB(2)+BB(3)*BB(3)
XNRM=SQRT(SUM1)
WRITE(6,97)SUM1,XNRM
97 FORMAT(1HQ/1X,74JVM SQD,E16.8/1X,8HSQRT JVM,E16.8)
DO 44 K=1,7
44 S(K)=CC(K)-A(K)*BB(1)-B(K)*BB(2)-C(K)*BB(3)
WRITE(6,43)(K,S(K),K=1,7)
43 FORMAT(1HQ/(1X,54RESID,I10,E15.8))

```

```

*
XLHS=
L (BB(1)-QB(1))**2+(BB(2)-QB(2))**2+(BB(3)-QB(3))**2
WRITE(6,21)(BB(I),QB(I),I=1,3)
21 FORMAT(1X,4HBBQB,2E16.8)
WRITE(6,24)XLHS,4OLD
24 FORMAT(1X,2H**2E16.8)
XLHS=SQRT(XLHS)/4OLD
WRITE(6,67)XLHS,XLHS,RHS,RHS
67 FORMAT(1X,4HXL4S,E15.8,F15.8,4H RHS,E15.8,F15.8)

```

```

C
DO 75 K=1,7
75 H(K)=SQRT(A(K)*A(K)+B(K)*B(K)+C(K)*C(K))
DO 75 I=1,6
JLIM=I+1
DO 75 J=JLIM,7
CJS=(A(I)*A(J)+B(I)*B(J)+C(I)*C(J))/(H(I)*H(J))
76 WRITE(6,77) I, J, CJS
77 FORMAT(1H0,5X,10HROW COSINE,2I3,E16.8)
DO 81 J=1,2
ILIM=J+1
DO 81 I=ILIM,3
81 WRITE(6,82) J, I, CCOS(I, J)
82 FORMAT(1H0,5X,13HCOLUMN COSINE,2I3,E16.8)

```

```

C
N=3
IA=IAA
ISW=1
CALL LSVLR(2,4,4,IA,IA,ISW,MKAREA,S,G,V)
WRITE(6,88) (K,S(K),K=1,3)
88 FORMAT(1H0/(10X,8HSING VAL,15,E16.8))
WRITE(6,89) ((G(I, J),J=1,N),I=1,M)
89 FORMAT(1H0/(1X,14U,3E15.8))
WRITE(6,90) ((V(I, J),J=1,N),I=1,N)
90 FORMAT(1H0/(1X,14V,3E15.8))
WRITE(6,94) ((TKK(I, J),J=1,3),I=1,3)
94 FORMAT(1H0/(1X,34TKK,3E15.8))
BQ(1)=BQ(2)=BQ(3)=0.0
CALL LLSQR(TKK,BQ,3,3,1,3,3,8,MKAREA,IER)
WRITE(6,93) IER
93 FORMAT(1X,5HIER =,I10)
WRITE(6,95) ((TKK(I, J),J=1,3),I=1,3)
95 FORMAT(1H0/(1X,74TKK PSI,3E16.8))

```

```

C
34 CONTINUE

```

```

C
STOP
END

```

```

SUBROUTINE VCOMP(V,Q,M,N)
DIMENSION V(7,7),VA(7,7),Q(7,7),INDEX(10),INV(10),C(10)
DIMENSION WKAREA(250)
IF (M.GT.7) GO TO 400
NPO=N+1
DO 33 J=NPO,M
DO 33 I=1,M
33 V(I,J)=.J
DO 2 I=1,M
DO 1 J=1,M
1 VA(I,J)=0.0
2 VA(I,I)=1.0
DO 3 I=1,N
3 INDEX(I)=I
DO 53 I=1,N
IF (V(I,I).NE.0.0) GO TO 49
DO 83 J=I,N
IF (V(I,J).EQ.0.0) GO TO 80
ISTORE=INDEX(I)
INDEX(I)=INDEX(J)
INDEX(J)=ISTORE
GO TO 81
80 CONTINUE
WRITE(6,83)
83 FORMAT(1H1/(1X,84ND PIVOT))
STOP
81 DO 82 K=1,M
STORE=V(K,I)
V(K,I)=V(K,J)
V(K,J)=STORE
STORE=VA(K,I)
VA(K,I)=VA(K,J)
82 VA(K,J)=STORE
49 STORE=V(I,I)
DO 51 J=I,N
V(I,J)=V(I,J)/STORE
51 VA(I,J)=VA(I,J)/STORE
IPO=I+1
DO 53 K=1,M
IF (K.EQ.I) GO TO 53
STORE=V(K,I)
DO 54 J=I,N
V(K,J)=V(K,J)-STORE*V(I,J)
54 VA(K,J)=VA(K,J)-STORE*VA(I,J)
53 CONTINUE
50 CONTINUE
IDGT=8
CALL LINV2F(VA,4,M,V,IDGT,WKAREA,IER)
WRITE(5,67) IER
67 FORMAT(1X,3HIER,I10)
DO 39 I=1,N
39 INV(INDEX(I))=I
WRITE(5,201)(INV(I),I=1,N)
201 FORMAT(1X,3HINV,10I10)
WRITE(6,200)(INDEX(I),I=1,N)
200 FORMAT(1X,5HINDEX,10I10)
DO 38 J=1,M

```

```

      DO 38 I=1,M
38  Q(I,J)=V(I,INV(J))
400 DO 402 J=N,M
      DO 411 I=1,M
401  Q(I,J)=0.0
402  Q(J,J)=1.
      DO 403 J=1,N
      DO 433 I=1,M
403  Q(I,J)=V(I,J)
      SUM=0.0
      DO 20 I=1,M
20  SUM=SUM+Q(I,1)**2
      SUM=SQRT(SUM)
      DO 21 I=1,M
21  Q(I,1)=Q(I,1)/SUM
      DO 22 L=2,M
      LMO=L-1
      DO 23 K=1,LMO
      C(K)=0.0
      DO 23 I=1,M
23  C(K)=C(K)+Q(I,L)*Q(I,K)
      DO 25 I=1,M
      DO 25 K=1,LMO
26  Q(I,L)=Q(I,L)-C(K)*Q(I,K)
      SUM=0.0
      DO 27 I=1,M
27  SUM=SUM+Q(I,L)**2
      SUM=SQRT(SUM)
      DO 28 I=1,M
28  Q(I,L)=Q(I,L)/SUM
22  CONTINUE
      DO 101 I=1,M
      DO 100 J=I,M
      SUM=0.
      DO 99 K=1,M
99  SUM=SUM+Q(K,I)*Q(K,J)
100  WRITE(5,101)I,J,SUM
101  FORMAT(1X,5HIJSUM,2I5,E15.8)
      RETURN
      END

```

ACKNOWLEDGMENTS

The author wishes to thank the following individuals for their interest and assistance: Dr. Elizabeth H. Cuthill (DTNSRDC, Code 1805); Dr. Surendra K. Dhir (DTNSRDC, Code 1844); and Mr. Robert P. Eddy (DTNSRDC, Code 1843).

REFERENCES

1. Nobis, D. and C.M. Vest, "Statistical Analyses of Error in Holographic Interferometry," *Applied Optics*, vol. 17, no. 14 (Jul 15, 1978) pp. 2198-2204.
2. Stewart, G.W., "On the Perturbation of Pseudo-Inverses, Projections and Linear Least Squares Problems," *SIAM Review*, vol. 19, no. 4 (Oct 1977) pp. 634-662.
3. Stewart G.W., "Introduction to Matrix Computations," Academic Press, New York (1973).
4. Wilkinson, J.H., "Rounding Errors in Algebraic Processes," Prentice-Hall, Englewood Cliffs, N.J. (1963).
5. International Mathematical & Statistical Libraries, Inc., "Computer Subroutine Libraries in Mathematics and Statistics," IMSL Library 3 (CDC 6000 series), vol. 2, Houston, Texas (Apr 1977).

INITIAL DISTRIBUTION

Copies		Copies	Code	Name
1	NRL/Code 8441 (J. Hansen)	10	1844	D.A. Gignac
2	NAVPGSCOL	1	1850	T. Corin
	1 59C1/G. Cantin	1	1870	M. Zubkoff
	1 M. Kelleher			
12	DDC	2	1890	G.R. Gray J.D. Strickland
1	Air Force Aero Res Lab P. Nikolai	1	1892	S.E. Good
2	BUSTAND	2	1966	J.R. Caspar Y.-N. Liu
	1 H. Oser			
	1 M. Cordes			
1	U Maryland/Prof. G.W. Stewart	1	522.1	Unclassified Lib (C)
		1	522.2	Unclassified Lib (A)

CENTER DISTRIBUTION

Copies	Code	Name
1	1552	J. Bai
4	1725	R.F. Jones, Jr. J.E. Roderick P.N. Roth L.N. Gifford, Jr.
1	1730.5	J.H. Ma
2	1740.4	M. Salive N.E. Fioravante
1	1745	C. Ng
1	1800	G.H. Gleissner
1	1802.2	F.N. Frenkiel
1	1805	E.H. Cuthill
2	1809.3	D. Harris
1	1820	A.W. Camara
2	1826	L.K. Meals L.M. Culpeper
1	1840	J.W. Schot
1	1843	R.P. Eddy
4	1844	S.K. Dhir G.C. Everstine F.M. Henderson M.M. Hurwitz

DTNSRDC ISSUES THREE TYPES OF REPORTS

1. DTNSRDC REPORTS, A FORMAL SERIES, CONTAIN INFORMATION OF PERMANENT TECHNICAL VALUE. THEY CARRY A CONSECUTIVE NUMERICAL IDENTIFICATION REGARDLESS OF THEIR CLASSIFICATION OR THE ORIGINATING DEPARTMENT.

2. DEPARTMENTAL REPORTS, A SEMIFORMAL SERIES, CONTAIN INFORMATION OF A PRELIMINARY, TEMPORARY, OR PROPRIETARY NATURE OR OF LIMITED INTEREST OR SIGNIFICANCE. THEY CARRY A DEPARTMENTAL ALPHANUMERICAL IDENTIFICATION.

3. TECHNICAL MEMORANDA, AN INFORMAL SERIES, CONTAIN TECHNICAL DOCUMENTATION OF LIMITED USE AND INTEREST. THEY ARE PRIMARILY WORKING PAPERS INTENDED FOR INTERNAL USE. THEY CARRY AN IDENTIFYING NUMBER WHICH INDICATES THEIR TYPE AND THE NUMERICAL CODE OF THE ORIGINATING DEPARTMENT. ANY DISTRIBUTION OUTSIDE DTNSRDC MUST BE APPROVED BY THE HEAD OF THE ORIGINATING DEPARTMENT ON A CASE BY CASE BASIS.