

AD-A074 182

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 11/6
PHOTOACTIVATION OF COBALT CARBONYL CATALYSTS: GENERATION OF REA--ETC(U)
SEP 79 C L REICHEL, M S WRIGHTON N00014-75-C-0880
TR-18-ONR NL

UNCLASSIFIED

| OF |

AD
A074182




END

DATE

FILMED

10-19

DDC



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

12

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS BEFORE COMPLETING FORM

1. REPORT NUMBER ONR-TR-18		2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Photoactivation of Cobalt Carbonyl Catalysts: Generation of Reactive Mononuclear Fragments from Dinuclear, Metal-Metal Bonded Complexes		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report	
7. AUTHOR(s) Carol L. Reichel and Mark S. Wrighton		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0680	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Massachusetts Institute of Technology Cambridge, MA 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-579	
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, Virginia 22217		12. REPORT DATE September 12, 1979	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 16	
LEVEL		15. SECURITY CLASS. (of this report) UNCLASSIFIED	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	

AD A 074182

16. DISTRIBUTION STATEMENT (of this Report)

Approved for public release; reproduction is permitted for any purpose of the United States Government; distribution unlimited.

17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)

Distribution of this document is unlimited.

DDC
RECEIVED
SEP 20 1979
UNCLASSIFIED

18. SUPPLEMENTARY NOTES

Prepared for publication in the Journal of the American Chemical Society.

19. KEY WORDS (Continue on reverse side if necessary and identify by block number)

photocatalysis, metal radicals, metal-metal bond complexes, photogenerated hydrides

20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

The metal-metal bonded complexes $[Co_2(CO)_6L_2]$ ($L = P(n-Bu)_3, P(OPh)_3$) and $[Fe(\eta^5-C_5H_5)(CO)_2Co(CO)_3P(OPh)_3]$ all undergo efficient ($\phi > 0.1$ at 355 nm) photochemical metal-metal bond cleavage to produce reactive 17-valence electron $[Co(CO)_3L]$ fragments. In the presence of 1-pentene no alkene isomerization is found for $L = P(n-Bu)_3$; low, but significant, photocatalytic activity is found for $L = P(OPh)_3$ or when the $P(n-Bu)_3$ complex is irradiated in the presence of $P(OMe)_3$. The $[Co(CO)_3L]$ precursors are strikingly more active when irradiation (over)

DDC FILE COPY

DD FORM 1473 1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

is carried out in the presence of HSiEt_3 , a hydride source. Both alkene isomerization and n-pentane formation are observed, along with small quantities of Si-containing products. The $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ radical photogenerated independently from $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ is not active under any conditions used thus far. The activity of 17-valence electron radicals in this application is $[\text{Co}(\text{CO})_3(\text{P}(\text{OPh})_3)] > [\text{Co}(\text{CO})_3(\text{P}(\text{n-Bu})_3)] > [\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$. These species are not themselves effective catalysts but do seem to react with HSiEt_3 to form catalytically active, mononuclear Co-hydride complexes.

Accession For	
NTIS Grant	<input checked="" type="checkbox"/>
DIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Available/or special
A	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract ¹⁵ N00014-75-C-0880

Task No. NR 051-579

⁹ TECHNICAL REPORT NO. 18

⁶ Photoactivation of Cobalt Carbonyl Catalysts:
Generation of Reactive Mononuclear Fragments from
Dinuclear, Metal-Metal Bonded Complexes.

by

¹⁴ TR-18-CNR

¹⁰ Carol L. Reichel and Mark S. Wrighton

Prepared for Publication

in the

Journal of the American Chemical Society

¹² 16

¹¹ 12 Sep 79

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

September 12, 1979

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

79 09 14 047

200 007

JLB

greater than

Photoactivation of Cobalt Carbonyl Catalysts: Generation of Reactive Mononuclear Fragments From Dinuclear, Metal-Metal Bonded Complexes

the fifth power

Abstract: The metal-metal bonded complexes $[\text{Co}_2(\text{CO})_6\text{L}_2]$ ($\text{L} = \text{P}(\text{n-Bu})_3$, $\text{P}(\text{OPh})_3$) and $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Co}(\text{CO})_3(\text{P}(\text{OPh})_3)]$ all undergo efficient ($\Phi > 0.1$ at 355 nm) photochemical metal-metal bond cleavage to produce reactive 17-valence electron $[\text{Co}(\text{CO})_3\text{L}]$ fragments. In the presence of 1-pentene no alkene isomerization is found for $\text{L} = \text{P}(\text{n-Bu})_3$; low, but significant, photocatalytic activity is found for $\text{L} = \text{P}(\text{OPh})_3$ or when the $\text{P}(\text{n-Bu})_3$ complex is irradiated in the presence of $\text{P}(\text{OMe})_3$. The $[\text{Co}(\text{CO})_3\text{L}]$ precursors are strikingly more active when irradiation is carried out in the presence of HSiEt_3 , a hydride source. Both alkene isomerization and n-pentane formation are observed, along with small quantities of Si-containing products. The $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ radical photogenerated independently from $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ is not active under any conditions used thus far. The activity of 17-valence electron radicals in this application is $[\text{Co}(\text{CO})_3(\text{P}(\text{OPh})_3)] > [\text{Co}(\text{CO})_3(\text{P}(\text{n-Bu})_3)] > [\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$. These species are not themselves effective catalysts but do seem to react with HSiEt_3 to form catalytically active, mononuclear Co-hydride complexes.



to the fifth power

Photoactivation of Cobalt Carbonyl Catalysts: Generation of Reactive Mononuclear Fragments From Dinuclear, Metal-Metal Bonded Complexes

Sir:

Metal-metal bonded complexes are generally photosensitive with respect to cleavage of the metal-metal bond,¹ and certain dicobalt complexes are known² to be hydroformylation catalyst precursors under thermal conditions. We report herein our preliminary results concerning the photogeneration of catalytically active mononuclear cobalt carbonyl fragments from dinuclear, metal-metal bonded complexes. The results illustrate the potential utility of photoinduced metal-metal bond cleavage in probing catalytic mechanisms and in initiating catalytic chemistry under thermal conditions where there would be no reaction without light activation. The complexes studied thus far are $[\text{Co}_2(\text{CO})_6\text{L}_2]$ ($\text{L} = \text{P}(\underline{n}\text{-Bu})_3, \text{P}(\text{OPh})_3$) and $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Co}(\text{CO})_3\text{P}(\text{OPh})_3]$,³ and the catalytic probe chemistry has been reaction of 1-pentene/ HSiEt_3 mixtures. The cobalt systems have been chosen for study because it is believed that the dinuclear $[\text{Co}_2(\text{CO})_6(\text{P}(\underline{n}\text{-Bu})_3)_2]$ actually forms mononuclear $[\text{HCo}(\text{CO})_3\text{P}(\underline{n}\text{-Bu})_3]$ under hydroformylation conditions.² The ability to photogenerate $[\text{Co}(\text{CO})_3\text{L}]$ under mild thermal conditions offers an opportunity to gain insight into the mechanisms of cobalt carbonyl catalyzed reactions.

All three complexes studied here undergo efficient photochemical cleavage of the metal-metal bond, as reflected in the photochemistry that we find, equations (1) and (2). $[\text{Mn}_2(\text{CO})_{10}]$ ⁴



and $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]^5$ are known to undergo efficient metal-metal bond cleavage, and the photoinduced formation of heterodinuclear complexes is believed to result from coupling of two different 17-valence electron radicals. The forward and reverse photoreactions represented in equations (1) and (2) are chemically clean, and quantum yields for near-ultraviolet light induced cleavage of the metal-metal bonds are all >0.1 .

Irradiation of $[\text{Co}_2(\text{CO})_6\text{L}_2]$ in the presence of 1-pentene in benzene or n-octane solution results in little net reaction of any kind. The disappearance of the Co-Co bonded species is very slow, particularly for $\text{L} = \text{P}(\underline{n}\text{-Bu})_3$. Presumably, the Co-Co bond is cleaved efficiently, but recoupling of the $[\text{Co}(\text{CO})_3\text{L}]$ fragments obtains to result in little net chemical change. Irradiation of the Fe-Co bonded species under the same conditions results in the redistribution reaction shown in equation (2). Table I shows the effect on 1-pentene by irradiating the $[\text{Co}_2(\text{CO})_6\text{L}_2]$ species. For $\text{L} = \text{P}(\underline{n}\text{-Bu})_3$ there is no detectable reaction in 94 h, but for $\text{L} = \text{P}(\text{OPh})_3$ there is considerable alkene isomerization and a small amount of n-pentane forms. Further, adding $\text{P}(\text{OMe})_3$ to a 1M 1-pentene solution containing $[\text{Co}_2(\text{CO})_6(\text{P}(\underline{n}\text{-Bu})_3)_2]$ results in photocatalytic activity comparable to that found for the $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$ complex.





Addition of HSiEt_3 to the 1-pentene solutions dramatically affects the photocatalytic activity, Table II. Under the same conditions, except for the presence of HSiEt_3 , the photolysis of $[\text{Co}_2(\text{CO})_6(\text{P}(\underline{n}\text{-Bu})_3)_2]$ essentially completely equilibrates the linear pentenes to the thermodynamic ratio,⁶ whereas no reaction obtains in the absence of the HSiEt_3 . A significant amount of n-pentane is formed and small quantities of Si-containing products are also detectable but have not been analyzed quantitatively. The addition of $\text{P}(\text{OMe})_3$ further accelerates the photocatalytic activity of the $\text{P}(\underline{n}\text{-Bu})_3$ complex. The effect of added HSiEt_3 on the photocatalytic activity of the $[\text{Co}(\text{CO})_3\text{P}(\text{OPh})_3]$ precursors is equally striking. The heterodinuclear Fe-Co complex is about as active as the homodinuclear precursor. Note that the $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ is essentially non-active and is, of course, the homodinuclear source of the 17-valence electron radical $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$. However, the related radical, $[\text{Fe}(\eta^3\text{-C}_3\text{H}_5)(\text{CO})_3]$, is active as an isomerization catalyst.⁷

While none of the systems exhibit significant thermal activity on the same time scale as used in the photochemical experiments, long thermal reaction time at 25° C does result in similar catalytic chemistry, and the effects of added HSiEt_3 or $\text{P}(\text{OMe})_3$ are similar. The most thermally active species is the $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$; the $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Co}(\text{CO})_3(\text{P}(\text{OPh})_3)]$ is qualitatively less active, but the least active complex is $[\text{Co}_2(\text{CO})_6(\text{P}(\underline{n}\text{-Bu})_3)_2]$.

The data suggest that the 17-valence electron radicals $[\text{Co}(\text{CO})_3\text{L}]$ ($\text{L} = \text{P}(\underline{n}\text{-Bu})_3, \text{P}(\text{OPh})_3$) or $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ are not themselves active catalysts for alkene isomerization. However, these species apparently can react with HSiEt_3 , a hydride source, to give mononuclear hydride catalysts for isomerization and alkene reduction. The enhanced catalytic activity, n-pentane formation, and formation of Si-containing products strongly implicate such a role for the HSiEt_3 . Irradiation of metal-metal bonded compounds in the presence of silicon hydrides is known to result in the

formation of silyl- and hydride complexes.⁸ Note that $[M_2(CO)_8H_2]$ (M=Mn, Re) can be formed via photolysis of $[M_2(CO)_{10}]$ in the presence of H_2 .⁹ Ir and uv-vis spectral changes in our system containing $HSiEt_3$ reveal conversion to mononuclear Co complexes, and catalytic activity persists when all metal-metal bonded compounds are exhausted. Molecular H_2 at 2 atm does appear to serve as a source of hydride for $[Co_2(CO)_6(P(OPh)_3)_2]$ but the catalytic activity in the presence of 1 M 1-pentene is not as great as that found with $HSiEt_3$. A priori H_2 should be less active as a hydride source, since the H-H bond is stronger than the Si-H bond. Whether mononuclear hydrides are responsible for reaction in the absence of H_2 or Si-H sources, Table I, is not clear. Our studies of this low activity situation have not been definitive to date.

Mononuclear Co-hydride complexes possibly arise from reaction of 15-valence electron Co-containing species¹⁰ with $HSiEt_3$ followed by hydrogen abstraction according to reactions (3)-(6), not unlike the processes proposed for the



photochemical formation of $[M_2(CO)_8H_2]$.^{9,11} That reaction (4) is crucial is consistent with our finding that 10 psi CO pressure effectively suppresses photocatalytic activity.

Hydrides like that indicated in equation (6) are known catalysts for alkene reactions.¹² Independent preparation¹³ of $[HCo(CO)_3P(OPh)_3]$ by addition of $HBF_4 \cdot Et_2O$ to $Na[Co(CO)_3P(OPh)_3]$ in tetrahydrofuran (THF) solution has been carried out, and we find thermal catalytic chemistry (isomerization and alkene reduction at 25°C in $HSiEt_3$ /1-pentene, 1.4 M each, in THF), and the distribution of products is similar to that found from irradiating $[Co_2(CO)_6(P(OPh)_3)_2]$ under the same conditions. We find that the $[HCo(CO)_3P(OPh)_3]$

has an intense ir absorption at 1992 cm^{-1} (THF), and that at 25°C the complex decomposes¹³ to form $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$ (1978 cm^{-1}) in THF solution. Further, the catalytic activity of our THF solutions of $[\text{HCo}(\text{CO})_3\text{P}(\text{OPh})_3]$ is enhanced (~factor of three in rate) by near-uv irradiation. This last result would indicate that our photocatalytic experiments beginning with $[\text{Co}_2(\text{CO})_6\text{L}_2]$ likely have a component of light-accelerated catalysis by mononuclear compounds. Consistent with this conclusion we find that ~15 min of irradiation of $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$ followed by dark reaction yields catalytic chemistry at a rate substantially lower than with continuous irradiation. Comparison of relative dark and light rates accords well with the dark vs. light rates when $[\text{HCo}(\text{CO})_3\text{P}(\text{OPh})_3]$ is used.

Preliminary results using 1/1 DSiEt_3 /1-pentene with $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$ as the photocatalyst reveals the rapid formation of HSiEt_3 (detected by pmr and gc mass spec) accompanying formation of pentane and cis- and trans-2-pentene. The deuterium is found in all of the hydrocarbons. This experiment confirms an important role for the Si-H(D) system and adds further to the understanding of the mechanism. Thus a Co species containing alkene, hydride, and the tri-alkylsilyl group is implicated in the catalytic cycle. Inasmuch as we do observe significant amounts of alkane product it is reasonable that a key species is in fact $[\text{H}_2\text{Co}(\text{CO})_2(\text{SiEt}_3)\text{L}]$ from oxidative addition of HSiEt_3 to $[\text{HCo}(\text{CO})_3\text{L}]$ or from disproportionation of the 17-valence electron species $[\text{HCo}(\text{CO})_2(\text{SiEt}_3)\text{L}]$ indicated in equation (5).

Data in Tables I and II reveal that the ordering of the activity of the three 17-valence electron radicals is $[\text{Co}(\text{CO})_3\text{P}(\text{OPh})_3] > [\text{Co}(\text{CO})_3\text{P}(\underline{n}\text{-Bu})_3] > [\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$. The lower thermal activity of the Fe-Co system compared to $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$ is logically a consequence of the more inert metal-metal bond. The effect of added $\text{P}(\text{OMe})_3$ on the activity of the $\text{P}(\underline{n}\text{-Bu})_3$ complex is in accord with the notion that the photogenerated $[\text{Co}(\text{CO})_3\text{L}]$ radicals are substitution labile,¹⁰ and the various substituted radicals will have a different reactivity towards HSiEt_3 .¹⁴ The activity of the systems studied do

not seem to be great under the conditions employed, but we note that the observed quantum yields (molecules of product/photon incident) are essentially unity for the alkene isomerization in the presence of HSiEt_3 using $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$. Increased light intensity increases the observed rates, and we find that the linear pentenes can be equilibrated to the thermodynamic ratio in <3 h for a neat 1/1 HSiEt_3 /1-pentene solution. The turnover rate thus exceeds $3 \times 10^2/\text{h}$, and may be much higher at higher intensities. If a 1/1 HSiEt_3 /1-pentene is exposed to 10 psi H_2 and illuminated at high intensities the turnover number for pentane formation exceeds $10^2/\text{h}$. Finally, at the high intensities, large yields of hydrosilation products are observable on the timescale of experiments detailed in Table II. Further data concerning these optical effects on turnover rate will be included in the full paper.

Acknowledgements.

We thank the Office of Naval Research for partial support of this research. MSW acknowledges support as a Dreyfus Teacher-Scholar grant recipient, 1975-1980.

Carol L. Reichel and Mark S. Wrighton*

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

References

1. Abrahamson, H. B.; Wrighton, M. S. Inorg. Chem., 1978, 17, 1003 and references cited therein.
2. (a) Parshall, G. W. J. Molecular Catalysis, 1978, 4, 243; (b) Paulik, F. E. Catal. Rev., 1972, 6, 49.
3. $[\text{Co}_2(\text{CO})_6\text{L}_2]$ complexes were synthesized and identified according to the literature procedure: Manning, A. R. J. Chem. Soc. (A), 1968, 1135. The Fe-Co species was prepared by near-uv irradiation of a mixture of $[\text{Co}_2(\text{CO})_6(\text{P}(\text{O}^i\text{Pr})_3)_2]$ and $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$ in benzene solution followed by chromatography on neutral alumina. Elemental analysis by Alfred Bernhardt, West Germany, was satisfactory; calcd. for $\text{C}_{28}\text{H}_{10}\text{O}_8\text{PFeCo}$: C, 53.36; H, 3.20; P, 4.91; found: C, 53.12; H, 3.30; P, 4.66. The compound melts with decomposition 140-142°C. The full characterization will appear in the full paper but ir evidence reveals no bridging CO's.
4. Wrighton, M. S.; Ginley, D. S. J. Am. Chem. Soc., 1975, 97, 2065.
5. (a) Giannotti, C.; Merle, G. J. Organometal. Chem., 1976, 105, 97; (b) Abrahamson, H. B.; Palazzotto, M. C.; Reichel, C. L.; Wrighton, M. S. submitted for publication.
6. Bond, G. C.; Hellier, M. J. Catal., 1965, 4, 1.
7. Putnik, C. F.; Welter, J. J.; Stucky, G. D.; D'Aniello, Jr., M. J.; Sosinsky, B. A.; Kirner, J. F.; Muetterties, E. L. J. Am. Chem. Soc., 1978, 100, 4107.
8. (a) Wrighton, M. Chem. Rev., 1974, 74, 401; (b) Hoyano, J. K.; Graham, W. A. G. Inorg. Chem., 1972, 11, 1265.
9. Byers, B. H.; Brown, T. L. J. Am. Chem. Soc., 1975, 97, 947 and 3260; 1977, 99, 2527.
10. Absi-Halabi, M.; Brown, T. L. J. Am. Chem. Soc., 1977, 99, 2982 provides evidence for $15e^-$ -Co species from $17e^-$ fragments.
11. The oxidative addition of H_2 to $\text{Co}(0)$ in the equilibrium $\text{H}_2 + \text{N}_2\text{Co}(\text{PPh}_3)_3 \rightleftharpoons \text{N}_2 + \text{H}_2\text{Co}(\text{PPh}_3)_3$, Speier, G.; Marko, L. Inorg. Chim. Acta, 1969, 3, 126, may be via $15e^-$ species. However, direct abstraction of H· from HSiEt_3 by the $17e^-$ radical may be possible if there is ligand redistribution to produce $[\text{Co}(\text{CO})_n\text{L}_{4-n}]$ species; see also Kidd, D. R. and Brown, T. L. J. Am. Chem. Soc., 1978, 100, 4103.
12. See: Taqui Khan, M. M.; Martell, A. E. "Homogeneous Catalysis by Metal Complexes", Vol. II, Academic Press, New York, 1974, for discussion of various catalytic reactions of alkenes using cobalt carbonyl precursors.

13. (a) Hieber, W.; Lindner, E. Chem. Ber., 1961, 94, 1417; (b) Wender, I.; Pino, P. "Organic Syntheses via Metal Carbonyls", Interscience, New York, 1968, pp. 188-189; (c) Piacenti, F.; Bianchi, M.; Benedetti, E. La Chimica e l'Industria, 1967, 49, 245.
14. A reviewer suggested that $[\text{CoL}_4(\text{olefin})]^+$ may represent an actual catalyst involved here. Small quantities of very active species are of course difficult to rule out, but $[\text{CoL}_4]^+$, though a reactive species, has a short lifetime and gives decomposition to $[\text{CoL}_5]^+$ and $\text{Co}(0)$ under catalytic conditions: Muetterties, E. L.; Watson, P. L. J. Am. Chem. Soc., 1978, 100, 6978. Note too that we find photocatalytic formation of 3,3-dimethylpentane when $[\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2]$ is irradiated in the presence of HSiEt_3 /3,3-dimethyl-1-pentene evidencing catalytic activity for alkenes incapable of easy π -allyl-hydride formation believed to be important in the $[\text{CoL}_4]^+$ catalyzed isomerization. Use of DSiEt_3 results in the formation of HSiEt_3 and deuterated hydrocarbon alkane and alkene product.

Table I. Photocatalyzed 1-Pentene Reaction Using Dinuclear Co Complexes as Catalyst Precursors.

Catalyst Precursor, (conc.)	Conditions ^a	Irrdn Time, h	Analysis, % ^b			
			n-pentane	1-pentene	trans-2-pentene cis-2-pentene	
[Co ₂ (CO) ₆ (P(<u>n</u> -Bu) ₃) ₂] (2.64 x 10 ⁻³ M)	3.6M 1-pentene in n-octane	0	---	99.7	0.20	0.03
		94	---	99.7	0.23	0.09
		[thermal control]	---	99.7	0.21	0.06
[Co ₂ (CO) ₆ (P(OPh) ₃) ₂] (8.97 x 10 ⁻⁴ M)	1.0M 1-pentene + 1.6 x 10 ⁻² M P(OMe) ₃ in n-octane	0	---	99.7	0.18	0.16
		72	0.13	95.5	2.69	1.72
		[thermal control]	---	99.6	0.22	0.18
[Co ₂ (CO) ₆ (P(OPh) ₃) ₂] (8.97 x 10 ⁻⁴ M)	1.0M 1-pentene in benzene	0	---	99.3	0.28	0.38
		76	0.28	92.0	5.05	2.65
		[thermal control]	---	99.3	0.40	0.30

^aAll experiments run at 25° C using 1.0 ml samples freeze-pump-thaw degassed in at least four cycles and hermetically sealed in Pyrex ampules. Irradiation was with a GE Blacklite with principal output at 355 nm ± 15 nm providing 1.6 x 10⁻⁶ ein/min incident on the sample.

^bAnalysis of C₅ mixture by vpc against n-hexane as an internal standard.

Table II. Photocatalyzed 1-Pentene Reaction in the Presence of HSiEt_3 .^a

Catalyst Precursor, (conc.)	Conditions	Irrdn Time, h	Analysis, %		
			n-pentane	1-pentene	trans-2-pentene cis-2-pentene
[$\text{Co}_2(\text{CO})_6(\text{P}(\underline{n}\text{-Bu})_3)_2$] ($2.65 \times 10^{-3}\text{M}$)	3.6M HSiEt_3 / 3.6M 1-pentene (neat)	0 94 [thermal control]	0.0 ₁ 3.1 ₀ 0.6 ₄	99.6 3.30 97.9	0.27 74.5 1.0 ₇ 0.1 ₄ 18.0 0.3 ₆
	1.0M HSiEt_3 1.0M 1-pentene in n-octane	52	0.93	94.0	3.61 1.51
	1.0M HSiEt_3 1.0M 1-pentene $5.3 \times 10^{-3}\text{M}$ P(OMe) ₃ in n-octane	52	5.44	16.5	58.7 19.4
[$\text{Co}_2(\text{CO})_6(\text{P}(\text{OPh})_3)_2$] ($8.97 \times 10^{-4}\text{M}$)	1.0M HSiEt_3 1.0M 1-pentene in benzene	0 5 [thermal control]	--- 3.07 0.1 ₂	99.6 46.0 97.8	0.28 37.0 1.29 0.20 13.9 0.78
	1.0M HSiEt_3 1.0M 1-pentene in benzene	0 7 [thermal control]	--- 2.74 0.0 ₂	99.6 7.46 99.1	0.22 76.2 0.6 ₅ 0.20 13.6 0.22
[$(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2\text{-}$ $\text{Co}(\text{CO})_3(\text{P}(\text{OPh})_3)]$ ($1.79 \times 10^{-3}\text{M}$)	3.6M HSiEt_3 3.6M 1-pentene (neat)	0 96 [thermal control]	0.0 ₄ 0.77 0.0 ₄	99.6 98.3 99.3	0.19 0.60 0.4 ₁ 0.1 ₄ 0.3 ₅ 0.25

^aSee notes for Table I.

For NR 051-579

472:GAN:716:tam
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217 Attn: Code 472	2	Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12
ONR Branch Office 536 S. Clark Street Chicago, Illinois 60605 Attn: Dr. George Sandoz	1	U.S. Army Research Office P.O. Box 1211 Research Triangle Park, N.C. 27709 Attn: CRD-AA-IP	1
ONR Branch Office 715 Broadway New York, New York 10003 Attn: Scientific Dept.	1	Naval Ocean Systems Center San Diego, California 92152 Attn: Mr. Joe McCartney	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. J. Marcus	1	Naval Weapons Center China Lake, California 93555 Attn: Dr. A. B. Amster Chemistry Division	1
ONR Area Office One Hallidie Plaza, Suite 601 San Francisco, California 94102 Attn: Dr. P. A. Miller	1	Naval Civil Engineering Laboratory Port Hueneme, California 93401 Attn: Dr. R. W. Drisko	1
ONR Branch Office Building 114, Section D 666 Summer Street Boston, Massachusetts 02210 Attn: Dr. L. H. Peebles	1	Professor K. E. Woehler Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Washington, D.C. 20390 Attn: Code 6100	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217 Attn: Dr. Richard S. Miller	1
Commander, Naval Air Systems Command Department of the Navy Washington, D.C. 20360 Attn: Code 310C (H. Rosenwasser)	1	Naval Ship Research and Development Center Annapolis, Maryland 21401 Attn: Dr. G. Bosmajian Applied Chemistry Division	1
		Naval Ocean Systems Center San Diego, California 91232 Attn: Dr. S. Yamamoto, Marine Sciences Division	1

Encl

TECHNICAL REPORT DISTRIBUTION LIST, 051A

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. M. A. El-Sayed University of California, Los Angeles Department of Chemistry Los Angeles, California 90024	1	Dr. M. Rauhut American Cyanamid Company Chemical Research Division Bound Brook, New Jersey 08805	1
Dr. M. W. Windsor Washington State University Department of Chemistry Pullman, Washington 99163	1	Dr. J. I. Zink University of California, Los Angeles Department of Chemistry Los Angeles, California 90024	1
Dr. E. R. Bernstein Colorado State University Department of Chemistry Fort Collins, Colorado 80521	1	Dr. B. Schechtman IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. J. R. MacDonald Naval Research Laboratory Chemistry Division Code 6110 Washington, D.C. 20375	1		
Dr. G. B. Schuster University of Illinois Chemistry Department Urbana, Illinois 61801	1		
Dr. E. M. Eyring University of Utah Department of Chemistry Salt Lake City, Utah	1		
Dr. A. Adamson University of Southern California Department of Chemistry Los Angeles, California 90007	1		
Dr. M. S. Wrighton Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1		