

AD-A074 339

CASE WESTERN RESERVE UNIV CLEVELAND OHIO DEPT OF MA--ETC F/6 12/1
CONFIDENCE INTERVALS FOR THE COMMON VARIANCE OF EQUICORRELATED --ETC(U)
AUG 79 P F RAMIG, S ZACKS

UNCLASSIFIED

TR-37


N00014-75-C-0529

NL

| OF |

AD
A074339





ADA 074339

LEVEL

10

CONFIDENCE INTERVALS FOR THE COMMON VARIANCE OF EQUICORRELATED NORMAL RANDOM VARIABLES.

by

P. F. Ramig and S. Zacks
Case Western Reserve University

Technical Report No. 37

1 August, 1979

TR

DDC
R
E
SEP 27 1979

33p.

PREPARED UNDER CONTRACT

N 00014-75-C-0529 PROJECT NR 042-276

OFFICE OF NAVAL RESEARCH

This document has been approved for public release and sale; its distribution is unlimited.

Reproduction in Whole or in Part is Permitted for any Purpose for the United States Government

Department of Mathematics and Statistics
CASE WESTERN RESERVE UNIVERSITY
Cleveland, Ohio

403901

79 09 26 065

DDC FILE COPY

1. INTRODUCTION

The pooling of variance estimates, obtained under different experimental conditions, is a classical problem in statistical inference. The commonly used formulas for pooling in the normal case are based on the assumption that the samples are independent and the population variances are equal or their ratios are known. Previous studies were mainly oriented to the pooling problem when the assumption about the equality of variances is questionable. However, little is known concerning the situation when the samples may not be independent. In the present study we address the problem of estimating the common variance of several normal populations which may be correlated. More specifically we consider a model of several independent replicas from a multivariate normal distribution having a covariance matrix of the form $\sigma^2 \underline{R}$ where \underline{R} is a correlation matrix for equicorrelated random variables, with unknown variance, σ^2 , and correlation coefficient, ρ . To simplify we present the development under the assumption that the means are known. Straightforward generalization to the case of unknown means can be given for practical applications by substituting their maximum likelihood estimates and modifying the number of degrees of freedom. However, the theoretical problem of admissibility of the maximum likelihood procedures, as indicated by Stein [7], arises and Stein type estimators can be attempted.

We investigate the problem of determining efficient confidence intervals for ρ and σ^2 . Confidence intervals for ρ were previously determined by Olkin and Pratt [6]. Nontrivial confidence intervals for σ^2 , when ρ is unknown, are more difficult to obtain. There exists no uniformly most

accurate invariant (UMAI) confidence interval for σ^2 since ρ is a nuisance parameter. In the present study we develop interval estimators for σ^2 , which are asymptotically most efficient, by employing the best asymptotic normality, (BAN), of the maximum likelihood estimators (see Zacks [11] p.244). The problem is to evaluate the coverage probabilities of these MLE based interval estimators in small or medium size samples. Exact coverage probabilities were determined by employing distributional properties of the statistics involved and some numerical integrations. The results indicate that the actual coverage probabilities of the intervals proposed are close to the nominal ones, when the number of replicas is at least 20. For smaller samples the coverage probabilities may show greater deviation from the nominal ones, especially when $|\rho|$ is close to one. We have therefore investigated the properties of alternative estimators in small sample situations.

When ρ is known one can construct UMAI confidence intervals for σ^2 . The statistic used in this case suggests a system of confidence intervals in which proper estimates are substituted for the unknown ρ . This type of intervals will be called "estimated- ρ intervals". In order to secure the required coverage probability one may attempt to apply the Bonferroni inequality and use confidence limits for ρ , rather than point estimates. We show that such an approach results in very inefficient confidence intervals. By trial and error the development of more efficient intervals of this type might be possible. More research should be performed in order to derive universally good estimated- ρ type confidence intervals. In addition, we

Accession For	NTIS GRAB
	DDC TAB
	Unannounced
	Justification
By	
Distribution/	
Availability Codes	
Avail and/or	
special	
Dit	A

compare the MLE based confidence intervals with "naive" type of intervals, which employs in a non-optimal fashion the information on σ^2 obtained from the individual samples. We show in the present paper that despite the slight small sample deficiencies of the MLE based confidence intervals, they are generally more efficient than the other types of intervals considered here. For this purpose we introduce a measure of relative efficiency which is based on the actual coverage probability and the expected length of the intervals.

The determination of the actual coverage probabilities and the expected length is generally a difficult matter. We provide a method for exact numerical determination in the case of the MLE based intervals. For the "naive" type intervals an explicit formula for the expected length has been derived. For evaluating the efficiency of the "estimated- ρ type intervals" we have employed the Monte Carlo method. Finally, we have investigated the loss in actual coverage incurred by ignoring the possibility of correlation and employing the usual confidence intervals for the case of $\rho = 0$.

There are many practical problems for which the above model of equicorrelated homoscedastic observations applies. For example, the compressive strength of concrete cubes from the same batch are generally correlated. Since a batch is homogeneously mixed, all the cubes from the same batch will have strength values with equal variances and the correlation between any pair of cubes will be the same. Cubes prepared from different batches will generally be independent and will have the same variance if the manufacturing conditions are similar.

A frequent application of the equicorrelated, equal variance model is outlined by Winer [9] for the one-way repeated measures design.

2. THE MODEL, THE LIKELIHOOD FUNCTION AND
THE FISHER INFORMATION

Let \underline{X} be an $m \times 1$ random vector having an equicorrelated multinormal distribution with zero mean vector and covariance matrix

$$\underline{\Sigma} = \sigma^2[(1-\rho)\underline{I} + \rho\underline{J}], \quad (2.1)$$

$0 < \sigma^2 < \infty$, $-(m-1)^{-1} < \rho < 1$; where $m \geq 2$, and $\underline{J} = \underline{1}\underline{1}'$ where $\underline{1}' = (1, 1, \dots, 1)$. Consider the Helmert transformation $\underline{Y} = \underline{H}\underline{X}$ where \underline{H} is an orthogonal matrix with first row vector equal to $m^{-\frac{1}{2}}\underline{1}'$ (see Kendall and Buckland [3] p.126). The components of \underline{Y} are independent normal random variables, with zero means and variances

$$\text{Var}(Y_j) = \begin{cases} \sigma^2(1 + (m-1)\rho) & j = 1 \\ \sigma^2(1 - \rho) & j = 2, \dots, m. \end{cases} \quad (2.2)$$

Given n independent identically distributed observations on $\underline{X}_1, \dots, \underline{X}_n$, the likelihood function of (σ^2, ρ) can be expressed in terms of the \underline{Y} vectors as

$$L(\sigma^2, \rho; \underline{Y}_1, \underline{Y}_2, \dots, \underline{Y}_n) = c(\sigma^2)^{-mn/2}(1 + (m-1)\rho)^{-n/2} \cdot \\ (1-\rho)^{-(m-1)n/2} \exp\{-(2\sigma^2)^{-1}[(1 + (m-1)\rho)^{-1} \sum_{i=1}^n Y_{i1}^2 \\ + (1-\rho)^{-1} \sum_{i=1}^n \sum_{j=2}^m Y_{ij}^2]\}. \quad (2.3)$$

Define $Q_1 = \sum_{i=1}^n Y_{i1}^2$ and $Q_2 = \sum_{i=1}^n \sum_{j=2}^m Y_{ij}^2$.

It follows that Q_1 and Q_2 are independent and constitute complete sufficient statistics for the present model (see Lehmann [4] p.132).

Furthermore, let $\chi^2[v]$ designate a chi-squared r.v. with v degrees of freedom, then

$$\begin{aligned} Q_1 &\sim \sigma^2(1 + (m-1)\rho) \chi^2[n] \\ \text{and} & \\ Q_2 &\sim \sigma^2(1-\rho) \chi^2 [n(m-1)]. \end{aligned} \tag{2.4}$$

The log-likelihood (sample information) function can then be expressed as

$$\begin{aligned} l(\sigma^2, \rho; Y_1, Y_2, \dots, Y_n) &= \log c - \frac{mn}{2} \log \sigma^2 - \frac{n}{2} \log(1 + (m-1)\rho) \\ &\quad - \frac{(m-1)n}{2} \log(1-\rho) - (2\sigma^2)^{-1} [(1 + (m-1)\rho)^{-1} Q_1 + (1-\rho)^{-1} Q_2]. \end{aligned} \tag{2.5}$$

The score functions are given by

$$\begin{aligned} S_1(\sigma^2, \rho) &= \frac{\partial l}{\partial \sigma^2} = -mn/(2\sigma^2) + Q_1/[2\sigma^4(1+(m-1)\rho)] + Q_2/[2\sigma^4(1-\rho)] \\ S_2(\sigma^2, \rho) &= \frac{\partial l}{\partial \rho} = -n(m-1)/[2(1+(m-1)\rho)] + n(m-1)/[2(1-\rho)] \\ &\quad + (m-1)Q_1/[2\sigma^2(1+(m-1)\rho)^2] - Q_2/[2\sigma^2(1-\rho)^2]. \end{aligned} \tag{2.6}$$

The maximum likelihood estimator of $\theta = (\sigma^2, \rho)$ is obtained from (2.6) as the roots of the system of equations $S_1(\sigma^2, \rho) = 0$; $S_2(\sigma^2, \rho) = 0$.

These MLE's are

$$\begin{aligned} \hat{\sigma}^2 &= (Q_1 + Q_2)/(mn) \\ \text{and} & \\ \hat{\rho} &= (Q_1 - (m-1)^{-1}Q_2)/(Q_1 + Q_2). \end{aligned} \tag{2.7}$$

Since the present model satisfies the Cramér-Rao regularity conditions (Wijsman [8]), $E_{\theta} \{S_1(\sigma^2, \rho)\} = 0$ and $E_{\theta} \{S_2(\sigma^2, \rho)\} = 0$ and the variances and covariances of the score functions are given by

$$\begin{aligned} \text{Var}(S_1) &= nm/(2\sigma^4), \\ \text{Var}(S_2) &= n(m-1)/2\{((m-1)(1-\rho)^2 + (1 + (m-1)\rho)^2) \cdot \\ &\quad ((1-\rho)^2(1 + (m-1)\rho)^2)^{-1}\} \\ \text{Cov}(S_1, S_2) &= -n(m-1)m\rho/[2\sigma^2(1-\rho)(1 + (m-1)\rho)]. \end{aligned} \quad (2.8)$$

Accordingly, the Fisher information matrix is

$$\underline{I}(\sigma^2, \rho) = \begin{bmatrix} \text{Var}(S_1) & \text{Cov}(S_1, S_2) \\ \cdot & \text{Var}(S_2) \end{bmatrix} \quad (2.9)$$

and its inverse is

$$\underline{I}^{-1}(\sigma^2, \rho) = \begin{bmatrix} 2\sigma^4(1 + (m-1)\rho^2)/(nm) & 2\sigma^2\rho(1-\rho)(1+(m-1)\rho)/(nm) \\ \cdot & 2(1-\rho)^2(1+(m-1)\rho)^2/(nm(m-1)) \end{bmatrix} \quad (2.10)$$

Hence the asymptotic variance of any asymptotically normal estimator of σ^2 is at least $2\sigma^4(1+(m-1)\rho^2)/(nm)$.

3. POINT AND INTERVAL ESTIMATORS BASED ON THE MLE

The MLE of σ^2 given in (2.7) is uniformly minimum variance unbiased (UMVU) having a variance

$$V(\hat{\sigma}^2) = 2\sigma^4(1+(m-1)\rho^2)/(mn). \quad (3.1)$$

This variance coincides with the asymptotic variance of the MLE, which is presented in $\underline{I}^{-1}(\sigma^2, \rho)$. Furthermore, $\hat{\sigma}^2$ is best asymptotically normal estimator (BAN).

Since Q_1 and Q_2 have gamma distributions with different scale parameters, their sum does not have a gamma distribution. The exact distribution of $\frac{\hat{\Lambda}^2}{\sigma^2}$ can be expressed as a mixture of gamma distributions (see Neuts and Zacks [5]). This distribution is, however, too complicated to provide a simple system of confidence intervals when ρ is unknown. We therefore construct interval estimators, which are based on the asymptotic normality of the MLE's with the aim of attaining a $(1 - \alpha)$ coverage probability. These interval estimators are based on the fact that when n is large then

$$P\left(\left|\frac{\hat{\Lambda}^2 - \sigma^2}{\sigma^2 \delta(\hat{\rho})}\right| \leq z_\gamma\right) \approx 1 - \alpha \quad (3.2)$$

where $\gamma = 1 - \alpha/2$, z_γ is the γ -fractile of the standard normal distribution and

$$\delta(\hat{\rho}) = [2(1 + (m-1)\hat{\rho}^2)/mn]^{\frac{1}{2}}. \quad (3.3)$$

From (3.2) one obtains the system of interval estimators of the form $(\hat{\Lambda}^2/b_\gamma(\hat{\rho}), \hat{\Lambda}^2/a_\gamma(\hat{\rho}))$, where $a_\gamma(\hat{\rho}) = \max(0, 1 - z_\gamma \delta(\hat{\rho}))$ and $b_\gamma(\hat{\rho}) = 1 + z_\gamma \delta(\hat{\rho})$. Notice that in small samples $z_\gamma \delta(\hat{\rho})$ could exceed one with positive probability. This is not the problem in large samples since, as follows from (3.3), $\delta(\hat{\rho}) \leq (2/n)^{\frac{1}{2}}$. Thus, in small samples, the above system may have a very large or infinite expected length. In order to overcome this difficulty, we propose a modified system of MLE based interval estimators, namely $(\hat{\Lambda}^2_{A_\gamma(\hat{\rho})}, \hat{\Lambda}^2_{B_\gamma(\hat{\rho})})$, where $A_\gamma(\hat{\rho}) = \max(0, 1 - t_\gamma[v_{\hat{\rho}}] \delta(\hat{\rho}))$ and $B_\gamma(\hat{\rho}) = 1 + t_\gamma[v_{\hat{\rho}}] \delta(\hat{\rho})$. $t_\gamma[v_{\hat{\rho}}]$ is the γ -fractile of the t -distribution with $v_{\hat{\rho}}$ degrees of freedom where $v_{\hat{\rho}}$ is taken as the integer part of $n(1 + (m-1)(1-\hat{\rho}^2))$. We have replaced z_γ by $t_\gamma[v_{\hat{\rho}}]$ in order to compensate for interchanging the roles of $\hat{\Lambda}^2$ and σ^2 in (3.2). Thus, the

MLE based confidence intervals are of the form

$$(\hat{\sigma}^2/A_Y^*(\hat{\rho}), \hat{\sigma}^2/B_Y^*(\hat{\rho})) \quad (3.4)$$

where

$$A_Y^*(\hat{\rho}) = \begin{cases} b_Y(\hat{\rho}) & , \quad \text{MLE intervals} \\ 1/A_Y(\hat{\rho}) & , \quad \text{Modified-MLE intervals} \end{cases} \quad (3.5)$$

$$B_Y^*(\hat{\rho}) = \begin{cases} a_Y(\hat{\rho}) & , \quad \text{MLE intervals} \\ 1/B_Y(\hat{\rho}) & , \quad \text{Modified-MLE intervals.} \end{cases} \quad (3.6)$$

We conclude the present section with a comment on confidence intervals for ρ . From (2.4) it is immediately implied that

$$\frac{Q_1}{Q_2} \sim \frac{1+(m-1)\rho}{1-\rho} \cdot \frac{n}{n(m-1)} F[n, n(m-1)] \quad (3.7)$$

where $F[v_1, v_2]$ designates an F-statistic with v_1 and v_2 degrees of freedom. UMAI confidence interval for ρ can be directly obtained from (3.7) (see Lehmann [4]).

4. DETERMINATION OF THE EXACT COVERAGE PROBABILITY AND EXPECTED LENGTH OF THE MLE BASED INTERVALS

From (2.4) and the independence of Q_1 and Q_2 one deduces that

$$\frac{Q_1/(1+(m-1)\rho)}{Q_1/(1+(m-1)\rho) + Q_2/(1-\rho)} \sim \beta(n/2, n(m-1)/2) \quad (4.1)$$

where $\beta(p, q)$ designates a random variable having a beta distribution with

parameters p, q . For the purpose of clearly distinguishing between p and \hat{p} in what follows, we replace \hat{p} by R . According to (2.7)

$$Q_1 / (Q_1 + Q_2) = (1 + (m-1)R) / m \quad (4.2)$$

and it follows that the left hand side of (4.1) is equivalent to

$$\varphi(R; \rho) = \frac{(1-\rho)(1 + (m-1)R)}{(1-\rho)(1 + (m-1)R) + (1 + (m-1)\rho)(1-R)} \quad (4.3)$$

Accordingly, the probability distribution of R , for each ρ , can be determined from (4.1) and (4.3). In addition, it is easy to verify that $P(R < -(m-1)^{-1}) = 0$.

Consider the random variable

$$Q(\rho) = (1 + (m-1)\rho)^{-1} Q_1 + (1-\rho)^{-1} Q_2.$$

The distribution of $Q(\rho)$ is like that of $\sigma^2 \chi^2[mn]$.

Algebraic manipulations lead to the expression

$$Q(\rho) = \frac{Q_1 + Q_2}{mn} \cdot \left\{ \frac{n[(1-\rho)(1 + (m-1)R) + (m-1)(1-R)(1 + (m-1)\rho)]}{(1-\rho)(1 + (m-1)\rho)} \right\} \\ = \frac{\Lambda^2}{\sigma^2} \Psi(R; \rho). \quad (4.4)$$

That is, $Q(\rho)$ has been factored into the product of the MLE $\frac{\Lambda^2}{\sigma^2}$ and the function $\Psi(R; \rho)$. (Notice that for $R \geq -(m-1)^{-1}$, $\Psi(R; \rho) \geq 0$.)

We show now that R is independent of $Q(\rho)$. Fix ρ and consider the subfamily of distributions of (Q_1, Q_2) , \mathcal{F}_ρ , depending on the scale parameter σ^2 . The random variable $Q(\rho)$ is a complete sufficient statistic for \mathcal{F}_ρ . By Basu's theorem [1] $Q(\rho)$ and R are independent since R is ancillary (invariant with respect to the group of scale transformation).

Hence, $Q(\rho)$ is independent of $Y(R; \rho)$. It follows that the conditional distribution of $\hat{\sigma}^2 Y(R; \rho)$, given R , is like that of the marginal distribution of $Q(\rho)$, namely that of $\sigma^2 \chi^2[nm]$. Thus, given $R = r$, $\hat{\sigma}^2$ is distributed like $(\sigma^2/Y(r; \rho))\chi^2[nm]$. This result is the basis for the determination of the coverage probabilities and expected length.

The coverage probability, under σ^2 and ρ , of the intervals defined by (3.4) is

$$P_{\sigma^2, \rho} \{ \hat{\sigma}^2/A_Y^*(R) \leq \sigma^2 \leq \hat{\sigma}^2/B_Y^*(R) \} = \quad (4.5)$$

$$P_{\sigma^2, \rho} \{ \sigma^2 B_Y^*(R) \leq \hat{\sigma}^2 \leq \sigma^2 A_Y^*(R) \}.$$

This coverage probability, CP, can be determined according to the previous result and the law of iterated expectations by

$$CP = E\{P_{\sigma^2, \rho} \{ \sigma^2 B_Y^*(R) \leq \hat{\sigma}^2 \leq \sigma^2 A_Y^*(R) | R \} \} \quad (4.6)$$

$$= E\{P_{\rho} \{ Y(R; \rho) B_Y^*(R) \leq \chi^2[nm] \leq Y(R; \rho) A_Y^*(R) | R \} \}.$$

Notice that the coverage probability is a function of ρ only.

Let $Pos(k|\lambda)$ designate the c.d.f. of a Poisson distribution with mean λ . If nm is an even integer, the c.d.f. of the distribution of $\chi^2[nm]$ can be determined by $Pos(k|\lambda)$ according to the well known relationship

$$P\{\chi^2[nm] \geq x\} = Pos(nm/2 - 1 | x/2). \quad (4.7)$$

Hence,

$$P\{\sigma^2 B_Y^*(R) \leq \hat{\sigma}^2 \leq \sigma^2 A_Y^*(R) | R\} =$$

$$Pos(nm/2 - 1 | \lambda_1(r; \rho)) - Pos(nm/2 - 1 | \lambda_2(r; \rho)), \quad (4.8)$$

where

$$\lambda_1(r; \rho) = \frac{1}{2} B_Y^*(r) \Psi(r; \rho)$$

and

(4.9)

$$\lambda_2(r; \rho) = \frac{1}{2} A_Y^*(r) \Psi(r; \rho).$$

As shown earlier, the function $\varphi(R; \rho)$, given by (4.3), is distributed like $\beta(n/2, n(m-1)/2)$. Since $\varphi(R; \rho)$ is a strictly increasing function of R over the interval $(-(m-1)^{-1}, 1)$,

$$\begin{aligned} P_\rho \{r' \leq R \leq r''\} &= P_\rho \{\varphi(r'; \rho) \leq \varphi(R; \rho) \leq \varphi(r''; \rho)\} \\ &= I_{\varphi(r''; \rho)}^{(n/2, n(m-1)/2)} - \\ &\quad I_{\varphi(r'; \rho)}^{(n/2, n(m-1)/2)}, \end{aligned} \quad (4.10)$$

for any $-(m-1)^{-1} \leq r' \leq r'' \leq 1$; where $I_a(p, q)$ is the incomplete beta function.

To evaluate the coverage probability (4.6) exactly one needs to integrate (4.8) with respect to the distribution of R over $(-(m-1)^{-1}, 1)$. We approximate this integral by partitioning the interval $(-(m-1)^{-1}, 1)$ into a large number, k , of subintervals (r_{i-1}, r_i) , $i=1, 2, \dots, k$; where $r_0 = -(m-1)^{-1}$; $r_k = 1$. The CP function is approximated then by

$$\begin{aligned} CP \cong \sum_{i=1}^k & [\text{Pos}(mn/2 - 1 \mid \lambda_1(r_i^*; \rho)) - \text{Pos}(mn/2 - 1 \mid \lambda_2(r_i^*; \rho))] \cdot \\ & [I_{\varphi(r_i; \rho)}^{(n/2, n(m-1)/2)} - I_{\varphi(r_{i-1}; \rho)}^{(n/2, n(m-1)/2)}] \end{aligned} \quad (4.11)$$

where r_i^* is suitably chosen in (r_{i-1}, r_i) . In Table 1 and Table 2 we present the CP values of the MLE and Modified-MLE intervals. The

computations were performed with $k=100$ equal size subintervals with $r_i^* = (r_i + r_{i-1})/2$. The incomplete beta function was approximated using a Fourier series expansion (see Woods and Posten [10]).

The determination of the expected length of the MLE based intervals proceeds along similar lines. Let $L(R, \gamma)$ designate the length of such an interval having nominal coverage probability γ , i.e.

$$L(R, \gamma) = \sigma^2 \left(\frac{1}{B_\gamma^*(R)} - \frac{1}{A_\gamma^*(R)} \right). \quad (4.12)$$

The conditional expectation of $L(R, \gamma)$, given R , is

$$E\{L(R, \gamma) \mid R\} = \sigma^2 \frac{mn}{\Psi(R; \rho)} \left[\frac{1}{B_\gamma^*(R)} - \frac{1}{A_\gamma^*(R)} \right]. \quad (4.13)$$

Finally, the expected length of the MLE based intervals is approximated, similar to (4.11), by

$$E\{L(R, \gamma)\} = \sigma^2 mn \sum_{i=1}^k \frac{A_\gamma^*(r_i^*) - B_\gamma^*(r_i^*)}{\Psi(r_i^*; \rho) B_\gamma^*(r_i^*) A_\gamma^*(r_i^*)} \cdot \quad (4.14)$$

$$\left[I_{\varphi(r_i; \rho)}\left(\frac{n}{2}, \frac{n(m-1)}{2}\right) - I_{\varphi(r_{i-1}; \rho)}\left(\frac{n}{2}, \frac{n(m-1)}{2}\right) \right]$$

Expected lengths of MLE based intervals for $\sigma^2 = 1$, $m = 2$, and varying values of n , ρ and γ are given in Table 3.

5. ESTIMATED $-\rho$ INTERVALS

When ρ is known one can construct, on the basis of the statistic $Q(\rho)$, a system of UMAI confidence intervals for σ^2 at level $(1-\alpha)$.

These intervals are of the form

$$\left(\frac{Q(\rho)}{\chi_{1-\epsilon_1}^2 [mn]}, \frac{Q(\rho)}{\chi_{\epsilon_2}^2 [mn]} \right) \quad (5.1)$$

where ϵ_1, ϵ_2 are determined so that $\epsilon_1 + \epsilon_2 = \alpha$ and an additional condition is satisfied (see Lehmann [4]). In practice, equal tail intervals are employed. Estimated $-\rho$ type intervals are obtained from (5.1) by substituting suitable estimators of ρ . If the MLE $\hat{\rho}$ is substituted, one obtains the UMAI confidence intervals for the case of $\rho = 0$ (known). As shown later, this results in a loss of coverage probability when $|\rho|$ is close to 1. Another approach is based on the Bonferroni inequality

$$P_{\rho, \sigma^2} \left\{ L_\alpha \frac{1-\rho \hat{\rho}}{1-\rho^2} \leq \sigma^2 \leq U_\alpha \frac{1-\rho \hat{\rho}}{1-\rho^2}, \rho_\alpha \leq \rho \leq \bar{\rho}_\alpha \right\} \geq 1 - \alpha \quad (5.2)$$

for all (σ^2, ρ) , where

$$L_\alpha = \frac{mn\sigma^{\wedge 2}}{\chi_{1-\alpha/4}^2 [mn]}, \quad U_\alpha = \frac{mn\sigma^{\wedge 2}}{\chi_{\alpha/4}^2 [mn]} \quad (5.3)$$

and $\rho_\alpha, \bar{\rho}_\alpha$ are the lower and upper $(1-\alpha/2)$ confidence limits for ρ as obtained from (3.7). Consider the function

$$f(\rho; \hat{\rho}) = (1 - \rho \hat{\rho}) / (1 - \rho^2), \quad -(m-1)^{-1} < \rho < 1. \quad (5.4)$$

The function $f(\rho; \hat{\rho})$ is convex in ρ , $\lim_{\rho \rightarrow 1} f(\rho; \hat{\rho}) = \infty$,

$$\lim_{\rho \rightarrow -(m-1)^{-1}} f(\rho; \hat{\rho}) = \begin{cases} \infty & m=2 \\ \frac{(m-1)(m-1+\hat{\rho})}{m(m-2)} & m \geq 3 \end{cases}$$

and attains the minimum of $f_0(\hat{\rho}) = \frac{1}{2}(1 + (1-\hat{\rho}^2)^{\frac{1}{2}})$ at $\rho_0 = (1 - (1-\hat{\rho}^2)^{\frac{1}{2}}) / \hat{\rho}$. Note that if $\hat{\rho} = 0$, $\rho_0 = 0$. Furthermore, if $\hat{\rho} \geq 0$ then $\rho_0 \leq \hat{\rho}$, and vice versa. Inequality (5.2) prescribes a $(1-\alpha)$ level simultaneous

confidence region, $C_\alpha(\hat{\sigma}^2, \hat{\rho})$, for (σ^2, ρ) . The boundaries of this region as functions of ρ are (see Figure 1)

$$\begin{aligned} b_1(\rho) &= L_\alpha f(\rho; \hat{\rho}), \\ b_2(\rho) &= U_\alpha f(\rho; \hat{\rho}), \\ b_3(\rho) &= \underline{\rho}_\alpha, \\ b_4(\rho) &= \bar{\rho}_\alpha. \end{aligned} \tag{5.5}$$

A confidence interval for σ^2 , called the BF-interval, can be obtained by projecting $C_\alpha(\hat{\sigma}^2, \hat{\rho})$ on the σ^2 -axis. This projection yields the upper confidence limit

$$\bar{\sigma}_\alpha^2 = \begin{cases} U_\alpha f(\bar{\rho}_\alpha; \hat{\rho}), & \hat{\rho} \geq 0 \\ U_\alpha f(\underline{\rho}_\alpha; \hat{\rho}), & \hat{\rho} < 0 \end{cases} \tag{5.6}$$

and the lower confidence limit

$$\underline{\sigma}_\alpha^2 = L_\alpha g_\alpha(\hat{\rho}) \tag{5.7}$$

where

$$g_\alpha(\hat{\rho}) = \begin{cases} I(\underline{\rho}_\alpha \geq \rho_0) f(\underline{\rho}_\alpha; \rho) + I(\underline{\rho}_\alpha < \rho_0) f_0(\hat{\rho}), & \hat{\rho} \geq 0 \\ I(\bar{\rho}_\alpha \geq \rho_0) f_0(\hat{\rho}) + I(\bar{\rho}_\alpha < \rho_0) f(\bar{\rho}_\alpha; \hat{\rho}), & \hat{\rho} < 0 \end{cases} \tag{5.8}$$

and $I\{A\}$ is the indicator function of the set A . Notice that the intervals of σ^2 values, $(b_1(\hat{\rho}), b_2(\hat{\rho}))$ and $(b_1(0), b_2(0))$ coincide.

Furthermore, the BF-interval $(\underline{\sigma}_\alpha^2, \bar{\sigma}_\alpha^2)$ contains $(b_1(0), b_2(0))$.

The BF-intervals are, however, generally too conservative in the sense that their coverage probabilities are larger than $(1 - \alpha)$. Moreover, the BF-intervals could be considerably skewed around $\hat{\sigma}^2$ in cases of small n and $|\rho|$ close to 1. In Table 4 we present a few simulation estimates of

the coverage probabilities (CP) and the expected length (EL) of the (non-modified) MLE intervals and the BF-intervals for $1 - \alpha = .90$, $m = 2$, $n = 10$ and several values of ρ . These simulation estimates are based on 100 independent and identical replicas for the case $\sigma^2 = 1$.

We remark that the CP and the EL of these intervals can be determined exactly according to the method discussed in the previous section. However, a small scale simulation was employed for the purpose of obtaining preliminary estimates only. The above estimates indicate that, indeed, the BF-interval provides higher coverage than desired and is very inefficient compared to the MLE intervals with respect to the expected length. An extensive evaluation of the relative efficiency of the MLE intervals will be given later.

6. NAIVE INTERVALS FOR σ^2 AT $m = 2$.

We discuss in the present section another attempt to derive a system of confidence intervals for σ^2 , which may yield in small samples and for values of $|\rho|$ close to 1 more efficient results. The investigation is restricted to the case of $m = 2$. Accordingly, we consider n i.i.d. vectors (X_{1i}, X_{2i}) , $i=1,2,\dots,n$, having a joint bivariate normal distribution, with zero means, common variance σ^2 and correlation ρ .

$$\text{Let } S_1 = \sum_{i=1}^n X_{1i}^2, \quad S_2 = \sum_{i=1}^n X_{2i}^2, \quad \text{and} \quad S_{12} = \sum_{i=1}^n X_{1i} X_{2i}.$$

$(S_1 + S_2, S_{12})$ is a minimal sufficient statistic. A confidence interval for σ^2 , at level $1 - \alpha/2$, based on S_j only, $j = 1, 2$ is given by $(S_j / X_{1-\alpha/4}^2[n], S_j / X_{\alpha/4}^2[n])$. From the Bonferroni inequality we imply that

the intersection (if not empty) of these two intervals is a confidence interval for σ^2 at level not less than $(1 - \alpha)$. In other words, we consider the interval

$$\left(\frac{\max(S_1, S_2)}{\chi^2_{1-\alpha/4}[n]}, \frac{\min(S_1, S_2)}{\chi^2_{\alpha/4}[n]} \right). \quad (6.1)$$

The interval (6.1) is empty, whenever $\max(S_1, S_2) \geq (\chi^2_{1-\alpha/4}[n]/\chi^2_{\alpha/4}[n]) \cdot \min(S_1, S_2)$. The probability of this event is generally negligible. Even in cases of $\rho \approx 0$ and $\alpha \approx .20$ the probability that (6.1) is empty does not exceed .01 for all n . (The computation of this probability is based on a formula given in Johnson & Kotz [2] p.222). For completeness we define the interval estimator when (6.1) is empty to be $(S_1/\chi^2_{1-\alpha/2}[n], S_1/\chi^2_{\alpha/2}[n])$. Since (6.1) is not a function of the minimal sufficient statistic its efficiency may be improved. There is reason to expect it to be efficient, however, when $|\rho| \approx 1$. We therefore develop a formula for determining the expected length of (6.1) in order to compare its efficiency with that of the MLE based intervals given by (3.4).

We develop an approximate formula for the expected length of the interval estimator, disregarding the event that (6.1) is empty. This approximation may lead to an error which is bounded by 10^{-2} in the cases under consideration.

Let $V = \min(S_1, S_2)$ and $U = \max(S_1, S_2)$. Since $U + V = S_1 + S_2$ the expected length of (6.1) is

$$EL = \frac{2n \sigma^2 - E\{U\}}{\chi^2_{\alpha/4}[n]} - \frac{E\{U\}}{\chi^2_{1-\alpha/4}[n]} \quad (6.2)$$

$$= \frac{2n \sigma^2}{\chi^2_{\alpha/4}[n]} - \left(\frac{1}{\chi^2_{\alpha/4}[n]} + \frac{1}{\chi^2_{1-\alpha/4}[n]} \right) E(U)$$

Let $F(\eta; \rho, \sigma^2)$ be the c.d.f. of U under (σ^2, ρ) . It can be shown that (see Johnson & Kotz [2])

$$F(\eta; \rho, \sigma^2) = P_{\rho, \sigma^2}[S_1 \leq \eta, S_2 \leq \eta] \tag{6.3}$$

$$= \sum_{j=0}^{\infty} g(j|\rho^2, \frac{n}{2}) [P\{\chi^2[n+2j] \leq \eta|(1-\rho^2)\}]^2$$

where

$$g(j|\rho^2, \frac{n}{2}) = \frac{\Gamma(\frac{n}{2} + j)}{\Gamma(\frac{n}{2}) \Gamma(j+1)} \rho^{2j} (1-\rho^2)^{n/2}, j = 0, 1, \dots \tag{6.4}$$

is the p.d.f. of a negative-binominal distribution. The expected value of U can be obtained by evaluating $\int_0^{\infty} [1 - F(\eta; \rho, \sigma^2)] d\eta$. After several manipulations one obtains for even sample size n the formula.

$$E(U) = 2n-2(1-\rho^2) \sum_{j=0}^{\infty} g(j|\rho^2, \frac{n}{2}) \cdot \sum_{k=1}^{\frac{n}{2} + j} G(\frac{n}{2} + j - 1 | \frac{1}{2}, k), \tag{6.5}$$

where $G(\cdot | \nu, \nu)$ designates the c.d.f. corresponding to $g(\cdot | \nu, \nu)$.

7. THE LOSS OF COVERAGE IN ASSUMING $\rho = 0$.

Suppose the investigator ignores the possibility of correlation.

Under the assumption $\rho = 0$ a confidence interval for σ^2 could be obtained according to the distribution of $Q_1 + Q_2$ which is like that of $\sigma^2 \chi^2[mn]$.

The upper and lower limits, respectively, of a $1-\alpha$ confidence interval are

$$\frac{Q_1 + Q_2}{\chi^2_{\alpha/2}[mn]} \quad \text{and} \quad \frac{Q_1 + Q_2}{\chi^2_{1-\alpha/2}[mn]} \tag{7.1}$$

However, if the assumption $\rho = 0$ is incorrect the nominal coverage probability is unlikely to be attained. For the case where ρ is close to zero, actual coverage will be close to the nominal and thus the "cost of ignorance" is not great. But, for $|\rho|$ near unity actual coverage may be considerably less than the nominal resulting in substantial loss of coverage. In order to investigate the extent of loss in coverage probability when ρ is not zero we computed the actual coverage for $\rho = .1, .5, \text{ and } .9$ under varying nominal coverage, γ , and different m and n .

The actual coverage probability, AC, of the interval (7.1) is determined in a manner similar to that of determining the CP values of the MLE based intervals. Thus, for even mn ,

$$\begin{aligned}
 AC &= P \left\{ \frac{Q_1 + Q_2}{\chi^2_{1-\alpha/2}[mn]} \leq \sigma^2 \leq \frac{Q_1 + Q_2}{\chi^2_{\alpha/2}[mn]} \right\} \quad (7.2) \\
 &= E \left\{ P \left\{ \frac{\Psi(R; \rho) \chi^2_{\alpha/2}[mn]}{mn} \leq \chi^2[mn] \leq \frac{\Psi(R; \rho) \chi^2_{1-\alpha/2}[mn]}{mn} \mid R \right\} \right\} \\
 &\approx \sum_{i=1}^k \left[\text{Pos} \left(\frac{mn}{2} - 1 \mid \eta_1(r_i^*; \rho) \right) - \text{Pos} \left(\frac{mn}{2} - 1 \mid \eta_2(r_i^*; \rho) \right) \right] \cdot \\
 &\quad \left[I_{\varphi(r_i; \rho)} \left(\frac{n}{2}, \frac{n(m-1)}{2} \right) - I_{\varphi(r_{i-1}; \rho)} \left(\frac{n}{2}, \frac{n(m-1)}{2} \right) \right]
 \end{aligned}$$

where

$$\eta_1(r; \rho) = \frac{1}{2} \frac{\Psi(r; \rho)}{mn} \chi^2_{\alpha/2}[mn] \quad \text{and}$$

$$\eta_2(r; \rho) = \frac{1}{2} \frac{\Psi(r; \rho)}{mn} \chi^2_{1-\alpha/2}[mn].$$

AC values of (7.1) are given in Table 5.

8. THE RELATIVE EFFICIENCY OF INTERVAL ESTIMATORS

The efficiency of interval estimators is a function of two components: their coverage probabilities (compared to the nominal ones) and their expected length. If ρ is known, the UMAI confidence intervals (5.1) are most efficient in the class of all location invariant and scale preserving confidence intervals, in the sense that among all such intervals having at least γ coverage the UMAI intervals have minimal expected length. However, when ρ is unknown UMAI confidence intervals do not exist. Asymptotically, for large samples, the MLE based intervals are most efficient in a wide class of intervals. However, as seen in the tables, for relatively small samples the MLE based intervals are not always better than other competitors. In order to compare alternative interval estimators, especially in small samples cases, we propose the following measure of relative efficiency

$$RE = \frac{E\{Q(\rho)\} D(\tilde{\gamma}) (1 + (m-1)\rho^2)^{\frac{1}{2}}}{EL}, \quad (8.1)$$

where $\tilde{\gamma}$ is the actual coverage probability of the interval estimator,

$$D(\tilde{\gamma}) = (\chi^2_{(1-\tilde{\gamma})/2, [nm]})^{-1} - (\chi^2_{(1+\tilde{\gamma})/2, [nm]})^{-1} \quad (8.2)$$

and EL is its expected length. Notice that $E\{Q(\rho)\}D(\tilde{\gamma})$ is the expected length of (5.1) having the same coverage probability, $\tilde{\gamma}$, as the interval estimator under consideration. Finally since

$$\lim_{n \rightarrow \infty} \frac{E\{L(R, \gamma)\}}{E\{Q(\rho)\}D(\tilde{\gamma})} = (1 + (m-1)\rho^2)^{\frac{1}{2}} \quad (8.2)$$

the R.H.S. of (8.2) was introduced in the numerator of (8.1) in order

to provide a fair comparison. In Table 6 and Table 7 the relative efficiency values of the MLE and Modified MLE intervals are presented for various values of ρ , γ , n and m .

9. DISCUSSION AND CONCLUSION

In Table 1 we see that the coverage probabilities of the MLE intervals are uniformly close to the nominal ones. On the other hand, as shown in Table 2, the Modified-MLE intervals do exhibit deficient coverage, especially for large ρ values and small n . This deficiency is due to the modification which leads to shorter intervals. The loss in coverage that may result by ignoring the possibility of correlation is shown in Table 5 to be rather pronounced when ρ approaches 1 and m increases. For fixed m and ρ this loss of coverage is insensitive to n . The "naive" confidence interval (6.1) was designed to have coverage probability not less than the nominal one. It is therefore of interest to compare its expected length to that of the MLE estimators. This comparison is provided in Table 3. We see that for small n values and $\rho = .9$ the "naive" confidence interval may have a smaller expected length than the MLE interval. Such cases show areas of possible improvement over the MLE intervals. In Table 6 and Table 7 the relative efficiency, (8.1), of the MLE and Modified-MLE intervals is given. We see first that for small sample size ($n=10$) and large γ , the MLE interval may have infinite expected length. Furthermore, the relative efficiency of the Modified-MLE intervals is very close to 1 when $\rho = .1$ and decreases slowly as ρ approaches 1. The Modified-MLE intervals are considerably more efficient than the MLE intervals, despite

the fact that there is some loss in coverage.

In conclusion, confidence intervals ignoring the possibility of correlation should not be used. The estimated- ρ intervals discussed in Section 5 seem also to be inefficient. We do recommend the use of the Modified-MLE intervals (3.4). Although they may have some coverage deficiency in small samples these intervals are highly efficient. Finally, although the various tables present results for positive ρ values, similar results would be obtained for negative values of ρ .

REFERENCES

- [1] Basu, D. (1955). On statistics independent of a complete sufficient statistic. *Sankhyā*, 15, 377-380.
- [2] Johnson, N.L. and Kotz, S. (1972). *Distributions in Statistics: Continuous Multivariate Distributions*. New York: Wiley.
- [3] Kendall, M.G. and Buckland, W.R. (1957). *A Dictionary of Statistical Terms*. London: The International Statistical Institute, Oliver and Boyd.
- [4] Lehmann, E.L. (1959). *Testing Statistical Hypotheses*. New York: Wiley.
- [5] Neuts, M.F. and Zacks, S. (1967). On mixtures of χ^2 - and F-distributions which yield distributions of the same family. *Ann. Inst. Statist. Math.*, 19, 527-536.
- [6] Olkin, I. and Pratt, J.W. (1958). Unbiased estimation of certain correlation coefficients. *Ann. Math Statist.*, 29, 201-211.
- [7] Stein, C. (1964). Inadmissibility of the usual estimator for the variance of a normal distribution with unknown mean. *Ann. Inst. Statist. Math.*, 16, 155-160.
- [8] Wijsman, R.A. (1973). On the attainment of the Cramer-Rao lower bound. *Ann. of Statist.*, 1, 538-542.
- [9] Winer, B.J. (1971). *Statistical Principles in Experimental Design*. New York: McGraw-Hill.
- [10] Woods, J.D. and Posten, H.O. (1968). Fourier series and Chebyshev polynomials in statistical distribution theory. Research Report No. 37, University of Connecticut, Storrs, Connecticut.
- [11] Zacks, S. (1971). *The Theory of Statistical Inference*. New York: Wiley.




Figure 1 - Boundaries for simultaneous confidence region, $C_\alpha(\hat{\sigma}^2, \hat{\rho})$, for (σ^2, ρ) for case of $m=2$ and $\hat{\rho} \geq 0$.

TABLE 1 - Coverage probabilities of MLE intervals.

m	p		.10					.50					.90					
	n	v	.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99
2	10		.830	.923	.961	.988	.818	.916	.959	.988	.814	.911	.952	.983				
	20		.816	.912	.957	.985	.807	.906	.952	.986	.805	.904	.949	.984				
	30		.808	.904	.950	.986	.802	.900	.949	.987	.801	.900	.948	.985				
	40		.809	.903	.949	.987	.802	.901	.949	.987	.799	.899	.947	.986				
	50		.807	.906	.953	.988	.803	.902	.951	.988	.799	.899	.948	.986				
	60		.806	.905	.952	.988	.802	.902	.951	.989	.797	.898	.947	.987				
3	10		.822	.917	.959	.989	.807	.907	.955	.990	.809	.907	.950	.983				
	20		.808	.904	.950	.987	.800	.899	.948	.988	.802	.902	.948	.985				
	30		.808	.907	.953	.987	.801	.900	.949	.988	.801	.900	.948	.986				
	40		.806	.905	.952	.989	.800	.900	.950	.989	.799	.899	.948	.987				
	50		.805	.904	.952	.989	.800	.900	.950	.989	.793	.893	.948	.988				
	60		.804	.903	.952	.989	.800	.900	.950	.989	.796	.897	.947	.987				
4	10		.818	.914	.957	.986	.799	.899	.949	.988	.806	.903	.948	.983				
	20		.809	.905	.950	.987	.797	.897	.947	.988	.802	.901	.948	.985				
	30		.807	.905	.953	.989	.799	.898	.948	.989	.800	.900	.948	.986				
	40		.805	.904	.952	.989	.799	.899	.949	.989	.798	.899	.948	.987				
	50		.804	.903	.952	.989	.799	.899	.949	.989	.797	.897	.947	.987				
	60		.803	.903	.951	.989	.799	.899	.949	.989	.796	.896	.947	.987				
5	10		.812	.907	.952	.987	.792	.893	.945	.988	.804	.902	.947	.982				
	20		.808	.907	.953	.989	.796	.896	.946	.988	.802	.901	.948	.985				
	30		.806	.905	.952	.989	.797	.897	.947	.989	.800	.900	.948	.986				
	40		.804	.903	.952	.989	.798	.897	.948	.989	.798	.898	.948	.987				
	50		.804	.903	.952	.990	.798	.898	.948	.989	.797	.897	.947	.987				
	60		.803	.902	.951	.990	.798	.898	.948	.989	.795	.896	.946	.987				
6	10		.811	.906	.952	.987	.789	.890	.942	.987	.804	.902	.946	.982				
	20		.807	.906	.953	.989	.794	.894	.945	.988	.802	.901	.948	.985				
	30		.805	.904	.952	.989	.796	.896	.946	.988	.799	.899	.948	.986				
	40		.804	.903	.952	.990	.797	.897	.947	.988	.798	.898	.947	.987				
	50		.803	.902	.951	.990	.797	.897	.947	.989	.796	.897	.947	.987				
	60		.803	.902	.951	.990	.798	.897	.948	.989	.795	.896	.946	.987				

TABLE 2 - Coverage probabilities of Modified-MLE intervals.

m	P n \ y	.10					.50					.90					
		.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99
2	10	.816	.886	.921	.962	.804	.868	.904	.951	.793	.852	.889	.939	.810	.886	.923	.967
	20	.818	.899	.939	.977	.810	.891	.929	.970	.810	.886	.923	.967	.813	.898	.937	.976
	30	.813	.904	.946	.982	.810	.899	.939	.978	.813	.898	.937	.976	.814	.905	.944	.982
	40	.815	.910	.953	.985	.812	.904	.946	.982	.814	.905	.944	.982	.817	.909	.950	.986
	50	.815	.912	.957	.990	.814	.908	.951	.986	.817	.909	.950	.986	.817	.911	.953	.988
	60	.815	.913	.958	.991	.814	.910	.954	.988	.817	.911	.953	.988				
3	10	.815	.893	.929	.968	.800	.867	.903	.949	.790	.850	.886	.936	.806	.883	.920	.964
	20	.809	.900	.942	.979	.804	.888	.927	.968	.806	.883	.920	.964	.811	.896	.934	.975
	30	.810	.906	.952	.987	.807	.897	.938	.977	.811	.896	.934	.975	.812	.902	.942	.980
	40	.810	.908	.954	.989	.808	.901	.944	.981	.812	.902	.942	.980	.813	.905	.947	.983
	50	.810	.908	.955	.990	.809	.903	.947	.983	.813	.905	.947	.983				
	60	.810	.909	.955	.991	.809	.905	.949	.985	.813	.907	.950	.985				
4	10	.813	.893	.933	.972	.795	.865	.901	.947	.788	.848	.884	.933	.803	.881	.917	.961
	20	.808	.903	.947	.981	.801	.886	.925	.966	.803	.881	.917	.961	.808	.894	.932	.972
	30	.808	.905	.951	.988	.805	.895	.936	.975	.808	.894	.932	.972	.810	.899	.940	.978
	40	.808	.906	.953	.989	.806	.899	.941	.979	.810	.899	.940	.978	.810	.902	.944	.981
	50	.808	.906	.954	.990	.807	.901	.945	.982	.810	.902	.944	.981				
	60	.808	.907	.954	.990	.807	.902	.947	.983	.810	.904	.947	.983				
5	10	.807	.895	.936	.974	.793	.862	.898	.945	.786	.846	.882	.930	.802	.880	.916	.960
	20	.807	.903	.946	.986	.801	.885	.924	.965	.802	.880	.916	.960	.806	.892	.930	.971
	30	.807	.904	.946	.988	.803	.893	.934	.973	.806	.892	.930	.971	.807	.897	.938	.976
	40	.807	.905	.946	.989	.804	.897	.939	.977	.807	.897	.938	.976	.808	.900	.942	.980
	50	.807	.905	.953	.990	.805	.899	.943	.980	.808	.900	.942	.980				
	60	.807	.906	.953	.990	.805	.900	.945	.982	.807	.902	.945	.982				
6	10	.807	.896	.938	.976	.791	.860	.896	.943	.784	.845	.880	.928	.801	.878	.915	.958
	20	.807	.903	.949	.986	.802	.884	.922	.964	.801	.878	.915	.958	.804	.890	.929	.969
	30	.806	.904	.951	.988	.802	.891	.932	.972	.804	.890	.929	.969	.806	.895	.936	.975
	40	.806	.904	.952	.989	.803	.895	.938	.976	.806	.895	.936	.975	.806	.899	.941	.979
	50	.806	.905	.953	.990	.804	.898	.941	.979	.806	.899	.941	.979				
	60	.806	.905	.953	.990	.804	.899	.944	.981	.806	.899	.944	.981				

TABLE 3 - Expected length of the "Naive" and MLE-based intervals for $m=2, \sigma^2=1$.

γ	n	.10			.50			.90		
		Naive	MLE	M-MLE ^a	Naive	MLE	M-MLE	Naive	MLE	M-MLE
.80	10	1.241	1.038	.879	1.339	1.181	.960	1.665	1.559	1.162
	20	.772	.645	.610	.829	.723	.673	1.014	.907	.821
	30	.605	.506	.495	.648	.566	.548	.790	.700	.671
	40	.514	.430	.427	.550	.480	.474	.669	.590	.581
	50	.454	.381	.381	.486	.424	.423	.591	.519	.519
	60	.411	.345	.347	.440	.384	.386	.534	.469	.474
.90	10	1.736	1.559	1.145	1.850	1.858	1.253	2.230	2.769	1.531
	20	1.035	.883	.795	1.096	1.006	.878	1.295	1.314	1.082
	30	.801	.677	.644	.846	.764	.715	.995	.967	.883
	40	.676	.569	.556	.713	.639	.618	.837	.799	.765
	50	.595	.500	.496	.627	.561	.552	.735	.695	.684
	60	.538	.451	.452	.566	.505	.504	.663	.623	.625
.95	10	2.282	2.277	1.386	2.415	2.948	1.520	2.858	5.634	1.876
	20	1.296	1.131	.962	1.361	1.316	1.065	1.577	1.814	1.326
	30	.988	.843	.780	1.036	.963	.867	1.193	1.256	1.083
	40	.828	.700	.673	.867	.793	.750	.995	1.011	.938
	50	.726	.611	.601	.760	.689	.670	.871	.868	.839
	60	.654	.549	.547	.684	.618	.611	.783	.771	.765
.99	10	3.859	- ^b	1.888	4.052	-	2.208	4.695	-	2.325
	20	1.924	1.828	1.315	2.003	2.288	1.463	2.263	3.914	1.871
	30	1.413	1.248	1.066	1.467	1.476	1.191	1.645	2.116	1.528
	40	1.163	1.000	.920	1.206	1.160	1.030	1.347	1.569	1.323
	50	1.009	.856	.821	1.046	.983	.920	1.166	1.292	1.183
	60	.903	.760	.748	.935	.867	.839	1.041	1.120	1.080

a. M-MLE designates Modified MLE.

b. A dash, "-", indicates infinite expected length.

TABLE 4 - Simulation estimates of coverage probability (CP) and expected length (EL) of the MLE and BF intervals for $1-\alpha = .90$, $n=10$, $m=2$, 100 replicas.

Interval	$\rho=.10$		$\rho=.50$		$\rho=.90$	
	CP	EL	CP	EL	CP	EL
MLE	.91	1.574	.90	1.871	.88	2.799
BF	.95	2.599	.95	2.996	.96	3.658

TABLE 5 - Coverage probabilities of confidence intervals assuming $\rho=0$.

m	p	Y n	.10					.50					.90													
			.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99								
2	.798	10	.899	.949	.990	.752	.862	.924	.981	.657	.777	.855	.946	.798	10	.899	.949	.990	.752	.862	.924	.981	.657	.777	.855	.946
	.798	20	.898	.947	.987	.749	.860	.920	.977	.656	.775	.852	.943	.798	20	.898	.947	.987	.749	.860	.920	.977	.656	.775	.852	.943
	.794	30	.893	.944	.987	.748	.857	.918	.977	.655	.774	.850	.941	.794	30	.893	.944	.987	.748	.857	.918	.977	.655	.774	.850	.941
	.798	40	.897	.945	.988	.748	.858	.919	.977	.654	.773	.849	.941	.798	40	.897	.945	.988	.748	.858	.919	.977	.654	.773	.849	.941
	.798	50	.899	.949	.990	.749	.860	.921	.979	.654	.774	.851	.942	.798	50	.899	.949	.990	.749	.860	.921	.979	.654	.774	.851	.942
	.798	60	.899	.949	.990	.749	.859	.921	.979	.653	.773	.850	.941	.798	60	.899	.949	.990	.749	.859	.921	.979	.653	.773	.850	.941
3	.796	10	.897	.948	.989	.710	.826	.896	.968	.568	.687	.772	.889	.796	10	.897	.948	.989	.710	.826	.896	.968	.568	.687	.772	.889
	.792	20	.892	.943	.987	.706	.821	.890	.964	.567	.687	.770	.886	.792	20	.892	.943	.987	.706	.821	.890	.964	.567	.687	.770	.886
	.796	30	.897	.948	.988	.706	.823	.892	.965	.568	.687	.770	.886	.796	30	.897	.948	.988	.706	.823	.892	.965	.568	.687	.770	.886
	.796	40	.897	.948	.989	.706	.822	.892	.965	.567	.686	.770	.885	.796	40	.897	.948	.989	.706	.822	.892	.965	.567	.686	.770	.885
	.796	50	.897	.948	.989	.705	.822	.892	.965	.566	.685	.769	.884	.796	50	.897	.948	.989	.705	.822	.892	.965	.566	.685	.769	.884
	.796	60	.897	.948	.989	.705	.822	.891	.965	.565	.684	.768	.884	.796	60	.897	.948	.989	.705	.822	.891	.965	.565	.684	.768	.884
4	.794	10	.895	.945	.986	.672	.792	.867	.952	.506	.621	.706	.834	.794	10	.895	.945	.986	.672	.792	.867	.952	.506	.621	.706	.834
	.793	20	.894	.943	.987	.669	.788	.863	.950	.507	.621	.706	.833	.793	20	.894	.943	.987	.669	.788	.863	.950	.507	.621	.706	.833
	.793	30	.895	.947	.989	.669	.788	.864	.950	.507	.621	.706	.833	.793	30	.895	.947	.989	.669	.788	.864	.950	.507	.621	.706	.833
	.793	40	.895	.947	.989	.668	.788	.863	.950	.506	.621	.705	.832	.793	40	.895	.947	.989	.668	.788	.863	.950	.506	.621	.705	.832
	.793	50	.895	.947	.989	.668	.787	.863	.950	.506	.620	.704	.831	.793	50	.895	.947	.989	.668	.787	.863	.950	.506	.620	.704	.831
	.793	60	.895	.947	.989	.668	.787	.863	.949	.505	.619	.703	.830	.793	60	.895	.947	.989	.668	.787	.863	.949	.505	.619	.703	.830
5	.788	10	.889	.941	.986	.638	.760	.840	.936	.461	.570	.653	.786	.788	10	.889	.941	.986	.638	.760	.840	.936	.461	.570	.653	.786
	.791	20	.894	.946	.988	.637	.759	.838	.935	.463	.572	.655	.786	.791	20	.894	.946	.988	.637	.759	.838	.935	.463	.572	.655	.786
	.791	30	.894	.946	.988	.637	.757	.837	.934	.462	.571	.655	.785	.791	30	.894	.946	.988	.637	.757	.837	.934	.462	.571	.655	.785
	.791	40	.894	.946	.988	.636	.757	.836	.933	.462	.571	.654	.784	.791	40	.894	.946	.988	.636	.757	.836	.933	.462	.571	.654	.784
	.791	50	.893	.946	.988	.636	.756	.836	.933	.461	.570	.653	.783	.791	50	.893	.946	.988	.636	.756	.836	.933	.461	.570	.653	.783
	.791	60	.893	.946	.988	.636	.756	.835	.932	.460	.569	.652	.782	.791	60	.893	.946	.988	.636	.756	.835	.932	.460	.569	.652	.782
6	.787	10	.887	.940	.986	.611	.733	.816	.920	.427	.530	.612	.744	.787	10	.887	.940	.986	.611	.733	.816	.920	.427	.530	.612	.744
	.789	20	.892	.945	.988	.609	.731	.813	.918	.428	.532	.613	.745	.789	20	.892	.945	.988	.609	.731	.813	.918	.428	.532	.613	.745
	.789	30	.892	.945	.988	.608	.729	.811	.917	.428	.531	.612	.744	.789	30	.892	.945	.988	.608	.729	.811	.917	.428	.531	.612	.744
	.789	40	.892	.944	.988	.608	.729	.811	.916	.427	.531	.612	.743	.789	40	.892	.944	.988	.608	.729	.811	.916	.427	.531	.612	.743
	.789	50	.892	.944	.988	.608	.728	.810	.916	.426	.530	.611	.742	.789	50	.892	.944	.988	.608	.728	.810	.916	.426	.530	.611	.742
	.789	60	.892	.944	.988	.607	.728	.810	.915	.426	.529	.610	.741	.789	60	.892	.944	.988	.607	.728	.810	.915	.426	.529	.610	.741

TABLE 6 - Relative efficiency of MLE intervals.

n	p	.10					.50					.90					
		.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99	.80	.90	.95	.99
2	10	.945	.849	.712	a	.893	.771	.602	-	.807	.611	.364	-	.807	.611	.364	-
	20	.978	.940	.887	.684	.950	.897	.829	.620	.906	.821	.713	.424	.906	.821	.713	.424
	30	.979	.948	.909	.794	.962	.924	.880	.752	.924	.878	.809	.618	.924	.878	.809	.618
	40	.991	.963	.931	.845	.972	.945	.911	.815	.948	.905	.855	.712	.948	.905	.855	.712
	50	.993	.980	.959	.884	.981	.962	.937	.858	.957	.925	.885	.767	.957	.925	.885	.767
	60	.995	.984	.967	.904	.984	.968	.948	.884	.960	.934	.901	.803	.960	.934	.901	.803
3	10	.964	.908	.830	-	.884	.786	.664	-	.780	.589	.362	-	.780	.589	.362	-
	20	.977	.946	.906	.790	.940	.893	.839	.690	.890	.804	.700	.429	.890	.804	.700	.429
	30	.992	.976	.952	.860	.962	.933	.899	.802	.927	.870	.803	.620	.927	.870	.803	.620
	40	.994	.983	.966	.903	.972	.951	.926	.855	.942	.900	.850	.712	.942	.900	.850	.712
	50	.995	.987	.973	.924	.978	.961	.941	.885	.950	.917	.877	.766	.950	.917	.877	.766
	60	.996	.989	.978	.937	.981	.968	.951	.905	.954	.927	.894	.801	.954	.927	.894	.801
4	10	.973	.932	.875	-	.875	.784	.677	-	.765	.577	.360	-	.765	.577	.360	-
	20	.988	.962	.928	.841	.936	.894	.846	.722	.885	.797	.695	.431	.885	.797	.695	.431
	30	.993	.982	.965	.903	.960	.933	.902	.822	.921	.865	.798	.619	.921	.865	.798	.619
	40	.995	.987	.974	.928	.970	.950	.927	.868	.937	.895	.846	.710	.937	.895	.846	.710
	50	.996	.990	.980	.943	.976	.960	.942	.895	.946	.913	.873	.764	.946	.913	.873	.764
	60	.997	.991	.983	.953	.980	.966	.952	.912	.951	.923	.890	.798	.951	.923	.890	.798
5	10	.968	.929	.880	-	.866	.780	.682	-	.757	.571	.358	-	.757	.571	.358	-
	20	.990	.977	.955	.880	.937	.896	.852	.739	.881	.794	.693	.433	.881	.794	.693	.433
	30	.994	.985	.972	.923	.958	.931	.902	.829	.918	.861	.795	.618	.918	.861	.795	.618
	40	.996	.989	.979	.943	.968	.949	.927	.873	.935	.893	.843	.709	.935	.893	.843	.709
	50	.997	.991	.984	.955	.975	.959	.942	.898	.944	.910	.870	.762	.944	.910	.870	.762
	60	.997	.993	.986	.963	.979	.965	.951	.915	.949	.921	.888	.797	.949	.921	.888	.797
6	10	.972	.939	.898	-	.864	.780	.687	-	.753	.569	.358	-	.753	.569	.358	-
	20	.991	.980	.962	.901	.934	.894	.851	.744	.878	.791	.690	.433	.878	.791	.690	.433
	30	.994	.987	.976	.936	.956	.930	.902	.831	.916	.859	.792	.617	.916	.859	.792	.617
	40	.996	.991	.982	.953	.967	.947	.926	.874	.933	.890	.841	.708	.933	.890	.841	.708
	50	.997	.993	.986	.963	.974	.958	.941	.899	.943	.909	.868	.760	.943	.909	.868	.760
	60	.997	.994	.989	.969	.978	.965	.951	.916	.947	.920	.887	.795	.947	.920	.887	.795

a. A dash, -, indicates zero efficiency, due to infinite expected length.

TABLE 7 - Relative efficiency of Modified-MLE intervals.

m	p	.10					.50					.90				
		.80	.90	.95	.99	1.00	.80	.90	.95	.99	1.00	.80	.90	.95	.99	1.00
2	10	1.075 ^a	1.011	.947	.862	1.060	.971	.902	.832	1.027	.912	.834	.823			
	20	1.040	.999	.959	.875	1.030	.979	.922	.829	1.014	.942	.871	.763			
	30	1.015	.997	.966	.891	1.014	.983	.937	.848	1.002	.955	.898	.788			
	40	1.014	1.008	.982	.903	1.009	.988	.951	.866	.996	.963	.909	.809			
	50	1.011	1.007	.995	.950	1.009	.995	.967	.896	.997	.970	.926	.831			
	60	1.009	1.006	.996	.955	1.007	.996	.972	.905	.994	.971	.932	.843			
3	10	1.052	1.011	.955	.871	1.030	.944	.876	.786	.993	.880	.801	.768			
	20	1.015	.998	.968	.896	1.010	.967	.912	.820	.996	.926	.853	.744			
	30	1.012	1.008	.995	.949	1.007	.981	.939	.854	.995	.947	.884	.780			
	40	1.009	1.006	.996	.958	1.006	.986	.952	.874	.993	.957	.904	.803			
	50	1.007	1.005	.997	.964	1.004	.989	.960	.887	.990	.961	.916	.820			
	60	1.006	1.004	.998	.968	1.004	.990	.966	.898	.988	.964	.924	.832			
4	10	1.039	.998	.960	.880	1.008	.928	.860	.767	.978	.866	.788	.742			
	20	1.013	1.006	.985	.910	1.002	.959	.904	.812	.987	.917	.846	.738			
	30	1.009	1.006	.996	.958	1.003	.975	.932	.847	.990	.942	.880	.776			
	40	1.007	1.004	.997	.965	1.003	.981	.946	.867	.989	.952	.900	.800			
	50	1.005	1.004	.998	.971	1.002	.985	.955	.881	.987	.958	.913	.817			
	60	1.005	1.003	.998	.974	1.002	.987	.962	.892	.985	.961	.921	.830			
5	10	1.018	.996	.963	.888	.997	.917	.848	.754	.969	.859	.781	.728			
	20	1.010	1.006	.993	.951	1.000	.957	.901	.810	.986	.917	.845	.737			
	30	1.007	1.004	.996	.962	1.001	.971	.927	.841	.987	.939	.878	.774			
	40	1.005	1.003	.997	.970	1.001	.978	.942	.861	.987	.950	.898	.799			
	50	1.004	1.003	.998	.975	1.000	.982	.952	.877	.985	.956	.912	.817			
	60	1.004	1.002	.998	.978	1.000	.985	.959	.889	.983	.959	.921	.831			
6	10	1.015	.995	.965	.894	.989	.907	.838	.744	.964	.854	.776	.718			
	20	1.008	1.005	.993	.954	.998	.952	.895	.803	.983	.914	.843	.736			
	30	1.006	1.003	.996	.966	.999	.968	.923	.835	.985	.938	.877	.774			
	40	1.004	1.003	.997	.973	.999	.976	.939	.857	.985	.949	.898	.799			
	50	1.003	1.002	.998	.977	.999	.981	.949	.873	.984	.955	.911	.817			
	60	1.003	1.002	.998	.980	.999	.984	.957	.886	.982	.958	.921	.832			

a. Values of RE may exceed 1 owing to the definition of RE as well as computing error.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 37	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Confidence Intervals for the Common Variance of Equicorrelated Normal Random Variables.		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) P. F. Ramig and S. Zacks		8. CONTRACT OR GRANT NUMBER(s) N 00014-75-C-0529 PROJECT NR 042-276
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Mathematics and Statistics Case Western Reserve University Cleveland, Ohio		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS OFFICE OF NAVAL RESEARCH ARLINGTON, VIRGINIA 22217		12. REPORT DATE August 1, 1979
		13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Variance estimation; Equicorrelated normal variates; Maximum likelihood; Confidence intervals; Coverage probability.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Confidence intervals for the common variance of equicorrelated normal random variable are investigated. When the correlation coefficient is unknown no uniformly most accurate invariant confidence interval for the common variance exists. We develop interval estimators which are based on the asymptotically most efficient ones by employing the best asymptotic normality of the maximum likelihood estimators. In addition to their comparison with possible competitors, their small sample efficiency is		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

[20]

studied. A method for determining exact coverage probabilities and expected length is developed. As a result of these investigations we conclude that interval estimators based on maximum likelihood estimators are to be recommended.