

AD-A074 964

LETTERMAN ARMY INST OF RESEARCH SAN FRANCISCO CA
INVESTIGATION OF POSSIBLE ANTITHIAMIN PROPERTIES IN IRRADIATION--ETC(U)
AUG 79 E L MCGOWN, C M LEWIS, P P WARING

F/G 6/8

UNCLASSIFIED

LAIR-72

NL

| OF |


AD
A074964



END
DATE
FILMED

11-79

DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-2



12

LEVEL

DDC
RAPID
OCT 10 1979
RECEIVED
E

AD A 074964

INSTITUTE REPORT NO. 72

INVESTIGATION OF POSSIBLE ANTITHIAMIN PROPERTIES IN IRRADIATION STERILIZED CHICKEN


EVELYN L. McGOWN, PhD
CAROLYN M. LEWIS, MS
PAUL P. WARING, BS

DIVISION OF NUTRITION TECHNOLOGY

This document has been approved
for public release and sale; its
distribution is unlimited.

DDC FILE COPY

AUGUST 1979



LETTERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO CALIFORNIA 94129

Reproduction of this document in whole or in part is prohibited except with the permission of Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, DDC is authorized to reproduce the document for United States Government purposes.

Destroy this report when no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Revision of the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Research Council.

John M. ... 30 Aug 79
(Date)

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER LAIR Institute Report No. 72	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9
4. TITLE (and Subtitle) INVESTIGATION OF POSSIBLE ANTITHIAMIN PROPERTIES IN IRRADIATION STERILIZED CHICKEN		5. TYPE OF REPORT & PERIOD COVERED Final Report Sep 1978 - Feb 1979
7. AUTHOR(s) Evelyn L. McGown, PhD; Carolyn M. Lewis, MS; and Paul P. Waring, BS		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Division of Nutrition Technology (SGRD-ULN) Letterman Army Institute of Research Presidio of San Francisco, CA 94129		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick Frederick, MD 21701		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Project No. DL 47 Work Unit #008
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 1257		12. REPORT DATE August 1979
		13. NUMBER OF PAGES 47
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) THIS DOCUMENT HAS BEEN APPROVED FOR PUBLIC RELEASE AND SALE: ITS DISTRIBUTION IS UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 14 LAIR-72		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Irradiated Chicken, Antithiamin, antivitamin, thiamin, transketolase, thiamin pyrophosphate (TPP) effect		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Male and female rats (156 each) were made thiamin-deficient by feeding a semi-purified diet devoid of thiamin. They were then repleted with various test diets containing chicken which had been preserved by one of four methods: fro- zen, thermally processed, electron or gamma irradiated. All repletion diets contained carefully controlled (marginal or high) levels of thiamin. Recovery rates were monitored by growth (weight gain) and measurements of a thiamin- dependent blood enzyme (erythrocyte transketolase). The responses of rats fed		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

404912

LB

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. irradiated chicken were similar to those fed frozen or thermally processed chicken. No evidence was found of antithiamin substances in either gamma or electron irradiation sterilized chicken.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ABSTRACT

Male and female rats (156 each) were made thiamin-deficient by feeding a semi-purified diet devoid of thiamin. They were then repleted with various test diets containing chicken which had been preserved by one of four methods: frozen, thermally processed, electron or gamma irradiated. All repletion diets contained carefully controlled (marginal or high) levels of thiamin. Recovery rates were monitored by growth (weight gain) and measurements of a thiamin-dependent blood enzyme (erythrocyte transketolase). The responses of rats fed irradiated chicken were similar to those fed frozen or thermally processed chicken. No evidence was found of antithiamin substances in either gamma or electron irradiation sterilized chicken.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
A	

PREFACE

The experimental portions of the study covered in this report were conducted during the period of 1 September 1978 - 1 February 1979. All raw data are being stored at Letterman Army Institute of Research. Anyone wishing to examine the raw data or to obtain copies of tables containing individual values, may do so by contacting: Commander, Letterman Army Institute of Research, ATTN: SGRD-UJZ, Presidio of San Francisco, California 94129.

In addition to the personnel listed on page vi, the authors gratefully acknowledge the assistance of: LTC John T. Huber (MOBDES) who helped summarize and interpret animal weight data; Ms. Anne Regh and Ms. Marie Rogers, typists; and Ms. Lottie Applewhite, LAIR technical editor.

TABLE OF CONTENTS

	<u>Page</u>
Abstract	i
Preface	ii
Table of Contents	iii
Report of the Quality Assurance Unit	v
List of Personnel	vi
Signatures of Principal Scientists Involved in the Study	vii
BODY OF REPORT	
INTRODUCTION	1
METHODS	2
Processing of Chicken	2
Animal Care	3
Diet Preparations	3
Blood Sampling and Analyses	4
Data Acquisition and Handling	4
Statistical Analysis of Erythrocyte Transketolase Data	5
RESULTS	5
Observations on Growth	5
Results of Erythrocyte Transketolase Analyses	6
Statistical Analysis of Erythrocyte Transketolase Data	7
DISCUSSION	7
CONCLUSIONS	8
REFERENCES	9

	<u>Page</u>
APPENDICES	
Appendix A (Figures 1 through 20)	11
Appendix B (Tables 1 through 11)	33
DISTRIBUTION LIST	45

REPORT OF THE QUALITY ASSURANCE UNIT

Summary and Conclusions: This study and draft of the final report dated 30 April 1979 have been examined and found to have been conducted as described in the protocol, its addendum, and in the standard operating procedures. The data have been audited and the reported results were found to accurately reflect the raw data.

Inspections: Written records were not made of inspections until 20 March 1979. On that date record was made to the best of memory of previous inspections. Following that date written record was made of all inspections at the time of inspection.

16 October 1978: Inspected animal rooms and feeding and weighing operations. No recommendations were made.

November 1978: Inspected erythrocyte transketolase assay. No recommendations were made.

February 1979: Mr. Paul Varing advised the Quality Assurance Unit of a downward shift in the recovery of the controls used in the erythrocyte transketolase assay from December 12 onward to the conclusion of the study. He provided the Quality Assurance Unit with documentation and speculated that the samples and controls may have been warmed during a defrosting of the freezer in which they had been stored. He advised management that data collected from December 12 onward to the conclusion of the study be multiplied by a correction factor of 1.16. The Quality Assurance Unit requested an amendment to the standard operating procedure.

26 July 1979 to 1 August 1979: Draft of the report dated 30 April 1979 was reviewed and the data audited. The amendment to the erythrocyte transketolase requested in February had not yet been received. This was requested and received. This draft accurately described the methods and standard operating procedures used and the reported results accurately reflected the raw data.

Present Recommendation to Management: An archive unit needs to be established, in which to store the data.

William R. Wise, Jr.
WILLIAM R. WISE, JR.
QUALITY ASSURANCE MONITOR
30 Aug 1979

LIST OF PERSONNEL

Study Director

LTC John T. Fruin

Principal Investigators and Supervisory Personnel

Dr. Evelyn L. McGown

Mr. Paul P. Waring

Dr. James H. Skala

CPT John L. Szurek

LTC Jerome Goldsboro

Dr. Howerde E. Sauberlich

Data Summaries and Graphic Art

Ms. Carolyn M. Lewis

Animal Care and Feed Preparation

SSG Aldino Robles

SP5 Ronald Thompson

Mr. Wanless Hatcher

Laboratory Support

CPT Michael H. Dong

SP5 Dennis M. Fisher

SP5 John M. McDonnell

SP4 Michael G. Rusnak

Mr. Richard J. O'Connor

Ms. Mary F. Lyons

Mr. John K. Knudsen

Computer Programs

Mr. William H. Langley, Jr.

Mr. Francis G. Trout, Jr.

SP5 Hal Lewis

LTC William H. Amos, Jr.

Statistician

Dr. Virginia L. Gildengorin

Quality Assurance Monitor

Mr. William R. Wise, Jr.

Civilian Consultant

Dr. Nicholas Raica, Jr.; Denver, CO

U.S. Army Medical Research and Development Command

Liaison Advisory Officer

LTC Duane E. Milmas

Chief, Office for the Wholesomeness of Irradiated Foods

Fort Detrick, MD

Signatures of Principal Scientists
Involved in the Study

We, the undersigned, believe the study described in this report to be scientifically sound and the results and interpretations to be valid. The study was conducted to comply, to the best of our ability, with the Good Laboratory Practice Regulations for Nonclinical Laboratory Studies outlined by the Food and Drug Administration.

Evelyn L. McGown 30 Aug 79
EVELYN L. MCGOWN, Ph.D.
Research Chemist

Paul P. Waring 30 Aug 79
PAUL P. WARING, B.S.
Research Chemist

Carolyn M. Lewis 30 Aug 79
CAROLYN M. LEWIS, M.S.
Nutritional Scientist

Virginia L. Gildengorin 30 Aug 79
VIRGINIA L. GILDENGORIN, Ph.D.
Mathematical Statistician

John T. Fruin 30 Aug 79
JOHN T. FRUIN, LTC, VC, DVM, Ph.D.
Study Director

Houerde E. Sauberlich 30 Aug 79
HOUERDE E. SAUBERLICH, Ph.D.
Chief, Division of Nutrition
Technology

INTRODUCTION

The testing of control and irradiated meats for antimetabolite activity against vitamins B₁ and B₆ is a requirement of the protocol entitled "Animal Feeding Protocol for Irradiation Sterilized Test Foods" originated by the Office for the Wholesomeness of Irradiated Foods, US Army Medical Research and Development Command (USAMRDC), dated 21 October 1975.

The purpose of the study reported here was to determine whether irradiation (gamma or electron) or thermal processing of chicken produces factors which are antagonistic to vitamin B₁ (thiamin) in the diet of rats.

The protocol for the antithiamin study specified that rats were to be made deficient in thiamin (according to a pre-set weight gain criterion). They were then to be repleted with various chicken-containing or semi-purified diets which had identical (high or low) thiamin contents and the recovery rates monitored. A decreased recovery rate in animals fed irradiated meat relative to those fed control meat would indicate the presence of an antithiamin substance.

Resumption of weight gain was the obvious indicator of recovery from thiamin deficiency. The other (more sensitive and specific) parameter of thiamin status was specified to be erythrocyte transketolase (ETK) activity. This enzyme is active only in the presence of a thiamin derivative, thiamin pyrophosphate. The activity of the enzyme drops rapidly in red cells of animals fed diets deficient in thiamin. Also, in vitro addition of thiamin pyrophosphate (TPP) cofactor produces a greater (%) increase in enzymatic activity in hemolysates from deficient animals than from nondeficient controls. This in vitro stimulation, "TPP effect," is considered to be indicative of the percent apoenzyme which is not saturated with cofactor.

The protocol specified that the meat diets contain 35% test meat (on a dry weight basis). Furthermore, it specified that each meat be tested at 2 levels of vitamin intake, a marginal level and a high level. The high thiamin diets were included to determine whether or not any antithiamin substances (if detected) could be overcome by additional vitamin.

All test meats originated from one lot of chicken which had been heated to inactivate enzymes. This lot was divided into four parts and each was further prepared for storage by one of the following treatments: 1) frozen (control), 2) canned (thermally processed), 3) gamma irradiated, 4) electron irradiated. The last three treatments produce shelf-stable products and are known to cause decreased vitamin content. Finally, the protocol specified the inclusion of groups fed dry, semi-purified diets without chicken.

The original protocol had specified that the two thiamin levels be 5.0 and 20.0 mg thiamin/kg diet. However preliminary studies in this laboratory suggested that there was little difference in ETK response in thiamin-deficient rats repleted at 5.0 or 20.0 mg/kg. (For further comments concerning the levels of thiamin in repletion diets, see the previous report from this laboratory on the antithiamin properties of irradiated beef (1)). In the present study, the marginal thiamin level was lowered still further to 3.0 mg/kg diet to enhance the differences in response between the high and low vitamin groups and thus to ensure that any antithiamin properties (if present) could be detected.

METHODS

Processing of Chicken. All test meats were supplied by the U.S. Army Natick Research and Development Command, Natick, MA. They were processed according to the procedure outlined in Appendix A of the protocol described in paragraph 1 of page 1.

Enzyme Inactivation: Chicken breasts, thighs and legs were deboned, skinned and ground, then mixed with sodium chloride (7.5 g/kg chicken), sodium tripolyphosphate (3.0 g/kg chicken) and crushed ice (30 g/kg chicken). The ground chicken was then placed in fibrous casings and pressed into wire cages. It was cooked in this form for two hours at a Dry Bulb Temperature of 63° to 68°C and a Wet Bulb Temperature of 46° to 52°C, after which it was cooked at temperatures up to 88°C until the internal temperature reached 73° to 77°C. After cooking, the chicken was spray washed for 20 to 30 minutes with cold tap water and then refrigerated.

All chicken, except that to be electron irradiated, was placed in cans sealed under vacuum and frozen. Chicken to be electron irradiated was placed into rectangular pouches, sealed under vacuum and frozen until irradiated. Cans of chicken to be used as the frozen control were not processed further but remained frozen until used. To prepare thermally processed chicken, cans of chicken were heated to 115.6°C for 159 minutes according to conventional commercial practice resulting in a sterilization value of not less than Fo=6.0.

Gamma Irradiation: Chicken packaged in cans was irradiated in the frozen state using the Co-60 facility of the Radiation Laboratory at the US Army Natick Laboratories, Natick, Massachusetts. The dose of radiation absorbed was between 4.7 and 7.1 million rads (Mrads).

Electron Irradiation: Chicken packed in pouches was irradiated (while frozen) by the electron linear accelerator (Linac) facility of the Radiation Laboratory at the US Army Natick Laboratories, Natick, Massachusetts. The electrons were delivered in pulses of approximately 5 microseconds and with a pulse repetition rate of approximately 180 pulses per second. The electron energy spectrum was allowed to

peak between 9 and 10 million electron volts (MeV) during irradiation of the chicken with a full width of half maximum of 0.5 MeV or less.

Animal Care

Male and female weanling rats (156 each) were purchased from Charles River Breeding Laboratories, Wilmington, Mass. Each animal was identified by ear tag and individually caged in a room with a 12-hour light/dark cycle. All were given ad libitum water and fed a semi-purified diet (Table 1) containing 20.0 mg thiamin/kg diet.

The schedule and diet codes for the studies are outlined in Table 2. After one week of quarantine and adaptation (Phase 1), 24 rats were selected at random to remain on diet A. All other animals were placed on diet B, which was identical to diet A except that thiamin had been omitted (Phase 2). Growth was monitored throughout the study by thrice weekly weighings. Animals on diet B were considered to be deficient when the average daily weight gain was less than 0.5 grams. The deficient animals were then randomly divided into 11 groups of 12 animals each. The 24 diet A animals were also divided into 2 groups of 12 each. One group of 12 diet A rats and one group of 12 diet B rats were bled by cardiac puncture and removed from the study. The remaining 10 groups of deficient animals were placed onto 10 different repletion diets (C-L) and the remaining diet A group was continued on diet A. The repletion period (Phase 3) lasted 4 weeks.

Diet Preparations

Proximate analyses of the chicken (crude fat, protein, moisture, and ash) were done by standard methods (2). Calcium and phosphorus determinations were also done (3,4). Thiamin assays were done by a microbiological method which utilizes Lactobacillus viridescens as the test organism (5). The results of these assays are summarized in Table 3.

As specified by the protocol, the meat diets were formulated to contain 35% (dry weight) chicken. The fat and protein levels of the semi-purified diets (Table 1) were adjusted to be similar to the meat diets, based on calculations from proximate analysis data. For each of the meat treatment groups (E through L) a dry premix with fat and protein omitted and containing the proper amount of thiamin was prepared in advance. When mixed with the corresponding meat (35% dry weight basis) the complete diets contained the specified levels of thiamin and were similar to the semi-purified diets, except that ground chicken replaced the casein, lard, and corn oil. Analyzed thiamin contents of repletion diets are given in Table 4. The calcium/phosphate ratio of the meat diets was calculated to be 1.294 compared to 1.25 in the semi-purified diets; therefore, no adjustment was made in these two minerals.

Each meat was ground through a 1/4 inch plate and mixed with its

corresponding dry premix in the specified ratio. The diets were prepared no longer in advance of feeding than 24 hours. Each feed jar was weighed when placed into and removed from the cage to allow estimates of food consumption. The diets remained in the cages no longer than 48 hours before being replaced by fresh diet in clean jars. Feed jars containing each diet were also placed in empty cages to serve as evaporation controls.

Blood Sampling and Analyses

Blood samples were obtained by cardiac puncture from all rats on days 7, 14, and 27 of repletion. (The last sampling was set at day 27 rather than day 28 to avoid Thanksgiving holiday.) The animals were anesthetized with penthrane gas and samples (1.5 ml each) were collected into EDTA-containing syringes. Hematocrit determinations were done in duplicate on each sample. Aliquots of each were centrifuged, the red cells were washed, hemolyzed and stored frozen until assayed. Erythrocyte transketolase (ETK) activity was determined in the presence and absence of added thiamin pyrophosphate (TPP) by a method based on that of Smeets et al. (6) but adapted for the autoanalyzer (Waring, P.P., et al., In Preparation). Enzymatic activity was expressed as ETK = I.U./ml packed red cells where I.U. = μ mole glyceraldehyde-3-phosphate (G3P) produced per minute at 37°C. The stimulation in activity in the presence of TPP was calculated as:

$$\text{TPP Effect} = \frac{[\text{ETK stimulated} - \text{ETK unstimulated}] (100)}{\text{ETK unstimulated}}$$

Quality control summaries are presented graphically in Figures 1 and 2. In both figures, the dotted lines represent 2 standard deviations on either side of the means (solid lines). Figure 1 illustrates the day-to-day chart unit response to 3 different concentrations of G3P added to a carrier hemolysate. (The carrier was a composite of blood samples prepared in advance and frozen in aliquots for use throughout the study). In addition to the carrier, two other hemolysates were prepared in advance and frozen in aliquots for quality control purposes: FC, a composite of blood samples from non-deficient rats and FD, a composite of blood samples from thiamin-deficient rats. Figure 2 illustrates the day-to-day ETK activity values (and the coefficients of variation) obtained for FC, FD and the carrier. All three controls dropped by an average of 16% on assay days 17-27 compared to the first 16. (The reason for the drop is not known, but a transient warming during freezer defrosting is suspected). Since the chart unit responses remained constant throughout the study and apparent ETK activity in all three control hemolysates dropped equivalently, their values in Figure 2 and all samples assayed on those days have been adjusted up by 16%.

Data Acquisition and Handling

Animals were weighed using an electronic balance interfaced with a programmable calculator. The weight data were permanently printed

on paper tape and recorded on a miniature tape cartridge. The information on the magnetic tape was then transferred to a minicomputer (Data General Eclipse C330) to be processed and released in report format. Programs for transferring and processing the data were developed locally in the Information Sciences Group.

Erythrocyte transketolase data were recorded in digital print-out form, on paper tape from the auto-analyzer. These data, as well as the hematocrit values, were manually transferred to the computer through a remote terminal. All programs to process enzymatic data were developed by personnel in the Division of Nutrition Technology.

Statistical Analysis of Erythrocyte Transketolase Data

The measurements taken on the three sample days (7, 14, and 27) were analyzed separately for the effects of the design variables (i.e. diet, level of thiamin and sex) along with their interactions on ETK, ETK-stimulated and TPP effect.

Based on the assumptions of normality, statistical independence and the equality of subgroup variances, three-way analyses of variances were performed with a packaged computer program, BMDP Biomedical Computer Program P2V (7). The analyses revealed that the effects of diet and level of thiamin were different for the different sex groups. Thus it was decided to separate the analyses further between the sex groups and a two-way analysis of variance was performed on each of the three dependent variables to determine the significance of the two main effects (diet and thiamin level) and their interactions.

The following model was used in the two-way analyses of variances:

$$y = m + a_i + b_j + ab_{ij} + e$$

where y = the observed ETK
 m = overall mean effect
 a_i = effect due to diet group
 b_j = effect due to thiamin level
 ab_{ij} = effect due to diet by thiamin level interaction
 e = error term

Dunnetts method of multiple comparisons (8) was used to test potentially significant differences between specific treatment groups.

RESULTS

Observations on Growth

Growth (weight) curves for the males are shown in Figures 3-8 and for the females in Figures 9-14. (Phases 1 and 2 are represented in Figures 3 and 9.) Groups A and B were indistinguishable for at least 10 days after the beginning of phase 2 and then cessation of growth in

animals on the deficient diet was remarkably abrupt. Both males and females met the criterion for deficiency (weight gain less than 0.5 g/day) 14-16 days after being placed on the thiamin deficient diet.

Figures 4-8 and Figures 10-14 show the growth curves for all groups on the various repletion diets. The growth curve of Group A has been included in each figure (dashed line). A striking observation was the fact that when the thiamin-deficient animals were placed on the repletion diets, all chicken-fed groups gained faster than those repleted on semi-purified diets. In fact, within two weeks all groups on chicken diets had caught up to the non-deficient group and the females had strikingly surpassed their Group A.

A few accidental deaths occurred as a result of the anesthesia or cardiac puncture. These losses, as well as the stresses of bleeding all animals, resulted in potentially misleading shifts in some growth curves after days 7 and 14. To eliminate the potentially misleading effects of animal deaths on group means, the growth data were recalculated and expressed as mean weight gain per group per day. Tables 5 and 6 summarize the average daily weight gain for the males and females for weekly periods as well as the overall means for the four weeks of repletion. No consistent effect of vitamin level on weight gain was found, nor were there any differences observed between chicken-fed groups. As pointed out above, the average gains for animals fed semi-purified diets were less than those for the meat-fed groups. The daily mean weight gains from which the numbers in Tables 5 and 6 were calculated are included in Tables 7 and 8.

Results of Erythrocyte Transketolase (ETK) Analyses

Erythrocyte transketolase (ETK) data (unstimulated) are summarized in Table 9 and presented graphically by day in Figures 15-20. In each figure, the far left bar represents Group A (non-deficient animals). Group B means (representing the deficient animals at day 0, before repletion) have been dotted in to aid visual comparisons. The bars are paired together according to food group and in each pair the open and diagonally hatched bars represent low and high vitamin level, respectively.

Before repletion, the deficient animals (Group B) had ETK activities 20 - 25% of the levels observed in non-deficient controls. From Figures 13 and 16, it is obvious that within 7 days, ETK activity of these animals had increased strikingly regardless of the repletion diet. Smaller increases were observed in all groups on day 14 and in the females on day 27. The high vitamin group in each pair of repletion diets consistently had the higher ETK activity. At 27 days, all experimental group means (both high and low vitamin levels) were still lower than Group A means (Figures 15 and 18).

In contrast to the enzymatic activities, the in vitro effect of TPP cofactor was not greatly influenced by thiamin status or dietary treatment (Table 10). Although the Group B males did have a higher mean

TPP effect than Group A (and also a high standard deviation), there was no difference between the corresponding female groups.

Statistical Analysis of Erythrocyte Transketolase Data

Two-way analyses of variance (diet and vitamin level as grouping factors) were done on the ETK data from Groups C - L and P values are presented in Table 11. The enzymatic parameters, ETK, ETK-stimulated and TPP effect will be discussed separately.

1. ETK and ETK-stimulated (both related to the amount of enzyme present).

Significant differences were found between the high and low vitamin groups on all three days in both male and female groups. No effect of diet was found in the males, except on day 7 ($P = 0.03$ and 0.05 for ETK and ETK-stimulated). This was probably due to the high means of Groups J and L (both high vitamin, irradiated groups) on that day. The females showed no difference between diet groups except on day 14 ($P = 0.02$ and 0.05). This effect may be attributed to the difference between Groups C and G (low vitamin: semi-purified and thermal diets). Thus there was no consistent effect of diet on ETK activity.

Since the low vitamin, irradiated meat Groups I and K were considered the crucial test groups, these two were examined more closely with respect to the corresponding frozen chicken group. In no case was an irradiated group mean found to be sufficiently lower than the frozen group to warrant further scrutiny.

2. TPP Effect (related to the % unsaturation of enzyme with cofactor).

No effect of diet was found on TPP effect. There was a significant effect of vitamin level in the males only on day 7 ($P = 0.004$). Interactions appeared in the males on day 14 ($P = 0.04$) and females on day 7 ($P = 0.05$). Thus the TPP effect appeared to be unaffected by diet or thiamin status.

DISCUSSION

In the present study, absolute ETK enzyme activity rather than % stimulation by cofactor was the more sensitive index of thiamin status. ETK in deficient animals had dropped to 20-25% of that present in non-deficient animals. The responses of animals repleted with low thiamin intake were significantly less than those of animals repleted with high vitamin intake. ETK activity in the marginal thiamin groups, thus provided valid test systems for the presence of antithiamin substances. No evidence of any thiamin antagonism was detected.

ETK activity in thiamin deficient rats increased 2-3 fold within 7 days during repletion, but still averaged approximately 60% of the

activity of non-deficient animals. These facts indicate that not all of the decreased ETK activity in deficient animals was due to an absolute loss in enzyme. Nearly half of the enzyme remained circulating as inactive, but intact, apoprotein, ready to be reactivated by the appearance (*in vivo*) of the vitamin. The slower rise in ETK activity after the first week of repletion is presumably limited by red cell turnover rate, i.e. maximal ETK activity would not be attained until new erythrocytes replaced all those circulating red cells which contained reduced ETK apoprotein.

In vitro addition of thiamin pyrophosphate cofactor reactivated the apoenzyme little, if any. The small TPP effects were at first disconcerting, but the phenomenon is not without precedence (9). Brin (10) has pointed out that TPP stimulation may not always be elicited in hemolysates of rat erythrocytes and that it varies greatly among different strains of rats (11). The small, variable reactivation by TPP *in vitro* appears to be due to a lability of apotransketolase upon hemolysis of the red cells (McGown, E.L. and M.G. Rusnak, unpublished observations). The reasons for this apparent lability are not known but are currently under investigation in this laboratory. The magnitude of the TPP effect is greatly influenced by sampling, handling and storage procedures.

Thermal processing is known to cause destruction of vitamins. Table 3 reveals that gamma irradiation is also detrimental toward the thiamin content of chicken. Electron irradiated chicken fared better, but did not retain as much of the vitamin as the frozen control. However, the destruction of thiamin (75% in thermal and gamma irradiated, 30% in electron irradiated, relative to frozen control) did not result in antivitamin substances.

CONCLUSIONS

1. Under the conditions of these studies, ETK enzymatic activity rather than TPP effect was the more sensitive indicator of thiamin status; ETK activity was highly dependent on the amount of dietary thiamin while TPP effect was largely unaffected.

2. Thiamin-deficient rats were repleted with semi-purified diets or diets containing chicken (frozen, thermally processed, gamma irradiated or electron irradiated). No difference was found among the groups in growth (weight gain) or ETK response. Specifically, neither gamma, nor electron irradiated chicken had a detrimental effect on thiamin status in rats, compared to rats fed non-irradiated (stored frozen) test meat.

REFERENCES

1. MCGOWN, E.L., C.M. LEWIS, and P.P. WARING. Investigation of Possible Anti-Thiamin Properties in Irradiation Sterilized Beef. Report No. 71. Presidio of San Francisco, California: Letterman Army Institute of Research, August 1979
2. Official Methods of Analysis of the Association of Official Analytical Chemists, 12th Edition. Association of Official Analytical Chemists, Washington, D.C., 1975, para 24.002, 24.005, 24.006, and 24.024
3. GITELMAN, H.J. An improved automated procedure for the determination of calcium in biological specimens. *Anal Biochem* 18:521-531, 1967
4. KRAML, M. A semi-automated determination of phospholipids. *Clin Chim Acta* 13:442-448, 1966
5. PEARSON, W.N. Thiamine. In: Gyorgy, P. and W.N. Pearson (eds) *The Vitamins*, 2nd Edition. New York: Academic Press, Inc., 1967
6. SHEETS, E.H.J., H. MULLER, and J. DEWAELE. A NADH-dependent transketolase assay in erythrocyte hemolysates. *Clin Chim Acta* 33:379-386, 1971
7. Biomedical Computer Programs. University of California Press, Los Angeles, CA 1975
8. DUNNETT, C.W. New tables for multiple comparisons with a control. *Biometrics* 20:482-491, 1964
9. BRUNS, F.H., F. DUNWALD, and E. NOLTMANN. Ueber den Stoffwechsel von Ribose-5-Phosphat in Haemolysaten. III. Quantitative Bestimmung von Sedoheptulose-7-Phosphat und Einige Eigenschaften der Transketolase der Erythrocyten und des Blutserums.
10. BRIN, M. Transketolase (Sedoheptulose-7-phosphate: D-Glyceraldehyde-3-phosphate Dihydroxyacetone transferase, EC 2.2.1.1) and the TPP effect in assessing thiamine adequacy. In: *Methods in Enzymology*. Volume XVIII, pp. 125-133, 1971
11. BRIN, M. Erythrocyte as a biopsy tissue for function evaluation of thiamin adequacy. *J Am Med Assoc*. 187:762-766, 1964

	<u>Page</u>
Figure 1. Day-to-Day Precision of G-3-P Internal Standards	13
Figure 2. Day-to-Day Precision of Control Hemolysates.	14
Figure 3. Growth Curves, Males, Phases 1 and 2, Groups A and B	15
Figure 4. Growth Curves, Males, Phase 3, Groups A, C and D	16
Figure 5. Growth Curves, Males, Phase 3, Groups A, E and F	17
Figure 6. Growth Curves, Males, Phase 3, Groups A, G and H	18
Figure 7. Growth Curves, Males, Phase 3, Groups A, I and J	19
Figure 8. Growth Curves, Males, Phase 3, Groups A, K and L	20
Figure 9. Growth Curves, Females, Phases 1 and 2, Groups A and B	21
Figure 10. Growth Curves, Females, Phase 3, Groups A, C and D	22
Figure 11. Growth Curves, Females, Phase 3, Groups A, E and F	23
Figure 12. Growth Curves, Females, Phase 3, Groups A, G and H	24
Figure 13. Growth Curves, Females, Phase 3, Groups A, I and J	25
Figure 14. Growth Curves, Females, Phase 3, Groups A, K and L	26
Figure 15. Transketolase Activity, Males, Day 7	27
Figure 16. Transketolase Activity, Males, Day 14.	28
Figure 17. Transketolase Activity, Males, Day 27.	29

APPENDIX A

	<u>Page</u>
Figure 18. Transketolase Activity, Females, Day 7	30
Figure 19. Transketolase Activity, Females, Day 14	31
Figure 20. Transketolase Activity, Females, Day 27	32




Figure 1. Day-to-Day Precision of G-3-P Internal Standards.




Figure 2. Day-to-Day Precision of Control Hemolyzates. Controls corrected by 16%, assays #17-27.




Figure 3. Growth Curve, Males, Phase 1 and 2, Groups A and B.




Figure 4. Growth Curve, Males, Phase 3, Groups A, C and D.




Figure 5. Growth Curve, Males, Phase 3, Groups A, E and F.




Figure 6. Growth Curve, Males, Phase 3, Groups A, C and II.




Figure 7: Growth Curve, Males, Phase 3, Groups A, I and J.




Figure 8. Growth Curve, Males, Phase 3, Groups A, K and L.




Figure 9. Growth Curve, Females, Phase 1 and 2, Groups A and B.




Figure 10. Growth Curve, Females, Phase 3, Groups A, C and B.




Figure II. Growth Curve, Females, Phase 3, Groups A, E and F.




Figure 12. Growth Curve, Females, Phase 3, Groups A, C and H.




Figure 13. Growth Curve, Females, Phase 3, Groups A, I and J.




Figure 14. Growth Curve, Females, Phase 3, Groups A, K and L.




Figure 15. Transketolase Activity, Males, Phase 3, Day 7. I indicates S.D. ▨ Non-deficient group, 20 mg thiamin/kg diet. ▩ Thiamin deficient group, Day 0. ▪ Thiamin deficient group, Day 7.




Figure 16. Transketolase Activity, Males, Phase 3, Day 14. [] Non-deficient group, 20 mg thiamin/kg diet. [] Thiamin deficient group, Day 0. [] 3 mg thiamin/kg diet.




Figure 17. Transketolase Activity, Males, Phase 3, Day 27. [Hatched] Thiamin deficient group, Day 0. [Vertical lines] Non-deficient group, 20 mg thiamin/kg diet. [Dashed] 3 mg thiamin/kg diet.




Figure 18. Transketolase Activity, Females, Phase 3, Day 7. [] indicate S.D. [] Non-deficient group, 20 mg thiamin/kg diet. [] Thiamin deficient group, Day 0. [] 3 mg thiamin/kg diet. [] 20 mg thiamin/kg diet.




Figure 19. Transketolase Activity, Females, Phase 3, Day 14, indicates S.D., Non-deficient group, 20 mg thiamin/kg diet. Thiamin deficient group, Day 0. 3 mg thiamin/kg diet. 20 mg thiamin/kg diet.




Figure 20. Transketolase Activity, Females, Phase 3, Day 27. ▨ Non-deficient group, 20 mg thiamin/kg diet. ▩ Thiamin deficient group, Day 0. ▪ Thiamin deficient group, Day 27. ▫ Non-deficient group, 3 mg thiamin/kg diet.

	<u>Page</u>
Table 1. Composition of Diets	34
Table 2. Schedule and Diet Codes for Antithiamin Studies.	35
Table 3. Proximate Analyses, Calcium, Phosphorous and Thiamin Contents in Chicken Test Meats	36
Table 4. Thiamin Contents of Repletion Diets	37
Table 5. Growth of Thiamin-Deficient Rats Repleted with Semi-Purified or Chicken-Based Diets (Males)	38
Table 6. Growth of Thiamin-Deficient Rats Repleted with Semi-Purified or Chicken-Based Diets (Females)	38
Table 7. Mean Daily Weight Gains of Male Thiamin-Deficient. Rats During Repletion with Semi-Purified or Chicken-Based Diets	40
Table 8. Mean Daily Weight Gains of Female Thiamin- Deficient Rats During Repletion with Semi-Purified or Chicken-Based Diets	41
Table 9. Summary of Erythrocyte Transketolase Activity. in Thiamin-Deficient Rats During Repletion with Semi-Purified or Chicken-Based Diets	42
Table 10. Summary of Thiamin Pyrophosphate Stimulation (TPP Effect)	43
Table 11. Analysis of Variance Significance Levels	44

TABLE 1. Composition of Diets

	<u>Semipurified</u> %	<u>Chicken</u> %
Chicken (dry weight) ¹	-	35.0
Casein, vitamin free	21.8	-
Lard	8.8	-
Corn Oil	4.4	-
L-cystine	0.2	0.2
Vitamin mix ²	2.0	2.0
Choline chloride	0.2	0.2
Mineral mix ³	4.0	4.0
Cerolose	<u>58.6</u> 100%	<u>58.6</u> 100%

¹Lor labeled "Prod 3" and "Prod 3A."

²The vitamin premix was made up in a cellulose carrier and contributed to the final diet the following vitamins in mg/kg: gelatin coated retinal (500 IU/mg) 26; Cholecalciferol (400 IU/mg), 5; DL- α -tocopheryl-acetate powder (250 IU/g), 440; Menadione - sodium bisulfite trihydrate 1.0; Riboflavin 10; Pyridoxine · HCl 20; Niacin 60; Ca - D-Pantothenate 30; Folic Acid 2.0; Biotin 1.0; B₁₂, 0.1% triturate, 30. Thiamin · HCl was incorporated into a second premix and added to the diets to achieve the specified levels.

³The mineral mix contributed to the diet the following salts: in g/kg: CaCO₃, 4.78; CaHPO₄, 22.21; NaHCO₃, 1.164; NaCl, 1.494; K₂SO₄, 6.728; MgSO₄, 2.991; MnSO₄·H₂O, 0.258; in mg/kg: ZnCO₃, 37.6; KI, 0.337; FeSO₄·7H₂O, 292; CuSO₄·5H₂O, 33.2; Na₂SeO₃, 0.33; Cr(Acetate)₃·H₂O, 4.78; MoO₃, 1.51; CoSO₄·7H₂O, 4.79.

TABLE 2. Schedule and Diet Codes for Antithiamin Studies

<u>PHASE</u>	<u>LENGTH OF PHASE</u>	<u>DIET GROUPS</u>
1. (Quarantine)	1 week	A
2. (Depletion)	14-16 days	A,B
3. (Repletion)	4 weeks	A, C-L

<u>DIET CODE</u>	<u>DIET</u>	<u>THIAMIN LEVEL</u> <u>mg/kg dry weight</u>
A	Semipurified	20.0 (non deficient control group)
B	Semipurified	0 (deficient diet)
C	Semipurified	3.0
D	Semipurified	20.0
Chicken-Containing Diets		
E	Frozen Chicken	3.0
F	Frozen Chicken	20.0
G	Thermally Processed	3.0
H	Thermally Processed	20.0
I	Gamma Irradiated	3.0
J	Gamma Irradiated	20.0
K	Electron Irradiated	3.0
L	Electron Irradiated	20.0

TABLE 3. Proximate Analyses, Calcium, Phosphorus and Thiamin Contents in Chicken Test Meats

Test Meat	Moisture %	Protein %	Fat %	Ash %	Phosphorous %	Calcium %	Thiamin	
							mg/kg Wet Wt.	mg/kg Dry Wt.
Frozen								
Mean	64.45	18.74	13.92	1.51	0.248	0.005	0.370	1.040
SD	±.41	±0.54	±.36	±.09	±.014	±.001	±.0456	±0.128
n=8*								
Thermal								
Mean	64.00	18.86	14.34	1.54	0.251	0.004	0.083	0.230
SD	±.27	±0.32	±.15	±.06	±.019	±.001	±0.002	±0.004
n=8*								
Gamma Irradiated								
Mean	63.37	19.49	14.34	1.58	0.237	0.004	0.097	0.264
SD	±.95	±0.56	±.70	±.08	±.018	±.001	±0.009	±0.025
n=8*								
Electron Irradiated								
Mean	65.19	19.03	12.85	1.54	0.241	0.004	0.243	0.698
SD	±.61	±0.51	±.60	±.07	±.021	±.001	±0.046	±0.133
n=8*								

*For thiamin analyses, n=3. The meats were sampled three times and each sample was assayed at three different dilutions.

TABLE 4. Thiamin Contents of Repletion Diets

	<u>Thiamin</u>	
	<u>mg/kg wet weight¹</u>	<u>mg/kg dry weight²</u>
	<u>Mean ± SD³</u>	<u>Mean</u>
<u>3.0 mg/kg Diets</u>		
Frozen	1.82 ± 0.19	3.0
Thermal	1.68 ± 0.16	2.8
Gamma Irradiated	1.71 ± 0.18	2.8
Electron Irradiated	1.88 ± 0.06	3.1
Semi-Purified	3.38 ± 0.08	3.4
<u>20.00 mg/kg Diets</u>		
Frozen	11.6 ± 1.9	19.0
Thermal	11.6 ± 2.0	18.8
Gamma Irradiated	12.6 ± 1.3	20.2
Electron Irradiated	14.6 ± 1.3	24.2
Semi-Purified	22.7 ± 2.3	22.7

¹Microbiological assay

²Calculated by estimating the moisture content of each diet on the basis of the original proximate analyses of the test meats (meat diets contained 35% dry weight of the respective test meat).

³Two samples were removed from each diet. Each of these was extracted and assayed at three different dilutions. The means and SDs of each set of six values are presented.

TABLE 5. Growth of Thiamin-Deficient Rats Repleted with Semi-Purified or Chicken-Based Diets (Males)

Group	Treatment	Initial Weight (g) ¹	Final Weight (g) ¹	Average Daily Gains(g)				Overall Average (g)/day
				Week 1	Week 2	Week 3	Week 4	
A	Non-deficient	190.1 ± 13.3	377.5 ± 37.6	8.9	7.3	6.8	4.9	7.0
C	Dry - 3.00	159.6 ± 18.4	350.5 ± 35.0	9.3	6.9	6.4	5.2	6.9
D	Dry - 20.00	159.9 ± 21.2	346.7 ± 33.3	7.8	7.1	6.1	4.9	6.5
E	Frozen - 3.00	160.8 ± 22.4	386.9 ± 38.0	12.4	8.6	7.0	6.1	8.4
F	Frozen - 20.00	159.5 ± 25.5	392.0 ± 51.4	11.4	8.1	7.3	7.4	8.4
G	Thermal - 3.00	158.4 ± 19.3	383.3 ± 31.5	12.1	8.6	7.7	5.3	8.4
H	Thermal - 20.00	159.3 ± 19.6	398.7 ± 25.5	13.4	9.1	8.2	5.5	9.0
I	Gamma - 3.00	160.6 ± 21.4	392.8 ± 20.3	12.4	8.3	8.2	5.7	8.6
J	Gamma - 20.00	159.5 ± 22.9	380.7 ± 37.2	11.7	8.2	7.3	5.5	8.1
K	Electron - 3.00	159.2 ± 18.7	388.5 ± 36.1	11.3	8.5	8.1	5.7	8.4
L	Electron - 20.00	160.0 ± 20.8	375.9 ± 31.9	10.5	8.3	7.3	5.5	7.9

¹Mean ± S.D.

TABLE 6. Growth of Thiamin-Deficient Rats Repleted With Semi-Purified or Chicken-Based Diets (Females)

Group	Treatment	Initial Weight (g) ¹	Final Weight (g) ¹	Average Daily Gains(g)				Overall Average (g)/Day
				Week 1	Week 2	Week 3	Week 4	
A	Non-deficient	152.5 ± 9.9	239.5 ± 19.0	3.3	4.0	3.5	3.0	3.5
C	Dry - 3.00	143.2 ± 15.1	231.5 ± 28.1	4.4	4.0	2.4	2.7	3.4
D	Dry - 20.00	142.4 ± 13.3	234.6 ± 31.3	4.3	3.8	2.8	3.1	3.5
E	Frozen - 3.00	142.4 ± 13.4	261.2 ± 23.2	6.8	4.3	3.8	3.5	4.5
F	Frozen - 20.00	142.6 ± 15.7	268.2 ± 35.1	7.2	5.3	3.2	3.9	4.9
G	Thermal - 3.00	141.9 ± 13.2	258.1 ± 31.2	6.7	4.7	3.6	3.2	4.5
H	Thermal - 20.00	143.1 ± 13.1	262.0 ± 24.7	6.7	4.7	3.7	3.4	4.6
I	Gamma - 3.00	144.3 ± 12.6	255.3 ± 25.2	6.4	5.1	3.0	3.0	4.3
J	Gamma - 20.00	142.9 ± 12.8	254.8 ± 18.6	6.1	4.3	3.8	2.8	4.2
K	Electron - 3.00	141.3 ± 13.3	261.1 ± 26.5	7.5	5.2	3.4	3.3	4.8
L	Electron - 20.00	143.5 ± 11.8	253.9 ± 21.7	6.0	4.9	2.9	3.3	4.3

¹ Mean ± S.D.

TABLE 7. Mean Daily Weight Gains of Male Thiamin-Deficient Rats During Repletion With Semi-Purified or Chicken-Based Diets (Males)

Group	Dates of Weighings										
	10/26-27	10/27-30	10/30-11/1	11/1-3	11/3-6	11/6-8	11/8-14	11/14-16	11/16-17	11/17-20	11/20-22
	Mean Weight Gain (g)/Day ¹										
A	8.7	9.0	8.8	4.6*	10.6	4.9	7.2	5.7	-0.8	6.7	5.0
C	10.3	9.8	8.0	3.9**	9.9	5.4	6.5	6.0	-0.9	7.0	5.7
D	11.6	7.0	7.2	4.6	9.8*	5.5	6.3***	5.6	0.2	6.3	5.3
E	21.8	10.6	10.4	5.5	12.3	6.1	6.8*	7.7	-0.9	9.3	4.8
F	21.5	8.6	10.5	4.0	11.8	6.5	7.5**	6.6	0.1	10.8	6.1*
G	19.0	11.4	9.7	5.1**	11.8	7.1	7.9	7.3	-0.7	7.1	5.7
H	21.3	11.4	12.1	5.9	12.0*	8.1	8.3*	8.1	-2.3	8.4	5.0
I	21.3	11.2	9.6	4.6	9.4	10.3	8.6	7.1	-7.3	9.5	6.6
J	22.0	9.4	10.0	3.7*	11.1	8.3	7.6*	6.4	-5.7	9.3	5.4
K	21.9	9.3	8.9	5.3**	10.6	8.5	8.5	6.8	-4.7	8.4	6.9
L	20.4	8.4	8.7	4.9*	11.7	6.7	7.6*	6.5	-5.7	8.2	7.0

¹Time intervals between weighings varied from 1 to 6 days. Weekly averages for Table 5 were calculated for the following periods: Week 1 (10/26-11/1), Week 2 (11/1-11/8), Week 3 (11/8-11/16), Week 4 (11/16-11/22).

Indicates loss of one (), two (**), or three (***) animals after cardiac puncture.

TABLE 8. Mean Daily Weight Gains of Female Vitamin-Deficient Rats During Repletion With Semi-Purified or Chicken-Based Diets (Females)

Group	Dates of Weighings									
	11/1-3	11/3-6	11/6-7	11/7-14	11/14-16	11/16-17	11/17-20	11/20-22	11/22-24	11/24-27
	Mean Weight Gain (g)/Day									
A	2.2	4.4	2.2	4.0	1.9	1.3	4.1	2.3	3.8	3.0
C	4.4	5.3	1.8	4.0	1.3	0.7	3.8	2.6	2.7	2.8
D	4.5	4.8	2.2	3.8*	2.4	2.8	3.1	3.6	3.3	2.7
E	9.5	6.3	3.1	4.3	2.9	0.6	5.5	4.4	2.4	3.6
F	11.4	6.5	1.2	5.3	-0.4*	5.7	4.8	4.2	4.5	3.4
G	9.9	5.8	3.3	4.7*	1.9	3.5	4.7	3.2	4.0	2.6
H	9.4	6.4	2.2	4.7	2.2	2.3	5.1	3.3	3.3	3.5
I	8.7	6.5	1.3	5.1	0.9**	1.4	5.0	4.2	3.2	2.1
J	8.9	5.6	2.2	4.3*	3.5	0.9	5.0	2.7	5.3	1.3
K	9.8	7.0	4.5	5.2**	1.5	0.1	5.7*	1.3	7.1	2.1
L	8.0	6.4	1.1	4.9**	3.0*	-1.0	4.2	3.4	4.1	2.7

Time interval between weighings varied from 1 to 7 days. Weekly averages for Table 6 were calculated for the following periods: Week 1 (11/1-11/7), Week 2 (11/7-11/14), Week 3 (11/14-11/20), Week 4 (11/20-11/27).

Indicates loss of one () or two (**) animals after cardiac puncture.

TABLE 9. Summary of Erythrocyte Transketolase Activity in Thiamin-Deficient Rats During Repletion With Semi-Purified or Chicken-Based Diets¹

<u>Day of Repletion</u>	<u>Semi-Purified</u>	<u>Frozen</u>	<u>Thermal</u>	<u>Gamma</u>	<u>Electron</u>
<u>Males - 3 mg thiamin/kg</u>					
7	0.17 ± .04	0.17 ± .02	0.16 ± .03	0.17 ± .03	0.17 ± .03
14	0.23 ± .03	0.20 ± .02	0.20 ± .03	0.20 ± .02	0.20 ± .02
27	0.21 ± .04	0.19 ± .02	0.20 ± .02	0.18 ± .02	0.20 ± .01
<u>Males - 20 mg thiamin/kg</u>					
7	0.20 ± .04	0.19 ± .03	0.19 ± .02	0.21 ± .03	0.24 ± .06
14	0.25 ± .03	0.24 ± .02	0.26 ± .04	0.27 ± .03	0.26 ± .05
27	0.25 ± .03	0.26 ± .03	0.26 ± .03	0.25 ± .03	0.26 ± .04
<u>Females - 3 mg thiamin/kg</u>					
7	0.15 ± .04	0.15 ± .02	0.16 ± .01	0.15 ± .02	0.16 ± .02
14	0.18 ± .02	0.20 ± .03	0.21 ± .02	0.20 ± .05	0.19 ± .02
27	0.22 ± .03	0.21 ± .04	0.24 ± .04	0.23 ± .04	0.23 ± .03
<u>Females - 20 mg thiamin/kg</u>					
7	0.17 ± .03	0.18 ± .02	0.19 ± .03	0.18 ± .02	0.18 ± .02
14	0.21 ± .02	0.22 ± .03	0.22 ± .03	0.21 ± .03	0.21 ± .04
27	0.24 ± .03	0.25 ± .02	0.26 ± .04	0.25 ± .04	0.25 ± .06

¹IU/ml red cells; mean ± S.D.

TABLE 10. Summary of Thiamin Pyrophosphate Stimulation (TPP Effect)*

GROUP	MALES			FEMALES			
		<u>DAY 0</u>			<u>DAY 0</u>		
A	Mean	5.47			1.56		
	S.D.	4.59			3.46		
B	Mean	16.79			1.39		
	S.D.	17.47			4.81		
		<u>Day 7</u>	<u>Day 14</u>	<u>Day 27</u>	<u>Day 7</u>	<u>Day 14</u>	<u>Day 27</u>
A	Mean	1.76	5.53	5.88	2.26	4.85	2.45
	S.D.	3.17	2.03	4.22	3.58	0.84	4.06
C	Mean	1.89	2.17	5.74	4.01	3.86	2.39
	S.D.	3.44	2.84	2.22	4.62	4.05	4.18
D	Mean	3.20	5.85	3.95	3.51	4.83	4.80
	S.D.	3.72	3.78	3.60	4.37	3.17	4.08
E	Mean	2.64	7.35	4.84	6.13	4.31	5.68
	S.D.	3.99	3.92	3.15	5.48	5.88	3.92
F	Mean	1.26	5.35	5.36	3.79	3.00	4.68
	S.D.	2.65	3.31	2.99	4.71	3.16	3.52
G	Mean	6.57	5.28	5.17	1.52	4.22	2.80
	S.D.	5.98	4.06	2.79	3.37	4.53	3.28
H	Mean	1.35	4.85	3.18	2.54	3.71	5.33
	S.D.	3.00	3.01	2.79	4.95	3.37	3.14
I	Mean	4.15	3.68	5.06	3.58	2.29	3.89
	S.D.	4.43	3.95	4.73	5.24	4.34	4.57
J	Mean	1.37	5.53	4.73	2.52	3.06	2.56
	S.D.	2.48	3.35	2.75	3.52	3.54	2.97
K	Mean	4.37	7.95	4.51	2.30	3.80	3.89
	S.D.	6.03	4.83	4.29	3.98	4.02	5.38
L	Mean	1.51	5.34	4.55	8.14	3.17	3.43
	S.D.	2.69	4.66	3.95	6.25	3.44	4.28

*All stimulation values are expressed as percent.

TABLE 11. Analysis of Variance Significance levels¹

	<u>Parameters</u> ²	<u>Food</u>	<u>Vitamin</u>	<u>Interaction</u>
<u>Males</u>				
Day 7	ETK	0.03	0.00	0.40
	ETK Stim	0.05	0.00	0.43
	TPP Effect	0.54	0.00	0.10
Day 14	ETK	0.37	0.00	0.08
	ETK Stim	0.50	0.00	0.16
	TPP Effect	0.15	0.92	0.04
Day 27	ETK	0.43	0.00	0.55
	ETK Stim	0.52	0.00	0.38
	TPP Effect	0.94	0.30	0.68
<u>Females</u>				
Day 7	ETK	0.35	0.00	0.90
	ETK Stim	0.52	0.00	0.98
	TPP Effect	0.14	0.56	0.05
Day 14	ETK	0.02	0.00	0.64
	ETK Stim	0.05	0.00	0.64
	TPP Effect	0.60	0.83	0.85
Day 27	ETK	0.32	0.01	0.94
	ETK Stim	0.32	0.00	0.89
	TPP Effect	0.49	0.63	0.31

¹P values obtained by ANOVA on five food groups, two vitamin levels.
(Groups C through L).

²ETK, Erythrocyte transketolase activity.
ETK-Stimulated, ETK activity in the presence of added thiamin pyrophosphate
co-factor.
TPP Effect, % increase in ETK due to added TPP.

OFFICIAL DISTRIBUTION LIST

Commander
US Army Medical Research and
Development Command
Fort Detrick, Frederick, MD 21701

Defense Documentation Center
ATTN: DDC-TCA (12 cys)
Cameron Station
Alexandria, VA 22314

Director of Defense Research and Engineering
ATTN: Asst Dir (Environmental and Life Sciences)
Washington, DC 20301

The Surgeon General
ATTN: DASG-TLO
Washington, DC 20314

Superintendent
Academy of Health Sciences
US Army
ATTN: AHS-COM
Fort Sam Houston, TX 78234

Assistant Dean
Inst & Rsch Support
Uniformed Services University
of Health Sciences
6917 Arlington Road
Bethesda, MD 20014

Commander
US Army Environmental Hygiene Agency
Aberdeen Proving Ground, MD 21070

US Army Research Office
ATTN: Chem & Bio Sci Div
P.O. Box 1221
Research Triangle Park, NC 27709

Biological Sciences Division
Office of Naval Research
Arlington, VA 22217

Director of Life Sciences
USAF Office of Scientific Research (AFSC)
Bolling AFB
Washington, DC 20332

Director
Walter Reed Army Institute of Research
Washington, DC 2001

Commander
US Army Medical Research
Institute of Infectious Diseases
Fort Detrick, Frederick, MD 21701

Commander
US Army Research Institute
of Environmental Medicine
Natick, MA 01760

Commander
US Army Institute of Surgical Research
Brooke Army Medical Center
Fort Sam Houston, TX 78234

Commander
US Army Institute of Dental Research
Washington, DC 20012

Commander
US Army Medical Bioengineering
Research & Development Laboratory
Fort Detrick, Frederick, MD 21701

Commander
US Army Aeromedical Research Laboratory
Fort Rucker, AL 36362

Director
US Army Biomedical Laboratory
Aberdeen Proving Ground
Edgewood Arsenal, MD 21010

Commander
Naval Medical Research Institute
National Naval Medical Center
Bethesda, MD 20014

Commander
USAF School of Aerospace Medicine
Aerospace Medical Division
Brooks AFB, TX 78235

Commander
USAF Aeromedical Research Laboratory
Wright-Patterson AFB, OH 45433

OFFICIAL COOPERATING AGENCIES

Commander
U.S. Army Natick Res&Dev Command
Natick, MA 01760

Commandant
U.S. Marine Corps
HQ, US Marine Corps, LFS-4
Washington, DC 20389

Commander
U.S. Army Natick Res&Dev Com
ATTN: Director, Food Sciences
Laboratory
Natick, MA 01760

Commanding Officer
Navy Food Service Systems Office
Washington Navy Yard
Washington, DC 20374

Chairman, DOD Food Planning Board
Director, Supply Management Policy
OASD Manpower, Reserve Affairs &
Logistics
Pentagon, Room 3B730
Washington, DC 20301

Armed Forces Radiobiology Res Inst
Building 42, NNMC
Bethesda, MD 20014

Chairman, Joint Formulation Board
DOD Food RDT&ENG Program
HQ, US Marine Corps LFS-4
Washington, DC 20380

Commander
U.S. Army Natick Res&Dev Command
ATTN: US Navy Representative/JTS
Natick, MA 01760

Commander
U.S. Army Natick Res&Dev Command
ATTN: US Army Representative/JTS
Natick, MA 01760

Commander
HQDA, Office, Deputy Chief of Staff
for Res, Dev, and Acquisition
ATTN: DAMA-CSS-D
Washington, DC 20310

Commander
U.S. Army Natick Res&Dev Command
ATTN: US Marine Corps Representative/
JTS, USMC DRDNA-ZF
Natick, MA 01760

Commander
U.S. Army Natick Res&Dev Command
ATTN: Chief, Operations Res Sys Anl
Office
Natick, MA 01760

HQDA, Office, Deputy Chief of Staff
for Logistics
ATTN: Chief, Troop Support Division
Washington, DC 20310

Chairman, DOD Food Service Facility &
Equipment Planning Board
Office of the Chief of Engineers
Forrestal Building, Room 2-F-055
Department of the Army
Washington, DC 20314

Commander
U.S. Army Troop Support Agency
ATTN: DALO-TAD
Ft. Lee, VA 23801

Defense Logistic Agency
ATTN: DLA-LF, Room 3C325
Cameron Station
Alexandria, VA 22314

Commander
U.S. Army Natick Res&Dev Command
ATTN: Technical Director
Natick, MA 01760

Commander
U.S. Army Natick Res&Dev C54mand
ATTN: US Air Force Representative/JTS
Natick, MA 01760

OFFICIAL COOPERATING AGENCIES
(Continued)

Commander	(2 cys)	Commander	(25 cys)
US Army Natick Res&Dev Command		US Army Med Res&Dev Command	
ATTN: Chief, Radiation Preserva-		ATTN: Chief, Office for the Whole-	
tion of Food Division,		someness of Irradiated Foods	
Food Engineering Laboratory		Fort Detrick, MD 21701	
Natick, MA 01760			