

AD-A075 171

STANFORD UNIV CALIF DEPT OF COMPUTER SCIENCE
KHACHIAN'S ALGORITHM FOR LINEAR PROGRAMMING.(U)
JUL 79 P 6ACS, L LOVAS
STAN-CS-79-750

F/G 12/1

UNCLASSIFIED

N00014-76-C-0330
NL

1 OF 1
AD-
A075171



END
DATE
FILMED
11-79
DDC

1.0

2.8

2.5

3.15

3.15

2.2

1.1

2.0

2.0

1.8

1.25

1.4

1.6

NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

MA075171

(Handwritten mark)

LEVEL

KHACHIAN'S ALGORITHM FOR LINEAR PROGRAMMING

by

Peter Gács and Laszlo Lovász

STAN-CS-79-750
July 1979

DDC
RECEIVED
OCT 17 1979
A

DEPARTMENT OF COMPUTER SCIENCE
School of Humanities and Sciences
STANFORD UNIVERSITY

DDC FILE COPY

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

89 10 17 047



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 STAN-CS-79-750	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9 Technical report
4. TITLE (and Subtitle) 6 Khachian's Algorithm for Linear Programming		5. TYPE OF REPORT & PERIOD COVERED technical, July 1979
7. AUTHOR(s) 10 Peter/Gács and Laszlo/Lovász		6. PERFORMING ORG. REPORT NUMBER STAN-CS-79-750
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Computer Science Stanford University Stanford, California 94305 USA		8. CONTRACT OR GRANT NUMBER(s) 15 N00014-76-C-0330 VNSF-MCS-77-23738
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Va 22217 12 15 11		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) ONR Representative - Philip Surra Durand Aeromautics Building, Room 165 Stanford University Stanford, Ca. 94305		12. REPORT DATE July 1979
16. DISTRIBUTION STATEMENT (of this Report) Releasable without limitations on dissemination.		13. NUMBER OF PAGES 12
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) Unclassified
18. SUPPLEMENTARY NOTES		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (see reverse side)		

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

✓
L. G. Khachian's algorithm to check the solvability of a system of linear inequalities with integral coefficients is described. The running time of the algorithm is polynomial in the number of digits of the coefficients. It can be applied to solve linear programs in polynomial time.
↑

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	
Unannounced Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Availand/or special
A	

Khachian's Algorithm for Linear Programming

Peter Gács and Laszlo Lovász

Computer Science Department
Stanford University
Stanford, California 94305

Abstract.

L. G. Khachian's algorithm to check the solvability of a system of linear inequalities with integral coefficients is described. The running time of the algorithm is polynomial in the number of digits of the coefficients. It can be applied to solve linear programs in polynomial time.

This research was supported in part by National Science Foundation grant MCS-77-23738 and by Office of Naval Research contract N00014-76-C-0330. Reproduction in whole or in part is permitted for any purpose of the United States government.

L. G. Khachian [Doklady Akademii Nauk SSSR, 1979, Vol. 244, No. 5, 1093-1096] published a polynomial-bounded algorithm to solve linear programming. These are some notes on this paper. We have ignored his considerations which concern the precision of real computations, in order to make the underlying idea clearer, on the other hand, proofs which are missing from his paper are given in an appendix.

Let

$$(1) \quad a_i x < b_i \quad (i = 1, \dots, m, \quad a_i \in \mathbb{Z}^n, \quad b_i \in \mathbb{Z})$$

be a system of strict linear inequalities with integral coefficients. We present an algorithm which decides whether or not (1) is solvable, and yields a solution if it is.

Define

$$L = \sum_{i,j} \log (|a_{ij}| + 1) + \sum_i \log (|b_i| + 1) + \log nm + 1$$

L is the space needed to state the problem.

The Algorithm.

We define a sequence $x_0, x_1, \dots \in \mathbb{R}^n$ and a sequence of symmetric positive definite matrices A_0, A_1, \dots recursively as follows.

$x_0 = 0$, $A_0 = 2^L I$. Assume that (x_k, A_k) is defined. Check if x_k is a solution of (1). If it is, stop. If not, pick any inequality in (1) which is violated:

$$a_i x_k \geq b_i,$$

and set

$$x_{k+1} = x_k - \frac{1}{n+1} \frac{A_k a_i}{\sqrt{a_i^T A_k a_i}},$$

$$A_{k+1} = \frac{n^2}{n^2-1} \left(A_k - \frac{2}{n+1} \frac{(A_k a_i) \cdot (A_k a_i)^T}{a_i^T A_k a_i} \right).$$

(Note that the multiplication of vector $A_k a_i$ with itself in the second term results in an $n \times n$ matrix.)

In practice, we will compute only certain approximations of x_k and A_k by decimals of a certain precision. It can be shown that approximations within $\exp(-10nL)$ preserve the validity of the following theorem.

Theorem. If the algorithm stops, x_k is a solution of (1). If the algorithm does not stop in $6n^2L$ steps, then (1) is not solvable.

The first assertion is, of course, just a repetition of the stopping rule for the algorithm. To prove the crucial second statement, we shall need a series of lemmas, along with a geometric description of what's happening.

Let $x_0 \in \mathbb{R}^n$ and A a positive definite matrix. Then

$$(x-x_0)^T A^{-1} (x-x_0) \leq 1$$

defines an ellipsoid $E = (x, A)$ with center x . Let $a \in \mathbb{R}^n$, $a \neq 0$. Then we shall denote by E^a the ellipsoid (x', A') , where

$$x'_0 = x_0 - \frac{1}{n+1} A \frac{a}{\sqrt{a^T A a}},$$

$$A' = \frac{n^2}{n^2-1} \left(A - \frac{2}{n+1} \frac{(Aa)(Aa)^T}{a^T A a} \right) .$$

We shall denote the semi-ellipsoid

$$E \cap \{x: (x-x_0)a \leq 0\}$$

by $\frac{1}{2} E_a$.

Let us remark (although this is not needed in the proof) that geometrically this construction means the following. Take a hyperplane $ax = d$, $d < ax_0$, which is tangent to E at point y . Then

$$x_0 - y = A \frac{a}{\sqrt{a^T A a}} .$$

Now E^a will be the (unique) ellipsoid which touches the hyperplane $ax = d$ at y and intersects the hyperplane $ax = ax_0$ in the same ellipsoid as E .

So here come the lemmas. The first three are facts of number-theoretic nature which probably are familiar to many people who have investigated the complexity of algorithmic problems in linear algebra.

We use the notation $|x|_\infty = \max_i x_i$, $|x|_2 = \sqrt{\sum_i x_i^2}$.

Lemma 1. Every vertex v of the polyhedron

$$a_i \leq x_i \leq b_i \quad (i = 1, \dots, m)$$

$$x_i \geq 0$$

satisfies $|v|_\infty < 2^L/n$, and its entries are rational numbers with denominator at most 2^L .

Lemma 2. If (1) has a solution, then the volume of its solutions inside the cube $|x_i| \leq 2^L$ is at least 2^{-nL} .

Lemma 3. Suppose that the system

$$a_i x < b_i + 2^{-L} \quad (i = 1, \dots, m)$$

has a solution. Then

$$a_i x \leq b_i \quad (i = 1, \dots, m)$$

has a solution.

Lemma 4. $\frac{1}{2} E_a \subset E^a$.

Lemma 5. $\lambda(E^a) = c(n)\lambda(E)$,

where

$$c(n) = \left(\frac{n^2}{n^2-1} \right)^{(n-1)/2} \frac{n}{n+1} < e^{-(1/2)(n+1)} .$$

and $\lambda(X)$ is the volume of the set X .

The proof of the theorem is quite easy now. Suppose that the procedure does not stop after $k = 6n^2L$ steps, and yet (1) is solvable. Then by Lemma 2, the set P of its solutions x inside E_0 has $\lambda(P) \geq 2^{-nL}$. By Lemma 4, $P \subset E_k$. But by Lemma 5,

$$\lambda(E_k) < e^{-(k/2)(n+1)} \lambda(E_0) < e^{-(k/2)(n+1)} 2^{2Ln} < 2^{-nL} ,$$

a contradiction.

If one would like to decide the solvability of a system of the form

$$(2) \quad a_i x \leq b_i \quad (i = 1, \dots, n)$$

then we may consider instead the system

$$(3) \quad \lceil 2^L \rceil a_i x < \lceil 2^L \rceil b_i + 1 \quad (i = 1, \dots, n) .$$

By Lemma 3, this is solvable iff (2) is solvable.

If we want to solve a linear programming problem

$$\begin{aligned} & \text{maximize} && c^T x \\ & \text{subject to} && Ax \leq b \\ & && x \geq 0 \end{aligned}$$

then consider the system of inequalities

$$c^T x = b^T y$$

$$Ax \leq b$$

$$x \geq 0$$

$$A^T y \geq c$$

$$y \geq 0 .$$

This is solvable iff the original program has a feasible solution and a finite optimum, and for any solution (x,y) of this system, x is an optimal solution of the program.

Appendix

Proof of Lemma 1. Let $v = (v_1, \dots, v_n)$. By Cramer's rule, each v_i can be expressed as

$$v_i = D_i/D,$$

where D_i and D are determinants whose entries are $0, 1, a_{ij}$ or b_i . Hence D and D_i are integers, and

$$|D| \geq 1,$$

$$|D| \leq \prod (\text{norms of row vectors})$$

$$< 2^{L/nm} < 2^{L/n},$$

and the same holds for the D_i 's. This implies the assertion.

Proof of Lemma 2. We may assume that (1) has a solution $x_0 > 0$. So the polyhedron

$$(4) \quad \begin{aligned} a_i x &\leq b_i & (i = 1, \dots, m) \\ x &\geq 0 \end{aligned}$$

has an interior point. Since it contains no line, it also has a vertex $v = (v_1, \dots, v_n)$. By Lemma 1, we know that $v_i < 2^{L/n} < \lfloor 2^L \rfloor$. It follows that the polyhedron (4) has an interior point $x = (x_1, \dots, x_n)$ with $x_j < \lfloor 2^L \rfloor$, and so the polytope

$$(5) \quad \begin{cases} a_i x \leq b_i & (i = 1, \dots, m) \\ x \geq 0 \\ x_j \leq \lfloor 2^L \rfloor & (j = 1, \dots, n) \end{cases}$$

has an interior point. Hence, it has $n+1$ vertices v_0, \dots, v_n which

are not on a hyperplane. So (5) has volume at least

$$\frac{1}{n!} \det \begin{pmatrix} 1 & 1 & \dots & 1 \\ v_0 & v_1 & \dots & v_n \end{pmatrix} .$$

Here, by Lemma 1, we get that

$$v_i = \frac{1}{D_i} u_i ,$$

where u_i is an integer vector and D_i is an integer $< 2^{L/n}$. So

$$\begin{aligned} \det \begin{pmatrix} 1 & \dots & 1 \\ v_1 & \dots & v_n \end{pmatrix} &= \frac{1}{|D_1| \dots |D_n|} \det \begin{pmatrix} D_1 & \dots & D_n \\ u_1 & \dots & u_n \end{pmatrix} \\ &\geq \frac{1}{|D_1| \dots |D_n|} \geq 2^{-nL \cdot n^n} \end{aligned}$$

since the determinant in the second expression is a non-zero integer.

So the volume of the polytope (5) is at least $\frac{1}{n!} 2^{-nL \cdot n^n} > 2^{-nL}$.

Proof of Lemma 3. For $x \in \mathbb{R}^n$, set

$$\theta_i(x) = a_i x - b_i .$$

Let $x_0 \in \mathbb{R}^n$ be arbitrary.

Claim 1. There exists an $x_1 \in \mathbb{R}^n$ such that

$$(1) \quad \theta_i(x_1) \leq \max(0, \theta_i(x_0)) \quad (i = 1, \dots, m)$$

and

$$(2) \quad \text{The vectors } \{a_i : \theta_i(x_1) \geq 0\} \text{ span every other vector } a_i .$$

To prove the claim, it suffices to show that if x_0 does not satisfy (2) then we can find a vector x_1 such that x_1 satisfies (1) and $\theta_i(x_1) \geq 0$ holds, for more indices i than $\theta_i(x_0) \geq 0$. Repeating

this at most m times we must obtain an x_1 satisfying both (1) and (2).

Let, say $\theta_1(x_0), \dots, \theta_k(x_0) \geq 0$, $\theta_{k+1}(x_0), \dots, \theta_m(x_0) < 0$. Suppose that a_ν ($\nu > k$) is not a linear combination of a_1, \dots, a_k . Then the system of linear equations

$$a_i y = 0 \quad (i = 1, \dots, k)$$

$$a_\nu y = 1$$

is solvable. Let y_0 be a solution and consider

$$x_1 = x_0 + t y_0,$$

where

$$t = \max\{s \in \mathbb{R} : s a_j y_0 + \theta_j \leq 0 \ (j = k+1, \dots, m)\}$$

t is finite, in fact $t \leq -\theta_\nu$.

Then by the choice of t ,

$$\theta_i(x_1) = t a_i y_0 + \theta_i(x_0) \begin{cases} = \theta_i(x_0) & \text{if } 1 \leq i \leq k, \\ \leq 0 & \text{if } k+1 \leq i \leq m, \end{cases}$$

and equality holds for at least one $1 \leq i \leq m$. This proves the Claim.

Assume now that x_0 is such that

$$a_i x_0 < b_i + 2^{-L} \quad (i = 1, \dots, m)$$

Let, say $a_i x_0 \geq b_i$ for $i = 1, \dots, k$. Choose the labelling so that a_1, \dots, a_r are linearly independent but a_{r+1}, \dots, a_k are spanned by them. By the Claim, we may assume that a_{k+1}, \dots, a_n are also spanned by a_1, \dots, a_r .

Now let z be a solution of the system of linear equations

$$a_i z = b_i \quad (i = 1, \dots, r) .$$

We show that z satisfies

$$a_i z \leq b_i$$

for every $1 \leq i \leq m$. We know that

$$a_i = \sum_{j=1}^r \lambda_j a_j$$

with some real numbers λ_j . In fact by Cramer's rule we also know that

$$\lambda_j = D_j/D ,$$

where D_j and D are determinants formed by some entries of the vectors a_i and hence they are integers with absolute value less than $2^L/n$. Now

$$\begin{aligned} D(a_i z - b_i) &= \sum_{j=1}^r D_j a_j z - D b_i \\ &= \sum_{j=1}^r D_j b_j - D b_i . \end{aligned}$$

To estimate the right hand side, use that

$$\begin{aligned} \sum_{j=1}^r D_j b_j - D b_i &= \sum_{j=1}^r D_j (a_j x_0 - \theta_j(x_0)) - D(a_i x_0 - \theta(x_0)) \\ &= D \cdot \theta_i(x_0) - \sum_{j=1}^r D_j \theta_j(x_0) \\ &\leq D \cdot 2^{-L} + \sum_{j=1}^r |D_j| 2^{-L} < 1 , \end{aligned}$$

and since the left hand side is an integer,

$$\sum_{j=1}^r D_j b_j - D b_i \leq 0 ,$$

which proves the assertion.

Proof of Lemma 4. We may assume that $x_0 = 0$, $A = I$ (i.e., the ellipsoid is the unit sphere about 0) and that $a = (-1, 0, \dots, 0)^T$, since the contents of the lemma is invariant under affine transformations of the space.

Then

$$x'_0 = \left(\frac{1}{n+1}, 0, \dots, 0 \right)^T$$

and

$$A' = \text{diag} \left(\frac{n^2}{(n+1)^2}, \frac{n^2}{n^2-1}, \dots, \frac{n^2}{n^2-1} \right) .$$

Suppose $x \in \frac{1}{2} E_a$. Then $|x|_2 \leq 1$, $1 \geq \xi_1 = -a^T x \geq 0$. We have to show that

$$(x-x'_0)^T A'^{-1} (x-x'_0) \leq 1 .$$

But

$$\begin{aligned} (x-x'_0)^T A'^{-1} (x-x'_0) &= x^T A'^{-1} x - 2x^T A'^{-1} x'_0 + x'^T_0 A'^{-1} x'_0 \\ &= \frac{n^2-1}{n^2} x^2 + \frac{2n+2}{n^2} \xi_1^2 - 2 \frac{n+1}{n^2} \xi_1 + \frac{1}{n^2} \\ &= \frac{n^2-1}{n^2} (x^2-1) + \frac{2n+2}{n^2} \xi_1 (\xi_1-1) + 1 \leq 1 . \end{aligned}$$

Proof of Lemma 5. We may assume again that E is the unit sphere about 0 and $a = (1, 0, \dots, 0)^T$, since affine transformations do not change the proportion of volumes. By a well-known formula,

$$\begin{aligned} \lambda(E^a) &= \frac{\sqrt{\det A'}}{\sqrt{\det A}} \cdot \lambda(E) = \sqrt{\det A'} \lambda(E) \\ &= \frac{n}{n+1} \left(\frac{n^2}{n^2-1} \right)^{(n-1)/2} \cdot \lambda(E) = c(n) \cdot \lambda(E) . \end{aligned}$$

To estimate this factor use that

$$\frac{n^2}{n^2-1} = 1 + \frac{1}{n^2-1} < e^{1/(n^2-1)}$$

and

$$\frac{n}{n+1} = 1 - \frac{1}{n+1} < e^{-1/(n+1)} .$$

Substituting these bounds we get

$$c(n) < e^{-1/(2(n+1))} .$$