

AD-A077 106

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

A PRIORI ESTIMATES AND EXISTENCE RESULTS FOR SOME SEMILINEAR EL--ETC(U)

JUL 79 P L LIONS

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-1975

NL

| OF |
AD-
A077106



END
DATE
FILMED

12-79
DDC

~~LEVEL~~
~~LEVEL~~

(10) 8

AD A 077106

MRC Technical Summary Report #1975

A PRIORI ESTIMATES AND EXISTENCE
RESULTS FOR SOME SEMILINEAR ELLIPTIC
PROBLEMS

P. L. Lions

DDC
RECEIVED
NOV 26 1979
REGISTRY

**Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706**

July 1979

(Received March 26, 1979)

DDC FILE COPY

**Approved for public release
Distribution unlimited**

Sponsored by
U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

79 11 19 037

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

6 A PRIORI ESTIMATES AND EXISTENCE RESULTS FOR SOME
SEMILINEAR ELLIPTIC PROBLEMS.

10 P. L./Lions

9 Technical Summary Report #1975
July 1979

11
ABSTRACT

We study the existence of non-trivial, positive solutions of

$$(1) \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

We prove the existence of a priori estimates for positive solutions of (1), provided f does not increase too rapidly at infinity and provided Ω has some geometrical properties. From these a priori estimates we deduce the existence of non-trivial, positive solutions of (1).

14 MRC-TSR-1975

AMS (MOS) Subject Classifications: 35B45, 35B99

Key Words: Semilinear problems, Symmetry, Non-trivial solutions

Work Unit Number 1 (Applied Analysis)

- A -

* E.N.S. 45 rue d'Ulm, Paris 75005, FRANCE, and Lab. d'Anal. Num., 5 Place Jussieu, Paris 75230 Cedex 05, FRANCE.

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024

15

221 200

mt

SIGNIFICANCE AND EXPLANATION

Problems arising in, for example, physics, biology and chemical reactors lead to boundary value problems for semilinear elliptic equations. The question of identifying for which nonlinearities these semilinear problems have solutions is by no means settled. Roughly speaking, nonlinearities which grow too rapidly yield unsolvable problems while under other conditions on the growth existence can be proved. Here we improve some conditions which were previously known to guarantee the existence of positive solutions of such problems.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

-B-

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

I. Introduction:

In this paper we consider the following problem: find a solution of u

$$(1) \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where f is a given nonlinear function such that $f(0) = 0$, and Ω is a bounded domain in \mathbb{R}^N with a smooth boundary. Obviously zero is a solution of (1), so we look for a non-trivial solution of (1), and actually we shall restrict ourselves to positive solutions.

Under various conditions on f , the existence of a positive, non-trivial solution to (1) has been proved, for instance, by P. H. Rabinowitz [9], [10], A. Ambrosetti and P. H. Rabinowitz [8], R. E. L. Turner [11], H. Brézis and R. E. L. Turner [4], R. Nussbaum [7], H. Amann [1], S. I. Pohozaev [8].

We shall be concerned here with the superlinear case i.e., we shall assume that the function f satisfies the condition

$$(2) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t} > \lambda_1,$$

where λ_1 is the first eigenvalue of $-\Delta$, with Dirichlet boundary conditions. One way to solve (1), in this case, is to obtain L^∞ a priori estimates for positive solutions. This was done in [11], [4], assuming that

$$\lim_{t \rightarrow \infty} f(t)t^{-\frac{N+1}{N-1}} = 0.$$

We prove here that such estimates can be obtained if

$$(3) \quad \lim_{t \rightarrow \infty} f(t)t^{-\beta} = 0 \text{ for some } \beta < \frac{N}{N-2} \text{ (if } N = 2 \text{ for any } \beta)$$

* E.N.S. 45 rue d'Ulm, Paris 75005, FRANCE, and Lab. d'Anal. Num., 5 Place Jussieu, Paris 75230 Cedex 05, FRANCE.

provided Ω (when $N \geq 3$) has some geometrical properties. For example, we proved that there exists a constant C such that if u satisfies

$$-\Delta u = u^{\beta} \text{ in } \Omega, \quad u \geq 0 \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega$$

with $\beta < \frac{N}{N-2}$ and Ω convex, then $\|u\|_{L^{\infty}(\Omega)} \leq C$.

The main new ingredient used to obtain such estimates is the convenient application of some recent remarkable results of B. Gidas, Wei-Ming Ni and L. Nirenberg [5]. They prove that if u is a positive solution of (1), then the set of critical points of u is contained in a proper subset of Ω (independent of u and f), if Ω (for $N \geq 3$) has some geometrical properties. Using this result, an easy method is then used to give the needed a priori estimates and the existence of positive solutions of (1).

The author wishes to thank Profs. M. G. Crandall, B. Gidas, L. Nirenberg and R. E. L. Turner for various discussions on related topics.

II. A priori estimates:

Let Ω be a bounded, connected domain with a smooth boundary:

Theorem 1: We assume that f is locally Lipschitz from \mathbb{R}_+ to \mathbb{R}_+ , satisfies (2) and (3). We also assume that if $N \geq 3$, Ω is convex. Then there exists a constant C such that for all $u \in C^2(\bar{\Omega})$ satisfying (1')

$$(1') \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u \geq 0 & \text{in } \Omega, \quad u = 0 \text{ on } \partial\Omega; \end{cases}$$

we have

$$\|u\|_{L^{\infty}(\Omega)} \leq C.$$

Remark 1: Instead of assuming Ω convex, we could have assumed

$$(*) \quad \begin{cases} \exists \Omega' \text{ proper closed subset of } \Omega \text{ such that for all positive } u \text{ satisfying} \\ (1') \text{ the set } \{x \in \Omega \mid \nabla u(x) = 0\} \subset \Omega'. \end{cases}$$

In view of the results of B. Gidas, Wei-Ming Ni and L. Nirenberg [5], (*) is satisfied for every domain Ω in \mathbb{R}^2 , and for convex domains in \mathbb{R}^N ($N \geq 3$). Other cases

are given in [5]. The question of whether (*) is satisfied for any domain in \mathbb{R}^N ($N \geq 3$) is open and a positive answer would imply that Theorem 1 is true without the restriction Ω convex.

Proof of Theorem 1: The proof is divided into two steps: 1) we prove that u and f are bounded in $L^1_{loc}(\Omega)$. 2) we deduce the $L^\infty(\Omega)$ estimate.

1) Let σ_1 be a fixed eigenfunction of $-\Delta$ corresponding to the first eigenvalue i.e.:

$$-\Delta \sigma_1 = \lambda_1 \sigma_1 \text{ in } \Omega, \quad \sigma_1 = 0 \text{ on } \partial\Omega, \quad \sigma_1 \in C^2(\bar{\Omega}).$$

We may choose σ_1 so that $\sigma_1 > 0$ in Ω . Now (as in [11], [4], [7], [3]) we multiply (1) by σ_1 and we obtain

$$\lambda_1 \int_{\Omega} u \sigma_1 dx = \int_{\Omega} f(u) \sigma_1 dx.$$

But using (2) there exists $\lambda > \lambda_1$, $C > 0$ such that $f(t) \geq \lambda t - C$ for every $t \geq 0$. Thus we deduce $\int_{\Omega} u \sigma_1 \leq \text{const.}$, and $\int_{\Omega} f(u) \sigma_1 dx \leq \text{const.}$ Since $\sigma_1 > 0$ in Ω , we have an estimation of u and $f(u)$ in $L^1_{loc}(\Omega)$.

2) Thus, $-\Delta u$ and u are in a bounded subset of $L^1_{loc}(\Omega)$. But this implies by standard regularity results (localize and use $L^1 \subset W^{-\epsilon, p}$ $\forall p < \frac{N}{N-\epsilon}$) that u is bounded in $L^p_{loc}(\Omega)$ ($\forall p < \frac{N}{N-2}$ if $N \geq 3$, $\forall p < \infty$ if $N = 2$). Now we use (3) and we have

$$f(u) \text{ is bounded in } L^{q_1}_{loc}(\Omega) \text{ with } q_1 > 1.$$

This implies u is bounded in $W^{2, q_1}_{loc}(\Omega)$ and by a standard bootstrap argument we obtain after a finite number of iterations of the preceding argument

$$u \text{ is bounded in } W^{2, q}_{loc}(\Omega) \text{ with } q > \frac{N}{2}.$$

in particular u is bounded in $L^\infty_{loc}(\Omega)$. To conclude, we apply the previously mentioned result of [5]: indeed, there exists some open set Ω' (independent of f and u) such that if $u \neq 0$

$$\{x \in \Omega \mid \forall u(x) = 0\} \subset \Omega' \subset \bar{\Omega}' \subset \Omega.$$

Thus, $\|u\|_{L^\infty(\Omega)} = \|u\|_{L^\infty(\Omega')}$, and $\|u\|_{L^\infty(\Omega')} \leq \text{const.}$

Remark 2: 1) If $N = 2$, with a slight refinement of the second part of the proof, we can prove Theorem 1 with the assumption

$$(3') \quad \lim_{t \rightarrow \infty} f(t) \exp\{-t/C\} = 0,$$

where C is some constant depending only on Ω , and on $\lim_{t \rightarrow \infty} \frac{f(t)}{t}$. Indeed, if u is such that $-\Delta u$ is in bounded subset of $L'_{\text{loc}}(\Omega)$ (and $\Omega \subset \mathbb{R}^2$) then for some constant C'

$$\exp \frac{u}{C'} \in L^1(\Omega') \quad (\Omega' \text{ chosen as above}).$$

Now if $C > C'$, this implies that $f(u) \in L'_{\text{loc}}(\Omega')$. Thus $u \in W^{2,C/C'}_{\text{loc}}(\Omega')$ and this implies the L^∞_{loc} bound.

III. The existence result:

Theorem 2: Under the assumptions of Theorem 1 and if, in addition, we assume

$$(4) \quad \overline{\lim}_{t \rightarrow 0_+} \frac{f(t)}{t} < \lambda_1,$$

then there exists a positive solution of (1); that is, $u(x) > 0$ in Ω and u satisfies

$$(1) \quad -\Delta u = f(u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad u \in C^2(\bar{\Omega}).$$

Remark 3: Actually, Theorem 1 implies (as in [4]) the existence of a branch of positive solutions (λ, u) of the eigenvalue problem

$$-\Delta u = \lambda f(u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad u \in C^2(\bar{\Omega}),$$

provided f satisfies, instead of (4),

$$(4') \quad \lim_{t \rightarrow 0_+} \frac{f(t)}{t} = 0.$$

Remark 4: Theorem 1 can also be used in order to obtain existence results in the case where $f(0) > 0$ (see [6]).

Proof of Theorem 2: We use the same argument as in [4]. To this end we have only to establish the statement (5) below. We can then apply the proof of [4] (which uses

a simple fixed point argument):

$$(5) \quad \begin{cases} \text{If } (t,u) \text{ satisfy } -\Delta u = f(u) + t\sigma_1 \text{ in } \Omega, u \in C^2(\bar{\Omega}), u \geq 0 \text{ in } \Omega, \\ u = 0 \text{ on } \partial\Omega, t \geq 0 \text{ then } t \leq t_0, \|u\|_{L^\infty(\Omega)} \leq C_0. \end{cases}$$

where σ_1 is chosen as in the proof of Theorem 1.

To prove (5), we consider u a solution of

$$\begin{cases} -\Delta u = f(u) + t\sigma_1 \text{ in } \Omega, \\ u \in C^2(\bar{\Omega}), u \geq 0 \text{ in } \Omega, u = 0 \text{ on } \partial\Omega, \end{cases}$$

with $t \geq 0$. The first step of the proof of Theorem 1 then shows that we have

$$t \leq t_0.$$

Now, the second step of the proof applies, since the results of [5] still hold for u which is a solution of

$$\begin{cases} -\Delta u = f(x,u) \text{ in } \Omega, \\ u \in C^2(\bar{\Omega}), u \geq 0 \text{ in } \Omega, u = 0 \text{ on } \partial\Omega, \end{cases}$$

with $f(x,u) = f(u) + t\sigma_1(x)$. Indeed we have

$$\frac{\partial \sigma_1}{\partial \nu} \leq -a < 0 \text{ on } \partial\Omega,$$

where ν is the unit exterior normal vector, and this enables us to apply the results and the methods of [5].

REFERENCES

- [1] H. Amann: On the existence of positive solutions of non-linear elliptic boundary value problems, *Ind. Univ. Math. J.*, (1971), 125-146.
- [2] A. Ambrosetti and P. H. Rabinowitz: Dual variational methods in critical point theory and applications, *J. Functional Anal.* 14 (1973), 349-381.
- [3] H. Berestycki and P. L. Lions: A local approach to the existence of solutions of semilinear problems in \mathbb{R}^N . To appear.
- [4] H. Brézis and R. E. L. Turner: On a class of superlinear elliptic problems, *Comm. in PDE* 2 (6) (1977), 601-614.
- [5] B. Gidas, Wei-Ming Ni and L. Nirenberg: Symmetry and related properties via the maximum principle. To appear in *Comm. Math. Phys.*
- [6] P. L. Lions, to appear.
- [7] R. Nussbaum: Positive solutions of some nonlinear elliptic boundary value problems, *J. Math. Anal. Appl.* 51 (1975), 461-482.
- [8] S. I. Pohozaev: Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$, *Soviet Math.* 6 (1965), 1408-1411.
- [9] P. H. Rabinowitz: Some critical point theorems and applications to semilinear elliptic partial differential equations. To appear in *Ann. Scuola Norm. Pisa.*
- [10] P. H. Rabinowitz: Variational methods for nonlinear eigenvalue problems, in *Eigenvalues of Nonlinear Problems*, Edizioni Gemonese, Rome (1974), 141-195.
- [11] R. E. L. Turner: A priori bounds for positive solutions of nonlinear elliptic equations in two variables, *Duke Math. J.* 41, No. 4 (1974), 759-774.

PLL/scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1975	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A PRIORI ESTIMATES AND EXISTENCE RESULTS FOR SOME SEMILINEAR ELLIPTIC PROBLEMS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) P. L. Lions		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706 Wisconsin		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE July 1979
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Semilinear problems Symmetry Non-trivial solutions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We study the existence of non-trivial, positive solutions of (1) $\begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$ We prove the existence of a priori estimates for positive solutions of (1), provided f does not increase too rapidly at infinity and provided Ω has some geometrical properties. From these a priori estimates we deduce the existence of non-trivial, positive solutions of (1).		

AD-A077 106

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

A PRIORI ESTIMATES AND EXISTENCE RESULTS FOR SOME SEMILINEAR EL--ETC(U)

JUL 79 P L LIONS

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-1975

NL

| OF |
AD-
A077106



END
DATE
FILMED

12-79
DDC

~~LEVEL~~

(10) 8

MRC Technical Summary Report #1975

A PRIORI ESTIMATES AND EXISTENCE
RESULTS FOR SOME SEMILINEAR ELLIPTIC
PROBLEMS

P. L. Lions

AD A 077106

DDC
RECEIVED
NOV 26 1979
REGISTRY

**Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706**

July 1979

(Received March 26, 1979)

DDC FILE COPY

**Approved for public release
Distribution unlimited**

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

79 11 19 037

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

6 A PRIORI ESTIMATES AND EXISTENCE RESULTS FOR SOME
SEMILINEAR ELLIPTIC PROBLEMS.

10 P. L./Lions

9 Technical Summary Report #1975
July 1979

11 ABSTRACT

12 9

We study the existence of non-trivial, positive solutions of

$$(1) \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

We prove the existence of a priori estimates for positive solutions of (1), provided f does not increase too rapidly at infinity and provided Ω has some geometrical properties. From these a priori estimates we deduce the existence of non-trivial, positive solutions of (1).

14 MRC-TSR-1975

AMS (MOS) Subject Classifications: 35B45, 35B99

Key Words: Semilinear problems, Symmetry, Non-trivial solutions

Work Unit Number 1 (Applied Analysis)

- A -

* E.N.S. 45 rue d'Ulm, Paris 75005, FRANCE, and Lab. d'Anal. Num., 5 Place Jussieu, Paris 75230 Cedex 05, FRANCE.

Sponsored by the United States Army under Contract No. DAAG29-75-C-0024

15

221 200

mt

SIGNIFICANCE AND EXPLANATION

Problems arising in, for example, physics, biology and chemical reactors lead to boundary value problems for semilinear elliptic equations. The question of identifying for which nonlinearities these semilinear problems have solutions is by no means settled. Roughly speaking, nonlinearities which grow too rapidly yield unsolvable problems while under other conditions on the growth existence can be proved. Here we improve some conditions which were previously known to guarantee the existence of positive solutions of such problems.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
DDC TAB	
Unannounced	
Justification _____	
By _____	
Distribution/ _____	
Availability Codes	
Dist	Avail and/or special
A	

-B-

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

I. Introduction:

In this paper we consider the following problem: find a solution of u

$$(1) \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where f is a given nonlinear function such that $f(0) = 0$, and Ω is a bounded domain in \mathbb{R}^N with a smooth boundary. Obviously zero is a solution of (1), so we look for a non-trivial solution of (1), and actually we shall restrict ourselves to positive solutions.

Under various conditions on f , the existence of a positive, non-trivial solution to (1) has been proved, for instance, by P. H. Rabinowitz [9], [10], A. Ambrosetti and P. H. Rabinowitz [8], R. E. L. Turner [11], H. Brézis and R. E. L. Turner [4], R. Nussbaum [7], H. Amann [1], S. I. Pohozaev [8].

We shall be concerned here with the superlinear case i.e., we shall assume that the function f satisfies the condition

$$(2) \quad \lim_{t \rightarrow \infty} \frac{f(t)}{t} > \lambda_1,$$

where λ_1 is the first eigenvalue of $-\Delta$, with Dirichlet boundary conditions. One way to solve (1), in this case, is to obtain L^∞ a priori estimates for positive solutions. This was done in [11], [4], assuming that

$$\lim_{t \rightarrow \infty} f(t)t^{-\frac{N+1}{N-1}} = 0.$$

We prove here that such estimates can be obtained if

$$(3) \quad \lim_{t \rightarrow \infty} f(t)t^{-\beta} = 0 \text{ for some } \beta < \frac{N}{N-2} \text{ (if } N = 2 \text{ for any } \beta)$$

* E.N.S. 45 rue d'Ulm, Paris 75005, FRANCE, and Lab. d'Anal. Num., 5 Place Jussieu, Paris 75230 Cedex 05, FRANCE.

provided Ω (when $N \geq 3$) has some geometrical properties. For example, we proved that there exists a constant C such that if u satisfies

$$-\Delta u = u^\beta \text{ in } \Omega, \quad u \geq 0 \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega$$

with $\beta < \frac{N}{N-2}$ and Ω convex, then $\|u\|_{L^\infty(\Omega)} \leq C$.

The main new ingredient used to obtain such estimates is the convenient application of some recent remarkable results of B. Gidas, Wei-Ming Ni and L. Nirenberg [5]. They prove that if u is a positive solution of (1), then the set of critical points of u is contained in a proper subset of Ω (independent of u and f), if Ω (for $N \geq 3$) has some geometrical properties. Using this result, an easy method is then used to give the needed a priori estimates and the existence of positive solutions of (1).

The author wishes to thank Profs. M. G. Crandall, B. Gidas, L. Nirenberg and R. E. L. Turner for various discussions on related topics.

II. A priori estimates:

Let Ω be a bounded, connected domain with a smooth boundary:

Theorem 1: We assume that f is locally Lipschitz from \mathbb{R}_+ to \mathbb{R}_+ , satisfies (2) and (3). We also assume that if $N \geq 3$, Ω is convex. Then there exists a constant C such that for all $u \in C^2(\bar{\Omega})$ satisfying (1')

$$(1') \quad \begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u \geq 0 & \text{in } \Omega, \quad u = 0 \text{ on } \partial\Omega; \end{cases}$$

we have

$$\|u\|_{L^\infty(\Omega)} \leq C.$$

Remark 1: Instead of assuming Ω convex, we could have assumed

$$(*) \quad \begin{cases} \exists \Omega' \text{ proper closed subset of } \Omega \text{ such that for all positive } u \text{ satisfying} \\ (1') \text{ the set } \{x \in \Omega \mid \nabla u(x) = 0\} \subset \Omega'. \end{cases}$$

In view of the results of B. Gidas, Wei-Ming Ni and L. Nirenberg [5], (*) is satisfied for every domain Ω in \mathbb{R}^2 , and for convex domains in \mathbb{R}^N ($N \geq 3$). Other cases

are given in [5]. The question of whether (*) is satisfied for any domain in \mathbb{R}^N ($N \geq 3$) is open and a positive answer would imply that Theorem 1 is true without the restriction Ω convex.

Proof of Theorem 1: The proof is divided into two steps: 1) we prove that u and f are bounded in $L^1_{loc}(\Omega)$, 2) we deduce the $L^\infty(\Omega)$ estimate.

1) Let σ_1 be a fixed eigenfunction of $-\Delta$ corresponding to the first eigenvalue i.e.:

$$-\Delta \sigma_1 = \lambda_1 \sigma_1 \text{ in } \Omega, \quad \sigma_1 = 0 \text{ on } \partial\Omega, \quad \sigma_1 \in C^2(\bar{\Omega}).$$

We may choose σ_1 so that $\sigma_1 > 0$ in Ω . Now (as in [11], [4], [7], [3]) we multiply (1) by σ_1 and we obtain

$$\lambda_1 \int_{\Omega} u \sigma_1 dx = \int_{\Omega} f(u) \sigma_1 dx.$$

But using (2) there exists $\lambda > \lambda_1$, $C > 0$ such that $f(t) \geq \lambda t - C$ for every $t \geq 0$.

Thus we deduce $\int_{\Omega} u \sigma_1 \leq \text{const.}$, and $\int_{\Omega} f(u) \sigma_1 dx \leq \text{const.}$ Since $\sigma_1 > 0$ in Ω , we have an estimation of u and $f(u)$ in $L^1_{loc}(\Omega)$.

2) Thus, $-\Delta u$ and u are in a bounded subset of $L^1_{loc}(\Omega)$. But this implies by standard regularity results (localize and use $L^1 \subset W^{c,p}$ $\forall p < \frac{N}{N-c}$) that u is bounded in $L^p_{loc}(\Omega)$ ($\forall p < \frac{N}{N-2}$ if $N \geq 3$, $\forall p < \infty$ if $N = 2$). Now we use (3) and we have

$$f(u) \text{ is bounded in } L^{q_1}_{loc}(\Omega) \text{ with } q_1 > 1.$$

This implies u is bounded in $W^{2,q_1}_{loc}(\Omega)$ and by a standard bootstrap argument we obtain after a finite number of iterations of the preceding argument

$$u \text{ is bounded in } W^{2,q}_{loc}(\Omega) \text{ with } q > \frac{N}{2},$$

in particular u is bounded in $L^\infty_{loc}(\Omega)$. To conclude, we apply the previously mentioned result of [5]: indeed, there exists some open set Ω' (independent of f and u) such that if $u \neq 0$

$$\{x \in \Omega \mid \forall u(x) = 0\} \subset \Omega' \subset \bar{\Omega}' \subset \Omega.$$

Thus, $\|u\|_{L^\infty(\Omega)} = \|u\|_{L^\infty(\Omega')}$, and $\|u\|_{L^\infty(\Omega')} \leq \text{const.}$

Remark 2: 1) If $N = 2$, with a slight refinement of the second part of the proof, we can prove Theorem 1 with the assumption

$$(3') \quad \lim_{t \rightarrow \infty} f(t) \exp\{-t/C\} = 0,$$

where C is some constant depending only on Ω , and on $\lim_{t \rightarrow \infty} \frac{f(t)}{t}$. Indeed, if u is such that $-\Delta u$ is in bounded subset of $L^1_{\text{loc}}(\Omega)$ (and $\Omega \subset \mathbb{R}^2$) then for some constant C'

$$\exp \frac{u}{C'} \in L^1(\Omega') \quad (\Omega' \text{ chosen as above}).$$

Now if $C > C'$, this implies that $f(u) \in L^{C/C'}_{\text{loc}}(\Omega')$. Thus $u \in W^{2,C/C'}_{\text{loc}}(\Omega')$ and this implies the L^{∞}_{loc} bound.

III. The existence result:

Theorem 2: Under the assumptions of Theorem 1 and if, in addition, we assume

$$(4) \quad \overline{\lim}_{t \rightarrow 0^+} \frac{f(t)}{t} < \lambda_1,$$

then there exists a positive solution of (1); that is, $u(x) > 0$ in Ω and u satisfies

$$(1) \quad -\Delta u = f(u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad u \in C^2(\bar{\Omega}).$$

Remark 3: Actually, Theorem 1 implies (as in [4]) the existence of a branch of positive solutions (λ, u) of the eigenvalue problem

$$-\Delta u = \lambda f(u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad u \in C^2(\bar{\Omega}),$$

provided f satisfies, instead of (4),

$$(4') \quad \lim_{t \rightarrow 0^+} \frac{f(t)}{t} = 0.$$

Remark 4: Theorem 1 can also be used in order to obtain existence results in the case where $f(0) > 0$ (see [6]).

Proof of Theorem 2: We use the same argument as in [4]. To this end we have only to establish the statement (5) below. We can then apply the proof of [4] (which uses

a simple fixed point argument):

$$(5) \quad \begin{cases} \text{If } (t, u) \text{ satisfy } -\Delta u = f(u) + t\sigma_1 \text{ in } \Omega, u \in C^2(\bar{\Omega}), u \geq 0 \text{ in } \Omega, \\ u = 0 \text{ on } \partial\Omega, t \geq 0 \text{ then } t \leq t_0, \|u\|_{L^\infty(\Omega)} \leq C_0. \end{cases}$$

where σ_1 is chosen as in the proof of Theorem 1.

To prove (5), we consider u a solution of

$$\begin{cases} -\Delta u = f(u) + t\sigma_1 \text{ in } \Omega, \\ u \in C^2(\bar{\Omega}), u \geq 0 \text{ in } \Omega, u = 0 \text{ on } \partial\Omega, \end{cases}$$

with $t \geq 0$. The first step of the proof of Theorem 1 then shows that we have

$$t \leq t_0.$$

Now, the second step of the proof applies, since the results of [5] still hold for u

which is a solution of

$$\begin{cases} -\Delta u = f(x, u) \text{ in } \Omega, \\ u \in C^2(\bar{\Omega}), u \geq 0 \text{ in } \Omega, u = 0 \text{ on } \partial\Omega, \end{cases}$$

with $f(x, u) = f(u) + t\sigma_1(x)$. Indeed we have

$$\frac{\partial \sigma_1}{\partial \nu} \leq -\sigma_1 < 0 \text{ on } \partial\Omega,$$

where ν is the unit exterior normal vector, and this enables us to apply the results and the methods of [5].

REFERENCES

- [1] H. Amann: On the existence of positive solutions of non-linear elliptic boundary value problems, *Ind. Univ. Math. J.* (1971), 125-146.
- [2] A. Ambrosetti and P. H. Rabinowitz: Dual variational methods in critical point theory and applications, *J. Functional Anal.* 14 (1973), 349-381.
- [3] H. Berestycki and P. L. Lions: A local approach to the existence of solutions of semilinear problems in \mathbb{R}^N . To appear.
- [4] H. Brézis and R. E. L. Turner: On a class of superlinear elliptic problems, *Comm. in PDE* 2 (6) (1977), 601-614.
- [5] B. Gidas, Wei-Ming Ni and L. Nirenberg: Symmetry and related properties via the maximum principle. To appear in *Comm. Math. Phys.*
- [6] P. L. Lions, to appear.
- [7] R. Nussbaum: Positive solutions of some nonlinear elliptic boundary value problems, *J. Math. Anal. Appl.* 51 (1975), 461-482.
- [8] S. I. Pohozaev: Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$, *Soviet Math.* 6 (1965), 1408-1411.
- [9] P. H. Rabinowitz: Some critical point theorems and applications to semilinear elliptic partial differential equations. To appear in *Ann. Scuola Norm. Pisa.*
- [10] P. H. Rabinowitz: Variational methods for nonlinear eigenvalue problems, in *Eigenvalues of Nonlinear Problems*, Edizioni Gemonese, Rome (1974), 141-195.
- [11] R. E. L. Turner: A priori bounds for positive solutions of nonlinear elliptic equations in two variables, *Duke Math. J.* 41, No. 4 (1974), 759-774.

PLL/scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1975	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A PRIORI ESTIMATES AND EXISTENCE RESULTS FOR SOME SEMILINEAR ELLIPTIC PROBLEMS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) P. L. Lions		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706 Wisconsin		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE July 1979
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Semilinear problems Symmetry Non-trivial solutions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We study the existence of non-trivial, positive solutions of (1) $\begin{cases} -\Delta u = f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$ We prove the existence of a priori estimates for positive solutions of (1), provided f does not increase too rapidly at infinity and provided Ω has some geometrical properties. From these a priori estimates we deduce the existence of non-trivial, positive solutions of (1).		