

AD-A078 783

PACIFIC MISSILE TEST CENTER POINT MUGU CA
MARINE/CONTINENTAL HISTORY OF AEROSOLS AT SAN NICOLAS ISLAND DU--ETC(U)
APR 79 J ROSENTHAL + T E BATTALINO
PNTC-TP-79-32


F/6 4/2

UNCLASSIFIED

NL

OF |

AD-A078 783



END

DATE

FILED

1-80

BDC



Technical Publication TP-79-32

LEVEL

SUMMARY

MARINE/CONTINENTAL HISTORY OF AEROSOLS AT SAN NICOLAS ISLAND DURING CEWCOM-78 AND OSP III

(PREPARED FOR OPTICAL SIGNATURES PROGRAM (OSP)
NAVAL WEAPONS CENTER, and ELECTRO-OPTICAL
METEOROLOGY (EOMET) PROGRAM, NAVAL OCEAN
SYSTEMS CENTER)


By

J. Rosenthal,
T. E. Battalino, et al

Geophysics Division
Pacific Missile Test Center
Point Mugu, California 93042

and

V. R. Noonkester
Electromagnetic Propagation Division
Naval Ocean Systems Center
San Diego, California 92152



APRIL 1979

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

PACIFIC MISSILE TEST CENTER

Point Mugu, California

DDC FILE COPY



ADA 078783

79 12 27 202

UNCLASSIFIED

PACIFIC MISSILE TEST CENTER

AN ACTIVITY OF THE NAVAL AIR SYSTEMS COMMAND

This report was prepared by J. Rosenthal, T. E. Battalino, and H. Hendon, Geophysics Division, PACMISTESTCEN, and V. R. Noonkester, Electromagnetic Systems Division, NOSC.

Mr. D. A. Lea, Associate Geophysics Officer; Dr. T. C. Lockhart, Associate Range Operations Officer, and Mr. W. L. Miller, Associate Director, Range Directorate, have reviewed this report for publication.

THAD PERRY
Technical Director

J. M. TALLMAN, CAPT USN
Acting Commander Pacific Missile Test Center

Technical Publication TP-79-32

Published by Technical Information Division
Photography and Technical Information Department
Security Classification UNCLASSIFIED
First Printing 280 Copies

UNCLASSIFIED



DEPARTMENT OF THE NAVY

PACIFIC MISSILE TEST CENTER

POINT MUGU, CALIFORNIA 93042

IN REPLY REFER TO

4230

5605

Ser: M-184

10 December 1979

From: Commander, Pacific Missile Test Center

To: Distribution

Subj: Technical Publication TP-79-32; forwarding of

1. Enclosed is a PACMISTESTCEN Summary Document entitled "Marine/Continental History of Aerosols At San Nicolas Island During CEWCOM-78 and OSP III." The report summarizes detailed day by day analyses of meteorological conditions during measurement periods sponsored by the Optical Signatures Program (OSP), NWC, China Lake, CA, and the Electrooptical Meteorology (EOMET) Program, NOSC, San Diego, CA. A brief "EXECUTIVE SUMMARY" is provided at the beginning of this report.

2. A second more complete document of the same title has been published for a limited distribution for those desiring detailed analysis of each day of CEWCOM/OSP III covering 25 April - 1 June 1978. If desired, a copy of this complete document can be obtained by contacting Dr. Jon Wunderlich, OSP Program Manager, Code 39403, Naval Weapons Center, China Lake, CA 93555 or Mr. Jay Rosenthal, Code 3253, Pacific Missile Test Center, Point Mugu, CA 93042.

A. M. Torres

A. M. TORRES

By direction

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TP-79-32	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Marine/Continental History of Aerosols at San Nicolas Island During CEWCOM-78 and OSP III		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) J. Rosenthal, T. E. Battalino, and H. Hendon V. R. Noonkester		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Pacific Missile Test Center Point Mugu, California 93042		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS 123
12. REPORT DATE April 1979		13. NUMBER OF PAGES 32
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Meteorological and optical transmission data being collected at San Nicolas Island, categorized according to periods representing marine, continental or mixed aerosol sources. These data were based on a variety of sources. The aerosol source regions for the OSP III and CEWCOM-78 period were analyzed to perform such categorization. Based on the total		

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. Cont'd.

Integration of data, atmospheric conditions at San Nicolas Island during the CEWCOM/OSP period were marked by great variability.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

TABLE OF CONTENTS

	Page
SUMMARY	1
BACKGROUND	3
ATMOSPHERIC CHARACTERIZATION AT SNI	3
APPROACH	6
EXAMPLE OF ANALYSES: 10 May 1978	8
Synoptic Situation	9
Local Pressure Gradients - An Index of Maritime/Continental Influence	9
Vertical Transport	9
Horizontal Transport	16
SUMMARY/CONCLUSIONS	21
IMPLICATIONS FOR THE REPRESENTATIVENESS OF A MARITIME SITE	22
REFERENCES	24

LIST OF ILLUSTRATIONS

FIGURE		PAGE
1.	Southern California Coastline and Offshore Islands.....	4
2.	San Nicolas Island and the Marine Environment Test Range (METR).....	5
3.	NOAA Pacific Surface Analysis for 1800Z, 10 May 1978.....	10
4.	GOES Satellite Imagery of 2015Z (1215 PST) 10 May 1978 Showing Widespread Cellular Stratus and Frontal Band Off West Coast.....	11
5.	DMSP Satellite Imagery at 1917Z (1117 PST), 10 May 1978 Showing Extensive Stratus and Gravity Wave-induced Clearing Southeast of SNI.....	12
6.	Air Mass Flow Patterns Over Pacific Ocean Inferred from GOES Film Loops, 0006 - 1640Z, 10 May 1978.....	13
7.	Surface Pressure Gradients During CEWCOM, 8 to 17 May 1978....	14
8.	Time Cross-Section of Inversion Height (Shaded Region) Based on SNI Rawinsondes from 8 to 12 May 1978. (Thin Black Lines are Isentropes. Sounding Times (GMT) are Indicated at Bottom.).....	15
9.	Sea Surface Temperature Pattern Based on DMSP Infrared Data, 1947 PST, 11 May 1978.....	17
10.	Approximate Sea Level Air Trajectory to SNI During CEWCOM-78, 10 May 1978.....	18
11.	Marine Layer Trajectories and Flow Patterns Derived from GOES Satellite Film Loops, 10 May 1978.....	19
12.	Analysis of Visibility Patterns Over Coastal Southern California for 1800Z, 10 May 1978.....	20

LIST OF TABLES

TABLE

1.	Data Sources Used to Determine Aerosol History at SNI During CEWCOM/OSP III	7
2.	CEWCOM-78/OSP III Periods of Marine/Continental Influence at San Nicolas Island	21

EXECUTIVE SUMMARY

'Marine/Continental History of Aerosols At San Nicolas Island During CEWCOM-78 and OSP III'

Basic Point: No matter where EOMET measurements are conducted, air mass sources must be determined using techniques like those employed by PACMISTESTCEN in order to interpret marine aerosol data and its implications for electrooptical systems. Air mass differences in aerosol distributions may be much more significant than localized differences due to small scale features or island terrain.

Specific Conclusions:

1. Air mass flow pattern analysis based on actual weather episodes is essential for interpretation of aerosol data and its application to E-O studies.
2. Techniques have been developed by PACMISTESTCEN for doing such analysis using satellite data, weather charts, conventional surface and meteorological profile data, and other data sources considered together.
3. These techniques are being applied to the data at San Nicolas Island to distinguish data taken under maritime conditions from those taken under continental effects.
4. Using these techniques, it was found that during the first two hi-mode measurement periods at San Nicolas (CEWCOM/OSP III and November 1978), considerably more continental influence was experienced than would be expected from long-term climatologies. These were separated by periods of strong maritime influence as expected. During EOMET-79, conditions were maritime much of the time. Measurements conducted at SNI during the winter rainy season, and during summer when low clouds are most frequent should show a higher incidence of maritime conditions since the previous hi-modes were all conducted during spring or fall transition months.
5. Even when air is of mixed origin, as in typical "coastal" environments, atmospheric and transmission data is quite useful because it applies to conditions in which many naval operations are conducted.
6. Some continental influence will likely be experienced at almost any oceanic site of operational relevance, but it is especially likely at American and European mainland coastal sites which are subject to contamination by fresh or aged pollution from local, or remote sources.

PACIFIC MISSILE TEST CENTER
Point Mugu, California 93042

MARINE/CONTINENTAL HISTORY OF AEROSOLS
AT SAN NICOLAS ISLAND DURING
CEWCOM-78 AND OSP III

By

J. Rosenthal and T. Battalino
Pacific Missile Test Center

and

V. R. Noonkester
Naval Ocean Systems Center

Meteorological and optical transmission data being collected at San Nicolas Island must be categorized according to time periods representing marine, continental, or mixed aerosol sources. Based on a variety of data from sources such as surface and upper-air charts, high-resolution GOES and DMSP satellite imagery, climatological data, hourly surface observations, automatic weather station observations, and rawinsonde data, the aerosol source regions for the OSP III and CEWCOM-78 data period were analyzed to perform such categorization. Techniques were developed for assimilating much of this information into summaries or indices which reveal the relative influence of maritime or continental conditions. Air trajectories from low-level winds and satellite cloud monitoring indicated complex circulation patterns making source region identification difficult at times. Some climatological and DMSP sea-surface temperature data were also useful in describing transition conditions in the lower atmosphere near San Nicolas Island. Based on the total integration of data, atmospheric conditions at San Nicolas Island during the CEWCOM/OSP period were marked by great variability, with several episodes of strong maritime influence being separated by periods of continental influence. These changing conditions were created by moving weather systems, with transition periods characterized by air of mixed origins.

Publication UNCLASSIFIED.

Approved for public release; distribution unlimited.

BACKGROUND

Tactical use of Navy weapon systems employing imaging, detection and other electro-optical (EO) sensors, require a predictable knowledge of their effectiveness under various weather conditions experienced over or near the world's oceans. Significant progress has been made by Selby, et al (1978), in developing transmission codes applicable to continental environments with modifications that apply in part to maritime environments. To measure the quality and quantity of atmospheric effects on propagation of energy in the visible, infrared, and ultraviolet parts of the spectrum under the Navy's "real-world" oceanic conditions, a Navy maritime site is required where such transmission conditions can be accurately measured using both laboratory and operational or near-operational systems.

Similarly, to develop Navy models with which specific effects on particular EO systems can be predicted or assessed under "real world" oceanic conditions, a Navy maritime site is also required where the marine atmosphere can be characterized in detail.

Under the joint sponsorship of the Navy's Electro-optical-Meteorology (EOMET) and Optical Signatures Programs (OSP), both of these objectives are being completed at San Nicolas Island (SNI) approximately 100 km off the coast of southern California (figure 1). SNI is part of the Pacific Missile Test Center (PACMISTESTCEN) and as such, is the focal point for Navy development, test and evaluation (DT&E) of missile, aircraft, and weapon systems. The north end of the island on the windward side provides three overwater transmission paths measuring 4.1, 2.5, and 1.6 km (figure 2). Details of this facility are described by Matthews, Williams, et al (1978).

ATMOSPHERIC CHARACTERIZATION AT SNI

SNI was selected as the Navy site for these efforts because it has the best overall combination of attributes which satisfy the requirements for a maritime atmospheric transmission facility. SNI is swept by prevailing northwest winds which have a long fetch across the north Pacific Ocean. With a climate typical of the Mediterranean, the island experiences a variety of weather conditions throughout the year ranging from Pacific storms, rain-bearing fronts, and changing airmasses during the cooler half of the year, to spatial and temporal variations in extensive stratus and low cloud cover during the warmer months. This results in a wide range of atmospheric transparencies as well as wind conditions.

SNI's proximity to the continent also permits air of continental origin to occasionally reach the island whenever winds blow from land to sea. Thus, in terms of meeting the Navy objectives at SNI, it is important to identify and distinguish those periods




Figure 1. Southern California Coastline and Offshore Islands.

when measurements are made at SNI under typically maritime conditions from those when measurements are made under continental or near continental conditions. Only in this way can the results of all measurements at an atmospheric transmission facility like SNI be correctly interpreted.

In characterizing the maritime atmosphere at SNI, it is necessary to concentrate on those elements or characteristics which have the greatest bearing on EO systems performance. These are aerosols, low-level stability, and moisture. At the SNI measurement sites, these parameters are being monitored along with winds, temperatures, and other important parameters by means of automatic weather stations, aerosol collectors, and supplemental sensors. These data are then used to interpret simultaneous transmission measurements across the overwater paths throughout the year.

During several "high-mode" periods and special experiments, detailed meteorological profile and surface measurements of aerosols and atmospheric structure are made from the Naval Research Laboratory's (NRL) Micrometeorological facility at the northern tip of the island (Blanc, 1978), from the Naval Ocean Systems Center's (NOSC) instrumented aircraft (Jensen, 1978), from PACMISTESTCEN's high-resolution rawinsonde soundings, and from many other participants cooperating in the Navy's effort. The first such major experiment at SNI was the Cooperative Experiment of West Coast Oceanography and Meteorology (CEWCOM-78), which generally coincided with a period of extensive target signature and background measurements known as Optical Signatures Program III (OSP III). The total period extended from 25 April to 1 June 1978.

Of all the parameters being measured, the size, distribution, and nature of aerosols are most important because of their scattering and absorption effects on energy. To determine the representativeness of aerosol conditions encountered at SNI to conditions over the open sea, and to correctly interpret the in-situ measurements, it is therefore necessary to determine the marine/continental history of air arriving at SNI. This was accomplished during CEWCOM-78/OSP III through an examination of the large scale flow patterns, wherein airmasses of various origins traverse the northeast Pacific or western North America with SNI in their path.

APPROACH

A wide variety of data from diverse sources were used in the analysis of aerosol history at SNI. These included individual and statistically summarized surface data, upper-air measurements, synoptic weather charts for the surface and higher levels, and high-resolution meteorological satellite imagery. These data sources are listed in table 1.

Table 1. Data Sources Used to Determine Aerosol History at SNI
During CEWCOM/OSP III

Surface
Winds, temperature, humidity, visibility, and cloud cover at Sites A, B, and C Winds, temperature, humidity, visibility, and cloud cover at SNI Airfield Winds, temperature, humidity, visibility, and cloud cover at Mainland Coastal Sites Winds, temperature, humidity, visibility, and cloud cover at Island Sites Marine climatologies of cloud cover, visibility, and sea surface temperature Climatology of winds from SNI Airfield National Meteorological Center (NMC) Surface Weather Charts NRL radon measurements at SNI and aboard NPS Vessel ACANIA Visual observations at SNI
Upper-Air
National Meteorological Center (NMC) Upper-Air Charts at 850, 700, and 500 mb. Fleet Numerical Weather Central (FNWC) Upper-Air Charts for 500 mb GOES Satellite imagery of North Pacific and southern California areas at 30 minute intervals DMSP Satellite imagery PACMISTESTCEN rawinsonde measurements at SNI and Point Mugu Laguna Peak (1,400 feet msl) wind observations

First, surface weather parameters measured during CEWCOM/OSP III were compared with long term climatologies developed by PACMISTESTCEN for the SNI airfield. These comparisons indicated a higher incidence of continental flow for this time of year, overall, than is climatologically expected. Climatological mixing conditions were also inferred which portray the month of May at SNI as a general month of transition.

Next, a comprehensive analysis of meteorological conditions affecting SNI was made for each day of the CEWCOM/OSP III period. This detailed examination later permitted the separation or partition of the measurement period into episodes of maritime, continental, or mixed aerosol history. For each day, synoptic charts and satellite imagery were used to describe the prevailing large scale (synoptic) weather situation affecting SNI. This included the identification of major airmasses, cloud features and coverage, cold fronts, and moving large scale waves (100 - 1,000 mi) which determine SNI weather.

Next surface pressure gradients were calculated between west-east and north-south stations on the mainland, and used as indices of whether the air was flowing from the sea to the land (maritime at SNI) or from the land to the sea (continental at SNI).

Then, indications of vertical transport were considered using time sections of inversion height deduced from rawinsonde data. These were supplemented with important inferences on low-level vertical mixing conditions based on wind observations and DMSP-derived sea surface temperature patterns.

Lastly, horizontal transport conditions were analyzed. This involved principally, the construction and determination of low-level air trajectories ending at SNI for selected days. Two techniques of trajectory analysis were utilized. One involved trajectory computations in the customary manner, using hourly wind observations from coastal and island locations. Based on constructed streamline patterns, air displacements were calculated for hourly intervals extrapolated backward in time from northern SNI as a starting point. The second trajectory method employed cloud motion vectors or displacements for low (marine layer) cloud elements, determined from 16 mm film loops of GOES high resolution satellite imagery. Coverage over the area of interest was at 30 minute intervals showing movements of specific cloud elements or clusters.

Each of the two trajectory techniques has both strong points and weak points. They sometimes produce different trajectories since they are based on different data sources and slightly different altitudes. To aid in the determination of horizontal transport conditions, horizontal visibility analyses (used as an air mass tracer), Laguna Peak wind observations, surface radon measurements (Larson et al, 1979), and visual weather observations (Jeck, 1979, and Fairall et al, 1979), were also employed. When used with the aforementioned analyses of vertical transport, pressure gradients and synoptic weather conditions, a relatively confident determination of maritime versus continental aerosol history over the period of from one to several days could be made for each day of CEWCOM/OSP III at SNI. However, regions of strong subsidence introduced complications to the analysis by bringing air of uncertain origins from high- to low-levels.

EXAMPLE OF ANALYSES: 10 MAY 1978

To illustrate the techniques used in this effort, some results for 10 May 1978 are presented. This was determined to be a day of predominantly maritime conditions at SNI, and most of the data products were available for analysis.

Synoptic Situation

On this day, a strong Pacific anticyclone was located west of California with a general northwesterly flow along the west coast and at SNI. Weakening frontal systems moved towards SNI from the northwest across the top of the Pacific High. The locations of these features are shown in figure 3. The widespread cellular low stratus clouds over the eastern Pacific (near and to the west of SNI) and the weakening frontal cloud band advancing from the northwest are visible in the GOES picture in figure 4. A more detailed view of the stratus cover and its proximity to SNI is shown in the DMSP visual imagery presented as figure 5. This picture shows wave phenomena which reveal the northwest, maritime flow at SNI and daytime clearing to the north and east of SNI.


Based on filmloops of GOES imagery for the entire north Pacific ocean area, a composite analysis of cloud systems and derived flow patterns is provided in figure 6. This composite covers the period from the afternoon of 9 May through the morning of 10 May. It shows the northwest flow and stratus near the west coast, the front moving into the Pacific Northwest, and other higher-altitude features of less interest.

Local Pressure Gradients - An Index of Maritime/Continental Influence

Using surface pressure gradients between San Francisco and Point Mugu (SFO-NTD), Bakersfield and Point Mugu (BFL-NTD), and Tonopah, Nevada and Los Angeles (TPH-LAX), a measure of the tendency for sea to land (maritime) or land to sea (continental) airflow can be inferred. When the pressure differences are negative (higher pressure to the south and west), the tendency for a northwesterly maritime flow at SNI is greatest. The larger the negative value, the greater the maritime tendency and vice versa. As can be seen in the time sections of pressure gradient indices in figure 7, 10 May is a period when two of the three indices are strongly negative and the third reaches zero. In this figure, 10 May is at the center of one of two periods of maritime flow at SNI (the other is centered around 14 and 15 May and are separated by periods of continental or mixed air).

Vertical Transport

To determine the vertical transport conditions in the general vicinity of SNI on 10 May, a time section of inversion height is shown in figure 8 for the period 8 to 12 May based on SNI rawinsonde data. Stability is greatest where isentropes are horizontal and closely packed in the warm inversion layer. This region separates well mixed marine air below within the marine layer from drier (possibly continental) air above. When the marine layer is thick and the inversion strong, little chance exists for continental aerosols mixing down to the surface at SNI. These conditions existed



2015 10MY78 33A-1 00581 22131 SB6




Figure 4. GOES Satellite Imagery of 2015Z (1215 PST) 10 May 1978, Showing Widespread Cellular Stratus and Frontal Band Off West Coast.




Figure 5. DMSP Satellite Imagery at 1917Z (1117 PST), 10 May 1978 Showing Extensive Stratus and Gravity Wave-induced Clearing Southeast of SNI.





Figure 6. Air Mass Flow Patterns Over Pacific Ocean Inferred from GOES Film Loops,
0006 - 1640Z, 10 May 1978.



Dates centered at 12Z (0400 PST)

Figure 7. Surface Pressure Gradients During CEWC0M, 8 to 17 May 1978.

at SNI on 10 May when the marine layer reached a maximum depth of about 1,500 feet. While downward mixing of dry air cannot be precluded far at sea, such air seldom reaches the surface, and would almost certainly assume marine characteristics prior to reaching SNI.

Vertical mixing conditions near SNI can also be inferred by considering the effect of the wind flow superimposed on sea surface temperature patterns. Based on sea surface temperatures analyzed from DMSP infrared imagery on 11 May (figure 9), and assuming that these conditions existed on 10 May as well, SNI is shown to be located just to the west of a region of sharp sea surface temperature contrast separating cool water to the west from warmer water to the east. Such temperature gradients can have an important influence on stability and vertical mixing and therefore on the aerosol size distribution at SNI. On 10 May, since the wind at SNI is from the northwest, any such mixing effects are more likely to be east (downwind) of SNI than at the island itself, so that SNI conditions are relatively representative of conditions further out to sea (upwind to the west).

Horizontal Transport

Trajectories were constructed for 10 May from both surface wind reports (near sea level) and cloud motion vectors (within the marine layer). Figure 10 shows a 5-1/2 hour daytime trajectory, based on wind observations which show an over-water path to SNI with a possible transit over the Point Conception land area prior to entering the starting point of the trajectory. The trajectory also crosses the eastern edge of Santa Rosa Island.

However, the path of marine-layer cloud elements, as seen by satellite (figure 11), indicates an entirely over-ocean trajectory for a slightly longer time period. Also visible in figure 11, are streamlines (thin black arrows) and a second longer trajectory (due to stronger winds) further to the northwest which is in agreement with the general northwest maritime flow concluded to be present at SNI on this day. It should be pointed out that while trajectories of 5 to 10 hours duration may not in themselves be sufficient to determine aerosol or airmass history, continuity and careful monitoring of synoptic factors did permit air source characteristics to be confidently extrapolated further backwards in time in most instances.

One further indication of maritime conditions at SNI on 10 May is provided by an analysis of visibility conditions during mid-morning (figure 12). While such analyses are most questionable over ocean areas due to data voids, careful examination of winds, prevailing visibility, temperature, dewpoints, and hour-to-hour variations permits a reasonable delineation of low visibility areas (due to smog) from high visibility areas (due to clean air of either marine or continental origin), according to wind direction and geographical location. In figure 12, high visibilities within the marine layer are concluded to indicate clean marine air at SNI.




Figure 9. Sea Surface Temperature Pattern Based on DMSP Infrared Data, 1947 PST, 11 May 1978.

"APPROXIMATE SEA LEVEL AIR
TRAJECTORY TO SNI DURING CEWCOM-78"




Figure 10. Approximate Sea Level Air Trajectory to SNI During CEWCOM-78, 10 May 1978.




Figure 11. Marine Layer Trajectories and Flow Patterns Derived from GOES Satellite Film Loops, 10 May 1978.




Figure 12. Analysis of Visibility Patterns Over Coastal Southern California for 1800Z, 10 May 1978.

Additional support for maritime airflow at SNI is available from radon measurements provided by NRL, which indicate very low values on 10 May (Jeck, 1979 and Larson et al, 1979).

SUMMARY/CONCLUSIONS

The complete report summarizes the principal aerosol source regions for SNI during the five-week CEWCOM-78 and OSP III measurement periods (although, only analyses for 10 May 1978 appear in this summary report). It includes detailed day-by-day analyses of the meteorological features and processes responsible for aerosol transport to the island based on a variety of data sources such as surface and upper-air charts, high-resolution GOES and DMSP (satellite) imagery, climatological data, hourly surface observations, automatic weather station observations, rawinsonde data, information measured or recorded at the PACMISTESTCEN and other Navy sources.

Techniques were developed for assimilating much of this information into time cross-section analyses of inversion base height, marine layer conditions, indices of maritime/continental influence, and, most significantly, short-period trajectory analyses to describe or infer the vertical and horizontal transport of aerosols near SNI.

Based on these studies, it was found that the circulation controlling the airflow at SNI during CEWCOM-78/OSP was highly variable with a higher occurrence of offshore (continental) flow than expected based on long-term climatology. Several episodes of strong maritime flow were observed at SNI separated by periods when the air transport to SNI was from the land to sea or continental. These changes were controlled by moving weather systems separated by several days, comprising both synoptic and mesoscale circulations. The transition from one flow pattern to the other was sometimes rapid and there were days when air arriving at SNI had a mixed maritime and continental origin. Table 2 provides a breakdown of conditions during CEWCOM-78/OSP into periods of maritime and continental aerosol history at SNI.

Table 2. CEWCOM-78/OSP III Periods of Marine/Continental Influence at San Nicolas Island

Marine Influence	Continental Influence	Mixed, Modified, or Transition
25 - 27 April	4 May	28 - 30 April
1 - 3 May	11 - 12 May	5 - 7 May
8 - 10 May	17 - 18 May	13 May
14 - 15 May	26 - 29 May	16 May
19 - 24 May		25 May
30 May - 1 June		

Since the measurement period occurred during a month of transition from winter to summer with greater than normal offshore flow and variability in weather, and since aerosol loading at SNI may be affected by underlying gradients of sea surface temperature common to coastal regions, data taken at SNI during the CEWCOM-78/OSP period were at times not representative of open-ocean conditions likely to be found at much greater distances westward from SNI. However, meteorological analyses performed during this period, and observations made in other regions around the world show that during periods of large-scale offshore flow, effects on the marine layer may be observed hundreds of miles out into the open-ocean, so that the airflow at great distances from land are often affected. Thus, under strong offshore flow, conditions at SNI might still approximate some conditions experienced far at sea. During the periods of onshore maritime flow observed during CEWCOM/OSP, data taken at SNI are likely to be substantially representative of open-ocean conditions.

In attempting to delineate the relative influence of maritime or continental effects on aerosols at SNI, two types of trajectory analysis were used, one based on surface wind reports and the other upon satellite-derived cloud motion vectors. Due to data voids and inherent limitations of both techniques, conclusions on source regions did not always agree. To increase the accuracy, and reduce the effort required in this analysis for future measurement periods, techniques are being developed to automate the trajectory analysis process and integrate a variety of data sources into a single product. Thus, techniques can be applied to identify and describe low-level aerosol source regions at a maritime measurement site based on a variety of conventional weather data, satellite observations, in-situ measurements, and calculator-graphical techniques. These can be imperfectly applied to determine the relative maritime/continental influence at the site in question.

IMPLICATIONS FOR THE REPRESENTATIVENESS OF A MARITIME SITE

It should be emphasized that the maritime/continental nature of an ocean site cannot be established solely from geographical considerations and statistical climatologies of surface wind directions. Airmass flow pattern analysis (in 3-dimensions if possible) based on actual synoptic/mesoscale episodes and trajectory analysis is essential. Also of importance to establish the maritime/continental influence at an ocean site is the consideration of many other available parameters, conditions, or indices which are commonly used to describe an air parcel source. These include moisture, wind direction and speed, cloud cover, air source, and length of over-water/over-land trajectory, sea surface temperature patterns as it affects mixing, chemical composition, tracers (radon), aerosol distributions and their vertical profiles, and conditions aloft, as a function of season.

A volume of air is always undergoing modification from above and below as it moves across the Earth's surface; some of these subtle airmass changes may not be indicated by surface winds. Because of these horizontal and vertical transport mechanisms and slow airmass modification, "pure" marine or continental characteristics will seldom be found, even over the middle of the oceans and deserts; instead various degrees of transition and relative marine/continental nature will prevail. Also an EO system performance analysis cannot be based solely on considerations of marine/continental airflow; important mesoscale variations in humidity and wind structure often noted in satellite data must also be considered.

The question of suitability of SNI as a Navy site for EO studies in a maritime environment cannot be answered directly from an analysis of CEWCOM/OSP data. It has been pointed out that this period was marked by large fluctuations between marine and continental airmasses. Subsequent measurements made during November 1978 also show continental contributions, but measurements during April/May 1979 indicated a higher persistency of maritime conditions at SNI. Since all three measurement periods occurred during the transition seasons of spring or fall, an even higher incidence of maritime conditions can be expected for measurements conducted in winter when Pacific storms are most frequent, or during summer when stable marine layer conditions are nearly a daily occurrence at SNI. It appears that SNI is a good approximation to an open-ocean environment during periods of strong or persistent maritime flow, and unrepresentative of an open-ocean environment during periods of strong offshore flow. In between or during periods of transition conditions, SNI may be representative of a coastal environment, - an environment where many Naval operations are typically conducted. SNI appears to be distinctly more maritime than would be encountered on a populated mainland coast or larger island, particularly when compared with American and European coastline locations which are heavily contaminated by fresh and aged urban pollution.

REFERENCES

- Blanc, T. V., "Micrometeorological data for the 1978 Cooperative Experiment for West Coast Oceanography and Meteorology (CEWCOM-78) at San Nicolas Island, California", Naval Research Laboratory Report (NRL) Memorandum Report 3871, Vol. I, II, August 1978.
- Fairall, C.W.; G. E. Schacher; K. L. Davidson/and T. M. Houlihan, "Atmospheric Marine Boundary Layer Measurements in the Vicinity of San Nicolas Island During CEWCOM-78" Naval Postgraduate School, NPS 61-78-007, September 1978.
- Jeck, R. "Aerosol Particle Size Measurements at San Nicolas Island During CEWCOM-78", Naval Research Laboratory, Washington, D.C., NRL Memorandum Report 3931, 9 March 1979.
- Jensen, D. R., Aerosol-size distribution measurements in the marine boundary layer: OSP III and CEWCOM-78 data report, Naval Ocean Systems Center Technical Report TN 560, 15 November 1978.
- Larson, R. E.; W. Kasemir; and D. J. Bressan, "Measurements of Atmospheric ^{222}Rn at San Nicolas Island and Over Nearby California Coastal Areas During CEWCOM-78", NRL Memo Report 3941, Naval Research Lab., 1979.
- Matthews, G.; A. Akkerman; B. E. Williams; R. deViolini; and J. Rosenthal, "Atmospheric Transmission and Supporting Meteorology in the Marine Environment at San Nicolas Island", Semi-Annual Report, Jan 1978 through August 1978, Pacific Missile Test Center, PACMISTESTCEN Technical Publication No. TP-79-19, December 1978.
- Selby, J.; E. P. Shettle; and R. McClatchey, "Atmospheric Transmittance from 0.25 to 28.5 μm ", Supplement LOWTRAN 3B (1976), AFGL-TR-76-0258, Air Force Geophysics Lab, Hanscom AFB, Mass., 1 November 1978.

INITIAL DISTRIBUTION

EXTERNAL	Copies	EXTERNAL	Copies
Commander		Commanding Officer	
Naval Air Systems Command		Naval Research Laboratory	
Headquarters		Washington, DC 20375	
Washington, DC 20361		Attn: Code 1408 (N. Balthasar)	1
Attn: AIR-03P2 (J. Malley)	1	Code 1409 (J. MacCallum)	1
AIR-370 (K. Van Sickle)	1	Code 5553F (D. Friedman)	1
(M. Schefer)	2	Code 5560 (P. Ulrich)	1
AIR-5332 (M. Whiting)	1	Code 5567 (G. Trusty)	1
AIR-5333 (E. Cosgrove)	1	(J. Dowling)	1
AIR-53631F (D. Caldwell)	1	Code 5754 (E. Alexander)	1
AIR-954	2	Code 8320 (H. Gerber)	1
		(L. Ruhnke)	1
Defense Documentation Center		(T. Blanc)	1
Cameron Station		Code 8323 (R. Larson)	1
Alexandria, VA 22314		(R. Jeck)	1
Attn: TIOG	2	Code 8326 (J. Fitzgerald)	1
Commander		Code 8327 (S. Gathman)	1
Naval Material Command			
Headquarters		Commanding Officer	
Washington, DC 20360		Naval Environmental Prediction	
Attn: Code 08T22 (R. Gaylord)	1	Research Facility	
Code 08T2211 (C. Spinder)	1	Monterey, CA 93940	
Code 08T245 (G. Spalding)	1	Attn: A. Weinstein	1
		A. Gorogh	1
Commander		R. Fett	1
Naval Weapons Center		R. Nagle	1
China Lake, CA 93555			
Attn: Code 3173 (A. Shlanta)	1	Officer in Charge	
Code 39403 (T. Smith)	1	White Oak Laboratory	
(J. Wunderlich)	25	Naval Surface Weapons Center	
(L. Wilkins)	1	Silver Spring, MD 20910	
Code 3941 (D. Kummer)	1	Attn: Code R42 (B. Katz)	1
Code 533 (Library)	1	Code 213 (A. Hirshman)	1
Commander		Commander	
Naval Sea Systems Command		Naval Electronic Systems Command	
Headquarters		Headquarters	
Washington, DC 20362		Washington, DC 20360	
Attn: NAVSEA-03 (S. Marcus)	1	Attn: ELEX 035 (R. Golding)	1
PMS 405M (L.T.G. W. Levy)	1	ELEX 3102 (M. Parker)	1
Code 03415 (L. Tasaka)	1	PME 107 (J. Obrian)	1
		PME 107-53 (A. Ritter)	1
Commander		Commander	
Naval Materiel Command		Naval Ocean Systems Center	
Washington, DC 20360	1	San Diego, CA 92152	
High Energy Laser Program		Attn: Code 532 (J. Richter)	10
Washington, DC 20360		(D. Jensen)	1
Attn: Code PM-27	1	(H. Hughes)	1
		(V. Noonkester)	20

EXTERNAL	Copies	EXTERNAL	Copies
Chief of Naval Research 800 North Quincy Street Arlington, VA 22217		Commander Atmospheric Sciences Laboratory White Sands, NM 88002	
Attn: Code 221 (D. Lewis)	1	Attn: DELAS-AS (R. Bonner)	1
Code 421 (W. Condell)	1	(H. Holt)	1
Code 465 (J. Hughes)	1	DELAS-EO-MO (Dr. Gomez)	1
Superintendent		(J. Lindberg)	1
Naval Postgraduate School		(F. Niles)	1
Monterey, CA 93940		ERADCOM (T. Hall)	1
Attn: Code 61CR (A. Cooper)	1	Commander	
Code 61CT (E. Crittenden)	1	Air Force Avionics Laboratory	
Code 61SG (G. Schacher)	1	Wright-Patterson AFB	
Code 63DS (K. Davidson)	1	Dayton, OH 45433	
(D. Leipper)	1	Attn: RWI-3(WE) (CAPT B. Smith)	1
(C. Fairall)	1	RSP-5 (CAPT J. Pryce)	1
(T. Houlihan)	1	WRP-1 (R. Sanderson)	1
(R. Renard)	1	Commander	
(T. Williams)	1	Air Force Geophysics Laboratory	
Commanding Officer		Hanscom AFB, MA 02173	
Naval Ocean Research and Development Activity		Attn: OPA (R. Fenn)	1
National Space Technology Laboratories		(E. Shettle)	1
Bay St. Louis, MI 39520		OPI (R. McClatchey)	1
Attn: R. Goodman	1	OPR (B. Sandford)	1
Office of the Under Secretary of Defense (DR&E)		Commander	
The Pentagon		Army Electronics Command	
Washington, DC 20350		Night Vision Laboratory	
Attn: MAES (COL E. Friday)	1	Fort Belvoir, VA 22060	
(R. Davis)	1	Attn: DELNI-VI (R. Bergman)	1
Institute for Defense Analysis		(R. Moulton)	1
400 Army Navy Drive		Desert Research Institute	
Arlington, VA 22202		University of Nevada	
Attn: (L. Biberman)	1	Reno, NV 89507	
(R. Roberts)	1	Attn: J. Hudson	1
(H. Wolfhard)	1	Grumman Aerospace Corporation	
Massachusetts Institute of Technology		Research Department	
Lincoln Laboratory		Bethpage, NY 11743	
P.O. Box 73		Attn: Dr. J. Selby	1
Lexington, MA 02173		Science Applications, Inc.	
Attn: Dr. Keinman	1	1200 Prospect Street	
		P.O. Box 2351	
		La Jolla, CA 92038	
		Attn: Dr. M. Griggs	1

EXTERNAL	Copies	EXTERNAL	Copies
University of California at Los Angeles (UCLA) Department Atmospheric Sciences 405 Hilgard Avenue Los Angeles, CA 90024 Attn: (J. Edinger) (M. Wurtele)	1 1	Commander U.S. Army Missile Research Laboratory Advanced Sensors Directorate Restone Arsenal, AL 35809 Attn: Code DRDMI-TEI (H. Jackson)	1
Calspan Corporation Advanced Technology Center Buffalo, NY 14225 Attn: E. J. Mack T. A. Niziol	1 1	Defense Advanced Research Projects Agency 1400 Wilson Boulevard Arlington, VA 22209 Attn: S. Zakanyez	1
Megatek Corporation 1055 Shafter San Diego, CA 92117	1	Environmental Research Institute of Michigan P.O. Box 8618 Ann Arbor, MI 48107 Attn: R. Legault	1
Chief of Naval Operations-Pentagon Washington, DC 20350 Attn: Code CNO-982F3 (Capt Pellock) Code CNO-35E (L. Triggs)	1 1	Aeronautical Research Associates of Princeton 50 Washington Road Princeton, NJ 08540	1
David Taylor Naval Ship R&D Center Annapolis Laboratory Annapolis, MD 21402 Attn: Code 2833 (R. Burns)	1	Commanding Officer Naval Intelligence Support Center 4301 Suitland Road Washington, DC 20390 Attn: Code NISC-40 (P. Roberts) Code NISC-50 (R. Nelson)	1 1
Commander Air Force Armament Laboratory Eglin AFB, FL 32542 Attn: Code DLMT (M. Gay)	1	AIL, Division of Cutter-Hammer Melville, Long Island, NY 11746 Attn: A. DiNardo	1
Commander Naval Air Systems Command Headquarters Jefferson Plaza Arlington, VA 22212 Attn: Code PMA-262 (CDR J. Quinn)	1	Northrop Corporation, Aircraft Group 3901 West Broadway Hawthorne, CA 90250 Attn: L. Tanabe	1
Office Under Secretary of Defense Research & Engineering-Pentagon Washington, DC 20301 Attn: Code EPS (E. Myers)	1	Assistant Secretary of Navy Research Engineering and Systems The Pentagon Washington, DC 20350 Attn: Code 5 E 813 (Dr. Wang)	1
General Research Corporation SWL Division 7926 Jone Branch Drive McLean, VA 22101 Attn: R. Rollins	1	Commander Naval Air Systems Command JP Building #1 Arlington, VA 20360 Attn: Code AIR-630E (V. Tarulis)	1

INTERNAL	Copies	INTERNAL	Copies
Commander PACMISTESTCEN		Range Operations Department	
Code 0000		Code 3200	
RADM F. H. Baughman	1	CDR J. P. O'Leary	1
Technical Director		Dr. T. C. Lockhart	1
Code 0002		Code 3250	
Thad Perry	1	CDR J. Tupaz	1
Advanced/Special Projects		D. A. Lea	1
Code 0140		Code 3251	
CDR J. T. Larsen	1	AGCS R. Hicks	1
Code 0141		Code 3252	
W. Luttrell	1	R. Hixon	1
R. Diehl	1	R. Rutt	1
C. Elliott	1	W. Klemz	1
Systems Evaluation Directorate		Code 3253	
Code 1001		J. Rosenthal	20
CAPT R. L. Berg	1	R. de Violini	3
Systems Technology Department		B. Williams	3
Code 1200		T. Battalino	20
K. I. Lichti	1	R. A. Helvey	1
Code 1230		Y. K. Yamamura	1
M. R. Marson	1	R. Clark	1
Code 1232		R. Dixon	1
R. A. Mark	1	Code 3254	
G. B. Matthews	1	L. Hall	1
A. Akkerman	1	A. Bardusch	1
Fleet Weapons Engineering Directorate		M. Bahu	1
Code 2002		Engineering Applications Directorate	
G. R. Wachold	1	Code 4001	
Range Directorate		R. S. Walker	1
Code 3001		Technical Information Division	
CAPT H. M. J. Lewis, Jr.	1	Code 4230	
W. L. Miller	1	A. M. Torres	1
Range Management		Code 4230-1	
Code 3020		M. F. Hayes	30
F. Roth	1	Technical Reports Branch	
G. Keenan	1	Code 4231	
Capabilities Development Department		W. J. Kirkpatrick	1
Code 3140		Library Systems Office	
T. Carr	1	Code 6862	
Code 3143		Technical Reports Library	2
M. Bondelid	1	Patent Counsel	
J. Loos	1	Code PC	
		Dr. J. M. St. Amand	1




Figure 11. Marine Layer Trajectories