

ADA 079324

(12) LEVEL II

PITTSBURGH ATOMIC SCIENCES INSTITUTE

9 TECHNICAL PROGRESS SUMMARY NO. 26

ORDER NUMBER:

The Advanced Research Projects Agency  
ARPA Order No. 2686

CONTRACT NUMBER:

15 N00014-76-C-0098

PROGRAM CODE NUMBER:

6E20

NAME OF CONTRACTOR:

Department of Physics and Astronomy  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

DATE OF CONTRACT:

July 1, 1975

EXPIRATION DATE:

May 15, 1980

AMOUNT OF CONTRACT:

\$604,983

PROJECT SCIENTIST:

M. A. Biondi, Director  
Professor of Physics

Telephone: Area Code 412  
624-4354

SCIENTIFIC OFFICER:

Code N66014

TITLE OF WORK:

ATOMIC AND MOLECULAR PROCESSES

This research was supported by the Advanced Research Projects Agency of the Department of Defense and was monitored by ONR under Contract No. N00014-76-C0098.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Projects Agency or the U. S. Government.

DDC  
RECEIVED  
JAN 11 1980  
B

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

October 31, 1979

79 11 28 039

408 999

DDC FILE COPY

Pittsburgh Atomic Sciences Institute

Faculty Members:

J. N. Bardsley  
M. A. Biondi

Post-Doctoral Personnel:

R. Johnsen  
J. M. Wadehra

Graduate Research Assistants:

V. Jog  
J. Macdonald  
M. Whitaker

|                                 |               |                                     |
|---------------------------------|---------------|-------------------------------------|
| ACCESSION for                   |               |                                     |
| NTIS                            | White Section | <input checked="" type="checkbox"/> |
| DDC                             | Buff Section  | <input type="checkbox"/>            |
| UNANNOUNCED                     |               | <input type="checkbox"/>            |
| JUSTIFICATION                   |               |                                     |
| <b>PER LETTER</b>               |               |                                     |
| BY                              |               |                                     |
| DISTRIBUTION/AVAILABILITY CODES |               |                                     |
| Dist.                           | AVAIL. and/or | SPECIAL                             |
| <b>A</b>                        |               |                                     |

# I. Experimental Investigations of Laser Processes

## A. Electron-Ion Recombination under Laser Plasma Conditions (M. A. Biondi)

The 3-mode microwave afterglow apparatus and ultrahigh vacuum gas handling system was partially disassembled. Modifications were made to the vacuum system consisting of additional ultrahigh vacuum valves and stainless steel refrigerated traps to permit controlled vapor pressures of mercury and mercury compounds to be introduced to the quartz plasma-afterglow cylinder in the microwave cavity. In addition, desired amounts of noble gases (e.g. helium and xenon) can be introduced to the plasma region for control of electron heating/cooling rates and ion-molecule reaction paths.

Initial studies of mercury afterglows, with helium used as the inert buffer gas to control electron energy gain/loss rates and reduce ambipolar diffusion loss, were unsuccessful in achieving conditions where  $\text{Hg}_2^+$  ions dominated the afterglow for the desired recombination studies. As a result, a more complex ion generation sequence, based on the ion-molecule reaction studies described in Sec. I.B, was used to make  $\text{Hg}_2^+$  ions the principal afterglow ions. Xenon was added to the helium and mercury vapor, so that the ion generation/conversion sequence was  $\text{Xe}^+ \rightarrow \text{Xe}_2^+ \rightarrow \text{Hg}^+ \rightarrow \text{Hg Xe}^+ \rightarrow \text{Hg}_2^+$ .

At present, studies of the electron temperature ( $T_e$ ) dependence of the recombination coefficient  $\alpha(\text{Hg}_2^+)$  are in progress. The value at 300 K,  $\alpha \sim 5 \times 10^{-7} \text{ cm}^3/\text{sec}$ , agrees reasonably with earlier measurements at thermal energy. The excited state production by dissociative recombination is presently under investigation over the electron temperature range,  $300 \text{ K} < T_e < 10,000 \text{ K}$ . One of the principal products of recombination over the whole electron temperature range is the  $7^3\text{S}$  state, as evidenced by the  $5461 \text{ \AA}$  radiation from the afterglow.

### B. Ion-Molecule Reactions of Laser Interest (R. Johnsen)

The drift tube-mass spectrometer apparatus was modified to permit studies of ion-molecule reactions involving mercury or mercury compounds. Initially, the modified apparatus has been applied to studies of atomic and molecular rare gas ions with mercury atoms, usually via the additional residence time technique but in the case of the molecular ions by the change in arrival time spectra of the parent ions.

The results for the rare-gas atomic ions are summarized in Table I. Only

Table I. Rare-gas atomic-ion reactions with mercury at thermal energy

| <u>REACTION</u>                                               | <u>RATE COEFFICIENT</u><br><u>(cm<sup>3</sup>/sec)</u> |
|---------------------------------------------------------------|--------------------------------------------------------|
| $\text{He}^+ + \text{Hg} \rightarrow \text{He} + \text{Hg}^+$ | 2.5 (-9)                                               |
| $\text{Ne}^+ + \text{Hg} \rightarrow \text{Ne} + \text{Hg}^+$ | < 5 (-13)                                              |
| $\text{Ar}^+ + \text{Hg} \rightarrow \text{Ar} + \text{Hg}^+$ | < 5 (-13)                                              |
| $\text{Kr}^+ + \text{Hg} \rightarrow \text{Kr} + \text{Hg}^+$ | < 1 (-12)                                              |
| $\text{Xe}^+ + \text{Hg} \rightarrow \text{Xe} + \text{Hg}^+$ | < 1 (-13)                                              |


the  $\text{He}^+ + \text{Hg}$  reaction is fast; the others are undetectably slow.

The molecular rare-gas ion reactions with mercury are tabulated in Table II; two are undetectably slow.

Table II. Rare-gas molecular-ion reactions with mercury at thermal energy

| <u>REACTION</u>                                                   | <u>RATE COEFFICIENT</u><br><u>(cm<sup>3</sup>/sec)</u> |
|-------------------------------------------------------------------|--------------------------------------------------------|
| $\text{He}_2^+ + \text{Hg} \rightarrow 2 \text{He} + \text{Hg}^+$ | 4.5 (-10)                                              |
| $\text{Ne}_2^+ + \text{Hg} \rightarrow 2 \text{Ne} + \text{Hg}^+$ | 5 (-11)                                                |
| $\text{Ar}_2^+ + \text{Hg} \rightarrow 2 \text{Ar} + \text{Hg}^+$ | < 1 (-12)                                              |
| $\text{Kr}_2^+ + \text{Hg} \rightarrow 2 \text{Kr} + \text{Hg}^+$ | < 1 (-12)                                              |
| $\text{Xe}_2^+ + \text{Hg} \rightarrow 2 \text{Xe} + \text{Hg}^+$ | 5 (-10)                                                |


In addition, the reaction



proceeds with a rate coefficient  $k \approx 6 \times 10^{-10} \text{ cm}^3/\text{sec}$  at 300 K.

The xenon-mercury reaction sequence leading to production of molecular ions of mercury is found to involve a combination of fast 2-body and moderately fast 3-body reactions, as indicated in Table/Figure III.

Table/Figure III. Xenon-mercury reactions



At present the reactions involving Hg Br<sub>2</sub> molecules are under study and preliminary rate coefficient values have been obtained. It is expected that final values will be obtained during the next report period.

### III. Theoretical Investigations of Laser Processes

#### A. Ion-ion recombination (J. N. Bardsley)

Our basic computer program to evaluate the rate of ion-ion recombination at intermediate pressures is complete, and has been applied to rare-gas halide systems at several values of temperature and pressure.

We have confirmed our earlier speculation that an increase in the ambient temperature leads to a significant lowering of the recombination rate at low pressure but has little effect at high pressure. Due to these changes the peak in the recombination rate is pushed to higher pressure and is reduced in magnitude.

During this summer, in collaboration with B. Whitten of Miami University of Ohio, we began to investigate the dependence of the recombination rate upon the charge density by examining the effects of screening upon the recombining ion pair. The application of the Debye shielding theory suggests that this effect may lead to a reduction in the recombination rate for charge densities appropriate to high-power lasers or disturbed atmospheres.

Modelling calculations for rare-gas halide lasers, performed by L. Morgan at Livermore, suggest that the recombination rates may be significantly lower than those predicted by the chemical theories. We are planning to make one code available to them and to perform further analysis to check whether these preliminary results are consistent with our theory. This analysis will involve a more comprehensive study of screening effects.

B. Electron collisions with molecules (J. N. Bardsley)

We are adapting our resonance scattering code to permit us to make an analysis of dissociative attachment to  $\text{HCl}$ . Recent experimental results from Yale University suggest that the cross section may be very sensitive to the rotational and vibrational state of the  $\text{HCl}$  molecule. The necessary modifications to the code have been completed but have not yet been fully tested. We are currently developing the theoretical model on which our computations will be based.

C. Ion-molecule reactions (J. N. Bardsley)

The analysis of the response of the atmosphere to intense beams requires an understanding of the rates, over a broad range of temperatures, for many ion-molecule reactions. As part of our collaboration with the NOAA Aeronomy Group at Boulder we have completed a study of charge transfer from  $\text{O}^+$  in collisions with  $\text{O}_2$ . This work has helped to resolve the large differences between beam measurements of this reaction rate.

#### IV. Publications and Technical Presentations

##### A. Publications

Resonant Contributions to Single Charge Transfer between  $\text{He}^{2+}$  and He, J. N. Bardsley, James S. Cohen and J. M. Wadehra, Phys. Rev. A 19, 2129 (1979).

Molecular Resonance Phenomena, J. N. Bardsley, In Electron-Molecule and Photon-Molecule Collisions, Eds. T. Rescigno, V. McKay and B. Schneider (Plenum, 1979), 267.

The Mobility of  $\text{He}^+$  Ions in He, S. Sinha, S. L. Lin and J. N. Bardsley, J. Phys. B 12, 1613 (1979).

Cross Section for the Reaction  $\text{O}^- + \text{O}_2 \rightarrow \text{O}_2^- + \text{O}$  at Relative Kinetic Energies 0.04 - 2 eV, S. L. Lin, J. N. Bardsley, I. Dotan, F. C. Fehsenfeld and D. L. Albritton, Int. J. Mass. Spectrom. Ion Phys., in press.

Thermal-Energy Charge Transfer, Quenching and Association Reactions of Doubly Charged Ions in the Rare Gases, Rainer Johnsen and Manfred A. Biondi, Phys. Rev. A 20, 87, 1979.

##### B. Technical Presentations

Cross Section for the Reaction  $\text{O}^- + \text{O}_2 \rightarrow \text{O}_2^- + \text{O}$  at Relative Kinetic Energies 0.04-2 eV, presented by J. N. Bardsley at the Gaseous Electronics Conference, October 1979.

Charge-Transfer and Association Reactions Involving Mercury and Rare-Gas Ions, presented by R. Johnsen at the Gaseous Electronics Conference, October 1979.

#### V. Visiting Scientists

B. Whitten, Miami University of Ohio.

Senior InvestigatorEst. Funds Expended and Committed (Thous.)

|                                         |              |
|-----------------------------------------|--------------|
| J. N. Bardsley                          | 122.0        |
| M. A. Biondi                            | 230.0        |
| W. L. Fite                              | 56.7         |
| F. Kaufman                              | 147.4        |
| E. C. Zipf                              | <u>43.5</u>  |
| Total Expended and Committed            | 599.6        |
| Available Funds                         | <u>604.9</u> |
| Estimated Funds Remaining as of 9/30/79 | 5.3          |

---

## University Accounting of Funds

|                               |              |
|-------------------------------|--------------|
| Expended as of 9/30/79        | 599.6        |
| Available Funds               | <u>604.9</u> |
| Remaining Funds as of 9/30/79 | 5.3          |