

AD-A079 517

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF MATHEMATICS F/G 12/1
SOME PROPERTIES OF BEST LINEAR UNBIASED PREDICTORS AND RELATED --ETC(U)
NOV 79 A CABANNES

N00014-75-C-0555


UNCLASSIFIED

TR-16

NL

1 OF /
AD
A079517





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 079517

DDC FILE COPY

LEVEL II

12



SOME PROPERTIES OF BEST LINEAR
UNBIASED PREDICTORS AND RELATED
PREDICTORS

by

André Cabannes
DEPARTMENT OF MATHEMATICS
MASSACHUSETTS INSTITUTE OF TECHNOLOGY

TECHNICAL REPORT NO. 16

NOVEMBER 1979



PREPARED UNDER CONTRACT
N00014-75-C-0555 (NR-042-331)
FOR THE OFFICE OF NAVAL RESEARCH

DEPARTMENT OF MATHEMATICS
MASSACHUSETTS INSTITUTE OF TECHNOLOGY
CAMBRIDGE, MASSACHUSETTS

79 12 28 025

SOME PROPERTIES OF BEST LINEAR
UNBIASED PREDICTORS AND RELATED
PREDICTORS

by

André Cabannes
Massachusetts Institute of Technology

ABSTRACT

The variation of temperature or of pollutant concentrations over a geographic area are adequately represented by random fields. Given a real-valued random field $\{Z(x), x \in \mathbb{R}^2\}$ a basic problem is to interpolate Z over an area A from measurements taken at n stations $x_1, x_2, \dots, x_n$, when the distribution of Z is only partially specified. This is the motivation of the present paper.

It is shown that if the joint distributions are Gaussian the best linear unbiased predictor is (among other properties) admissible when used to predict Z at a single point, but inadmissible in general when used to predict the values of the field at several points. A Stein-like predictor is produced which is uniformly better than the B.L.U.E. in the latter case. A nonlinear predictor, based on relaxing the unbiasedness condition on the B.L.U.E., is also proposed and shown to be in some cases preferable.

Key words: Best linear unbiased predictor, multivariate normal distribution, admissibility, James-Stein estimation.

Accession For	
NTIS Grant	☒
DIC TAB	☐
Unannounced	☐
Justification	☐
Dist	Available/or special
A	

See 1473 in back

1. Introduction and summary:

The questions considered in this paper arise naturally when trying to interpolate a realization of a random field $\{Z(x), x \in \mathbb{R}^2\}$ from observations $Z(x_1), Z(x_2), \dots, Z(x_n)$ made at n points, for example interpolation of air pollutant concentration field from fixed monitoring stations. The first step is to study how to interpolate at a single point x ; and then, to study how to interpolate simultaneously at N points forming a fine grid, say, over an area of interest.

However in the present paper we study the abstract prediction problem, independently of the geometry of the underlying set of the random field. So consider $n+1$ real valued random variables $Z_1, Z_2, \dots, Z_n$ and Z with a joint distribution whose covariance matrix is specified and whose mean vector is the product of a specified matrix by an unspecified p -dimensional vector parameter a .

First of all, in Section 2, we are interested in the properties of the best linear unbiased predictor $\hat{Z}$ of Z based on the observed values of $Z_1, Z_2, \dots, Z_n$, when the risk function is $R(a, \hat{Z}) = E(\hat{Z} - Z)^2$. We show that if the joint distribution of $Z_1, Z_2, \dots, Z_n$ and Z is normal, then $\hat{Z}$ is extended Bayes, minimax, admissible and uniformly minimum variance unbiased.

Then in Section 3 we consider the problem of predicting

simultaneously N random variables $z_{n+1}, z_{n+2}, \dots, z_{n+N}$ (instead of a single Z) on the basis of the observed values of $z_1, z_2, \dots, z_n$. Under the risk function

$$\sum_{i=1}^N E(\hat{z}_{n+i} - z_{n+i})^2$$

we show that the pointwise use of the best linear unbiased predictor of Section 2 for each z_i , $i = n+1, \dots, n+N$ is not an admissible procedure in general under normality assumption. This is done by constructing a Stein-like predictor $(\tilde{z}_{n+1}, \tilde{z}_{n+2}, \dots, \tilde{z}_{n+N})$ for $(z_{n+1}, z_{n+2}, \dots, z_{n+N})$. A necessary and sufficient condition for the Stein-like predictor to be uniformly better than the simultaneous pointwise best linear unbiased predictor $(\hat{z}_{n+1}, \hat{z}_{n+2}, \hat{z}_{n+N})$ is given. It is derived as a consequence of the following result (proved in Section 3): If X is a p -dimensional normal random vector with distribution $N(\theta, I_p)$ and C is a fixed $N \times p$ matrix then $\exists \eta > 0$ such that $\forall \theta$

$$(1) \quad E \left\| CX \left(1 - \frac{n}{\|X\|^2} \right) - C\theta \right\|^2 < E \|CX - C\theta\|^2$$

if and only if

$$(2) \quad \max \text{ eigenvalue of } CC' < \frac{1}{2} \text{tr } CC'$$

This theorem extends the basic result of James and Stein (1961) which says that in the case $C = I_p$ the N.S.C. for (1) to hold is $p \geq 3$.

Finally, in Section 4, going back to the problem of predicting a single Z , we present a biased non-linear predictor Z^* obtained as a modification of $\hat{Z}$ when the unbiasedness condition is relaxed. We compute the risk function of Z^* and conclude that Z^* should be preferred to $\hat{Z}$ in some cases when the unspecified vector parameter a can be bounded.

2. The best linear unbiased predictor of a single random variable and its properties in the normal case:

Preliminaries:

Let $\underline{Z}$ denote the column vector formed by the observed values $(Z_1, Z_2, \dots, Z_n)$. Then the assumptions described in Section 1 can be written as follows:

$$E \underline{Z} = F'a, \quad \text{Cov } \underline{Z} = K$$

$$E Z = f'a, \quad \text{var } Z = \sigma^2$$

$$\text{Cov}(\underline{Z}, Z) = k$$

$$F \text{ (p} \times \text{n matrix), } f \text{ (p} \times \text{1), } K \text{ (n} \times \text{n), } k \text{ (n} \times \text{1) ,}$$

and σ^2 (scalar) are specified quantities.

a (p×1 vector) is an unspecified vector-parameter.

The best linear unbiased predictor

$$\hat{Z} = \lambda_0 + \lambda_1 Z_1 + \lambda_2 Z_2 + \dots + \lambda_n Z_n \quad (\text{that we write in short}$$

$\hat{Z} = \lambda_0 + \lambda' \underline{Z}$) of Z is obtained by solving the following system

$$\begin{cases} E(\hat{Z} - Z)^2 \text{ minimum} & 2.1.a \\ E \hat{Z} = f'a \text{ for all } a & 2.1.b \end{cases}$$

The minimization, in 2.1.a, a priori depends on a , but

condition 2.1.b. implies $\lambda_0 = 0$, $F\lambda = f$, and $E(\hat{Z}-Z)^2$ independent of a . The solution, which is classical, is

$$\lambda = K^{-1}k - K^{-1}F'(FK^{-1}F')^{-1}FK^{-1}k + K^{-1}F'(FK^{-1}F')^{-1}f \quad 2.2$$

and the mean square error of the best $\hat{Z}$ is

$$E(\hat{Z}-Z)^2 = \text{Var}(\hat{Z}-Z) = \sigma^2 - k'K^{-1}k + (FK^{-1}k - f)'(FK^{-1}F')^{-1}(FK^{-1}k - f) \quad 2.3$$

For subsequent use of these unwieldy formulas we introduce the following notation: $G = FK^{-1}F'$ (we assume it is non singular), and $\phi = FK^{-1}k - f$.

$\hat{Z}$ can be given a simple interpretation (Goldberger, 1962): since $\underline{Z} = F'a + \varepsilon$ where $E\varepsilon = 0$ and $\text{Cov } \varepsilon = K$. The generalized least square estimator of a based on $\underline{Z}$ is $\hat{a} = (FK^{-1}F')^{-1}FK^{-1}\underline{Z} = G^{-1}FK^{-1}\underline{Z}$ and the best linear unbiased predictor of Z can be rewritten

$$\hat{Z} = f'\hat{a} + k'K^{-1}[\underline{Z}-F'\hat{a}] \quad 2.4$$

Formula 2.4, in view of the best linear predictor

$$\hat{Z} = f'a + k'K^{-1}[\underline{Z}-F'a] \quad 2.5$$

of Z when a is specified, shows the natural origin of $\hat{Z}$.

Finally let us observe that $E(\hat{Z}-Z)^2 = \sigma^2 - k'K^{-1}k + \phi'G^{-1}\phi$ and $E(\tilde{Z}-Z)^2 = \sigma^2 - k'K^{-1}k$. Therefore $\phi'G^{-1}\phi$ represents the price we pay for not knowing a , while requiring unbiasedness of $\hat{Z}$. In Section 3 and Section 4 we show how this added term in the mean square error may be reduced.

Properties of $\hat{Z}$ in the normal case:

In addition to the assumptions laid out at the beginning of this Section, we now assume that Z and $\underline{Z}$ are jointly multivariate normal.

First of all suppose that the vector parameter a has the multivariate normal prior distribution $N(a, \Gamma)$

Lemma 2.1: Let $\hat{Z}_{\alpha, \Gamma}$ be the Bayes rule using mean squared error to predict Z , when we observe $\underline{Z}$, and when $a \sim N(a, \Gamma)$. It has the form

$$\hat{Z}_{\alpha, \Gamma} = k'K^{-1}\underline{Z} - \phi'E(a|\underline{Z}) \quad 2.6$$

and its Bayes risk is

$$r(\hat{Z}_{\alpha, \Gamma}) = \sigma^2 - k'K^{-1}k + \phi'(G+\Gamma^{-1})^{-1}G(G+\Gamma^{-1})^{-1}\phi \quad 2.7$$

Proof: The Bayes rule $\hat{Z}_{\alpha, \Gamma}$ is the mean of the posterior distribution of Z given $\underline{Z}$, after averaging of over a

$$\begin{aligned}\hat{z}_{\alpha, \Gamma} &= E\{E(Z|\underline{z}, a) | \underline{z}\} \\ &= f'E(a|\underline{z}) + k'K^{-1}\{\underline{z} - F'E(a|\underline{z})\} \\ &= k'K^{-1}\underline{z} - \emptyset'E(a|\underline{z})\end{aligned}$$

This establishes formula 2.6. The proof of formula 2.7 is omitted because the calculations are straightforward and tedious, using the fact that $E(a|\underline{z}) = (G+\Gamma^{-1})^{-1}\Gamma^{-1}\alpha + (G+\Gamma^{-1})^{-1}FK^{-1}\underline{z}$.

Theorem 2.1: The best linear unbiased predictor $\hat{z}$ of Z is extended Bayes and minimax.

Proof: The risk of $\hat{z}$ as a function of a is $R(a, \hat{z}) = \alpha^2 - k'K^{-1}k + \emptyset'G^{-1}\emptyset$. This expression does not depend on a , therefore $\hat{z}$ is an equalizer. Secondly, when all the eigen values of Γ tend to $+\infty$, the risk of $\hat{z}_{\alpha, \Gamma}$, given by 2.7, tends to $R(a, \hat{z})$. Therefore $\hat{z}$ is extended Bayes, and since it is an equalizer it must be minimax. ■

The next two problems are to show that $\hat{z}$ is admissible and UMVUE. So now we abandon the assumption that a has a prior distribution and return to the case when it is a free unspecified vector-parameter.

Lemma 2.2: For any predictor Z^* of Z based on $\underline{z}$ we have

$$E(Z^* - Z)^2 = \sigma^2 - k'K^{-1}k + E\{Z^* - E(Z|\underline{z})\}^2 \quad 2.8$$

Proof: $Z - Z^*$ can be written as the sum of $Z - E(Z|\underline{z})$ and $E(Z|\underline{z}) - Z^*$. Next, $Z - E(Z|\underline{z})$ is uncorrelated with $\underline{z}$;

hence, by normality it is independent of $\underline{z}$, and independent also of $E(\underline{z}|\underline{z}) - \underline{z}^*$ because this is a function of $\underline{z}$.

Formula 2.8 follows. ■

Lemma 2.3: $\hat{a} = G^{-1}FK^{-1}\underline{z}$ is a sufficient estimator for a , among the estimators based on $\underline{z}$.

This is a standard result in generalized regression.

We may just mention for completeness that $E(\underline{z}|\hat{a}) = F'\hat{a}$ and $\text{Cov}(\underline{z}|\hat{a}) = K - F'G^{-1}F$. Hence $\underline{z}$ given $\hat{a}$, which is normally distributed, has a distribution which does not depend on a .

Theorem 2.2: $\hat{z}$ is an admissible predictor of z .

Proof: Suppose that $\hat{z}$ is not admissible and that $z^* = g(\underline{z})$ is such that

$$E(z^* - z)^2 \leq E(\hat{z} - z)^2 \quad 2.9$$

for all a with strict inequality for at least one a .

Then from lemma 2.2 we deduce that

$$E\{z^* - E(z|\underline{z})\}^2 \leq E\{\hat{z} - E(z|\underline{z})\}^2 \quad 2.10$$

for all a , with strict inequality for at least one a .

$$E(z|\underline{z}) = k'K^{-1}\underline{z} - \emptyset'a, \text{ and} \\ \hat{z} = k'K^{-1}\underline{z} - \emptyset'\hat{a}$$

Therefore, pursuing our argument, we have.

$$E\{k'K^{-1}\underline{z}-g(\underline{z})-\phi'a\}^2 \leq E\{\phi'\hat{a}-\phi'a\}^2 \quad 2.11$$

for all a , with strict inequality for at least one a .

Then by lemma 2.3 and the Rao-Blackwell theorem, the quantity $E\{k'K^{-1}\underline{z}-g(\underline{z})|\hat{a}\}$ which is a function of $\hat{a}$ and does not depend on a , is a better estimator of $\phi'a$ than $\phi'\hat{a}$. Or, in other words, $\phi'\hat{a}$ is not an admissible estimator of $\phi'a$.

But this is a contradiction because $\hat{a} \sim N(a, G^{-1})$ and therefore it is elementary to see that $\phi'\hat{a}$ is admissible for $\phi'a$ among the class of estimators function of $\hat{a}$. (See also Cohen (1965).)

Theorem 2.3: $\hat{Z}$ is the uniformly minimum variance unbiased predictor of Z .

Proof: From lemma 2.2 we see that it is sufficient to show that the term $E\{E(Z|\underline{Z})-\hat{Z}\}^2$ is uniformly minimum in a , among unbiased predictors of Z .

An unbiased predictor $g(\underline{Z})$ of Z must satisfy $Eg(\underline{Z}) = EZ = E\{E(Z|\underline{Z})\}$; then $E\{f'a+k'K^{-1}[\underline{Z}-F'a]-g(\underline{Z})\} = 0$. Therefore it is such that $k'K^{-1}\underline{Z}-g(\underline{Z})$ is a unbiased estimator of $\phi'a$.

By the generalized version of Cramer-Rao inequality any

unbiased estimator of $\phi'a$ has its variance bounded from below by $\phi'I(a)^{-1}\phi$ where $I(a)$ is the Fisher information matrix of $\underline{Z}$ on a . It turns out that $I(a) = G$. Hence the Cramer-Rao lower bound is $\phi'G^{-1}\phi$. This is attained by $\phi'\hat{a}$, because $E\phi'\hat{a} = \phi'a$, and $E\{\phi'\hat{a} - \phi'a\}^2 = \phi'\text{Cov}(\hat{a})\phi = \phi'G^{-1}\phi$.

In conclusion for any unbiased predictor $g(\underline{Z})$ of Z we have $E\{E(Z|\underline{Z}) - g(\underline{Z})\}^2 = E\{k'K^{-1}\underline{Z} - g(\underline{Z}) - \phi'a\}^2 \geq \phi'G^{-1}\phi = E(\phi'\hat{a} - \phi'a)^2 = E\{E(Z|\underline{Z}) - \hat{Z}\}^2$ for all a .

3. A Stein like predictor for predicting simultaneously several random variables:

Suppose now that we want to predict N random variables $Z_{n+1}, Z_{n+2}, \dots, Z_{n+N}$ from the observations of $Z_1, Z_2, \dots, Z_n$.

The assumptions that we make here are

$$\underline{Z} \sim N(F'a, K)$$

$$Z_{n+i} \sim N(f_i'a, \sigma_i^2) \quad i = 1, 2, \dots, N$$

$$\text{Cov}(\underline{Z}, Z_{n+i}) = k_i \quad i = 1, 2, \dots, N$$

and all the Z_i 's (the observations and the values to be predicted) are jointly normal. As before all the parameters appearing in the distributions are specified, except for the free vector-parameter a .

The object of this section is to exhibit a predictor $\tilde{Z}$ of the type introduced by James and Stein (1961) to estimate the mean of a multivariate normal random vector, such that

$$\sum_{i=1}^N E(\tilde{Z}_{n+i} - Z_{n+i})^2 < \sum_{i=1}^N E(\hat{Z}_{n+i} - Z_{n+i})^2$$

for all values of a .

In Section 2 we saw that the admissibility of $\hat{Z}$ ultimately reduced to the admissibility of $\phi'\hat{a}$ to estimate $\phi'a$ when $\hat{a} \sim N(a, G^{-1})$. So it is natural that in the present section

the key result will be the inadmissibility of $\hat{\phi}a$ to estimate ϕa where ϕ is a $N \times p$ matrix satisfying certain conditions. This is stated in a general form in the following theorem.

Theorem 3.1: Consider $X \sim N(\theta, I_p)$, a p -dimensional normal random vector; let C be an $N \times p$ matrix then $\exists \eta > 0$ such that for all $\theta \in R^p$

$$E \left\| CX \left(1 - \frac{\eta}{\|X\|^2} \right) - C\theta \right\|^2 < E \left\| CX - C\theta \right\|^2 \quad 3.1$$

if and only if

$$\text{max eigenvalue of } CC' < \frac{1}{2} \text{tr } CC' \quad 3.2$$

Remark: Condition (2) implies $\text{rank } C \geq 2$. In particular if C is the identity, condition (2) is satisfied if and only if $p \geq 3$, which is a classical fact in Stein-estimation (see James and Stein (1961)).

Theorem 3.1 will be proved after the following two lemmas.

Lemma 3.1: If $X \sim N(\theta, I_p)$, for any row vector C and any function g such that the following expectations exist, we have

$$E(CXg(\|X\|^2)) = C\theta E g(X_{p+2K+2}^2) \quad 3.3$$

and also

$$E\{XX'g(\|X\|^2)\} = I_p E g(X_{p+2K+2}^2) + \theta\theta' E g(X_{p+2K+4}^2) \quad 3.4$$

where $K \sim \text{Poisson}\left(\frac{\|\theta\|^2}{2}\right)$.

For a proof the reader is referred to Stein (1966) where the essential technique is shown in order to prove formula 3.3; the same technique can be used to prove formula 3.4.

Lemma 3.2: Under the condition of theorem 3.1, we have

$$E\left\|CX\left(1 - \frac{\eta}{\|X\|^2}\right) - C\theta\right\|^2 = \text{tr } CC' \left[1 - 2\eta E \frac{1}{p+2K} + \eta^2 E \frac{1}{(p+2K)(p-2+2K)}\right] + \frac{\|C\theta\|^2}{\|\theta\|^2} (4\eta + \eta^2) E \frac{2K}{(p+2K)(p-2+2K)} \quad 3.5$$

where $K \sim \text{Poisson}\left(\frac{\|\theta\|^2}{2}\right)$.

Proof of Lemma 3.2: We start from

$$\begin{aligned} E\left\|CX\left(1 - \frac{\eta}{\|X\|^2}\right) - C\theta\right\|^2 &= E\left\|C(X-\theta) - \eta \frac{CX}{\|X\|^2}\right\|^2 \\ &= E(X-\theta)' C' C (X-\theta) - 2 E(X-\theta)' C' C \frac{X\eta}{\|X\|^2} + \eta^2 E \frac{X'C'CX}{\|X\|^4}. \end{aligned} \quad 3.6$$

Since for two vectors in $\mathbb{R}^p$ $x'y = \text{tr } yx'$, expression (3.6) can be rewritten

$$\text{tr } C E(X-\theta)(X-\theta)' C' - 2\eta \text{tr } C E \frac{X(X-\theta)'}{\|X\|^2} C' + \eta^2 \text{tr } C E \frac{XX'}{\|X\|^4} C' \quad 3.7$$

By Lemma 3.1 and using $E \frac{1}{X_n^2} = \frac{1}{n-2}$ and $E \frac{1}{X_n^2}^2 = \frac{1}{(n-2)(n-4)}$,

expression 3.7 is equal to

$$\begin{aligned} & \text{tr } CC' - 2\eta \text{tr } C \left\{ I_p E \frac{1}{X_{p+2K+2}^2} + \theta\theta' E \frac{1}{X_{p+2K+4}^2} - \theta\theta' E \frac{1}{X_{p+2K+2}^2} \right\} C' + \\ & \quad \eta^2 \text{tr } C \left\{ I_p E \left[\frac{1}{X_{p+2K+2}^2} \right]^2 + \theta\theta' E \left[\frac{1}{X_{p+2K+4}^2} \right]^2 \right\} C' \\ & = \text{tr } CC' - 2\eta \text{tr } CC' E \frac{1}{p+2K} - 2\eta \|C\theta\|^2 E \left\{ \frac{1}{p+2K+2} - \frac{1}{p+2K} \right\} + \\ & \quad \eta^2 \text{tr } CC' E \frac{1}{(p+2K)(p+2K-2)} + \eta^2 \|C\theta\|^2 E \frac{1}{(p+2K+2)(p+2K)} \\ & = \text{tr } CC' \left[1 - 2\eta E \frac{1}{p+2K} + \eta^2 E \frac{1}{(p+2K)(p+2K-2)} \right] + \\ & \quad \|C\theta\|^2 (4\eta + \eta^2) E \frac{1}{(p+2K+2)(p+2K)}. \end{aligned}$$

Finally, to get expression 3.5 given in the statement of

Lemma 3.2 we have to show

$$E \frac{1}{(p+2K+2)(p+2K)} = \frac{1}{\|\theta\|^2} E \frac{2K}{(p+2K)(p+2K-2)} .$$

This is done as follows:

$$\begin{aligned} E \frac{1}{(p+2K+2)(p+2K)} &= \sum_{k=0}^{\infty} e^{-\|\theta\|^2/2} \frac{\left(\|\theta\|^2/2\right)^k}{k!} \frac{1}{(p+2k+2)(p+2k)} \\ &= \frac{2}{\|\theta\|^2} \sum_{k=0}^{\infty} e^{-\|\theta\|^2/2} \frac{\left(\|\theta\|^2/2\right)^{k+1}}{(k+1)!} \frac{k+1}{(p+2(k+1))(p+2(k+1)-2)} \\ &= \frac{2}{\|\theta\|^2} \sum_{k=1}^{\infty} e^{-\|\theta\|^2/2} \frac{\left(\|\theta\|^2/2\right)^k}{k!} \frac{k}{(p+2k)(p+2k-2)} \\ &= \frac{1}{\|\theta\|^2} E \frac{2K}{(p+2K)(p+2K-2)} , \end{aligned}$$

which completes the proof of Lemma 3.2. Now we can turn to the proof of Theorem 3.1.

Proof of Theorem 3.1: The key to the proof is to observe that

$$\max_{\theta} \frac{\|C\theta\|^2}{\|\theta\|^2} = \text{max eigenvalue of } CC' .$$

First if

$$\text{max eigenvalue of } CC' \leq \left(\frac{1-\epsilon}{2}\right) \text{tr } CC', \quad 3.8$$

then $E\|CX\left(1 - \frac{\eta}{\|X\|^2}\right) - C\theta\|^2$ is bounded by

$$\begin{aligned} \text{tr } CC' \left\{ 1 - 2\eta E \frac{1}{p+2K} + \eta^2 E \frac{1}{(p+2K)(p-2+2K)} + \right. \\ \left. 2(1-\epsilon)\eta E \frac{2K}{(p+2K)(p-2+2K)} + (1-\epsilon)\eta^2 E \frac{K}{(p+2K)(p-2+2K)} \right\}. \quad 3.9 \end{aligned}$$

In turn to bound expression 3.9 we need $p \geq 3$; but, as observed in the remark on page 12, this is implied by 3.8. Then expression 3.9 is bounded by

$$\text{tr } CC' \left\{ 1 - 2\epsilon\eta E \frac{1}{p+2K} + \eta^2 E \frac{1}{p+2K} \right\}.$$

Therefore if we pick $\eta = \epsilon$, we get

$$E\|CX\left(1 - \frac{\epsilon}{\|X\|^2}\right) - C\theta\|^2 \leq \text{tr } CC' \left(1 - \epsilon^2 E \frac{1}{p+2K}\right) < \text{tr } CC' = E\|CX - C\theta\|^2.$$

Conversely if max eigenvalue of $CC' \geq \frac{1}{2} \text{tr } CC'$ we can choose the direction of θ such that

$$\frac{\|C\theta\|^2}{\|\theta\|^2} = \text{max eigenvalue of } CC'$$

and therefore such that

$$\begin{aligned} E\|CX\left(1 - \frac{\eta}{\|X\|^2}\right) - C\theta\|^2 &\geq \text{tr } CC' \left[1 - 2\eta E \frac{1}{p+2K} + \right. \\ &\quad \left. 2\eta E \frac{2K}{(p+2K)(p-2+2K)} + \eta^2 E \frac{K+1}{(p+2K)(p-2+2K)}\right] \\ &= \text{tr } CC' \left[1 - 2\eta E \frac{p-2}{(p+2K)(p-2+2K)} + \eta^2 E \frac{K+1}{(p+2K)(p-2+2K)}\right]. \end{aligned}$$

Now, for any $\eta > 0$, there exists θ large enough such that the above quantity is strictly greater than $\text{tr } CC'$, because when $\|\theta\| \rightarrow +\infty$

$$E \frac{p-2}{(p+2K)(p-2+2K)} = o\left(E \frac{K+1}{(p+2K)(p-2+2K)}\right).$$

This completes the proof of Theorem 3.1.

Application of Theorem 3.1 to the prediction of z_{n+1} , $z_{n+2}, \dots, z_{n+N}$ using $z_1, z_2, \dots, z_n$:

Parallel to the notation $\emptyset = FK^{-1}k-f$, let us define $\emptyset_i = FK^{-1}k_i - f_i$ $i=1, 2, \dots, N$.

From the results of Section 2, we know that the individual best linear unbiased predictors of each z_{n+i} , $i=1, 2, \dots, N$, based on $\underline{z}$, are

$$\begin{aligned}\hat{z}_{n+i} &= f_i' \hat{a} + k_i' K^{-1} [z - F' \hat{a}] \\ &= k_i' K^{-1} \underline{z} - \phi_i' \hat{a} \quad i=1,2,\dots,N\end{aligned}$$

And we know that they are individually admissible. The fact that it is not the case, in general, when they are used simultaneously, is the object of the following theorem:

Theorem 3.2: For $i=1,2,\dots,N$, define

$$\tilde{z}_{n+i} = k_i' K^{-1} \underline{z} - \phi_i' \hat{a} \left(1 - \frac{\eta}{\hat{a}' G \hat{a}}\right) \quad 3.10$$

then $\exists \eta > 0$ such that, for all a

$$\sum_{i=1}^N E(\tilde{z}_{n+i} - z_{n+i})^2 < \sum_{i=1}^N E(\hat{z}_{n+i} - z_{n+i})^2 \quad 3.11$$

if and only if

$$\text{max eigenvalue of } [\phi_i' G^{-1} \phi_j] < \sum_{i=1}^N \phi_i' G^{-1} \phi_i \quad 3.12$$

where $[\phi_i' G^{-1} \phi_j]$ is the $N \times N$ matrix whose (i,j) element is $\phi_i' G^{-1} \phi_j$.

Proof: Since $\underline{z} \sim N(F'a, K)$ we have $K^{-\frac{1}{2}} \underline{z} \sim N(K^{-\frac{1}{2}} F'a, I_p)$.

Consider an orthogonal matrix $H = \begin{pmatrix} H_1 \\ H_2 \end{pmatrix}$ such that

$v(H_1') = v(K^{-\frac{1}{2}} F')$ where $v(H_1')$ stands for the vector space generated by the columns of H_1' . Then

$$HK^{-\frac{1}{2}} \underline{Z} \sim N \left(\begin{pmatrix} H_1 K^{-\frac{1}{2}} F' a \\ 0 \end{pmatrix}, I_p \right).$$

let $M = H_1 K^{-\frac{1}{2}} F'$; it is of size $p \times p$ and non singular since we assumed that $FK^{-1}F'$ is of full rank. Let $\alpha = Ma$ and $\hat{\alpha} = M\hat{a}$, then $\hat{\alpha} \sim N(\hat{\alpha}, MG^{-1}M')$. Let us show that $MG^{-1}M'$ is actually I_p : we start from

$$\begin{aligned} M'M &= FK^{-\frac{1}{2}} H_1' H_1 K^{-\frac{1}{2}} F' \\ &= FK^{-\frac{1}{2}} (I - H_2' H_2) K^{-\frac{1}{2}} F' \\ &= FK^{-\frac{1}{2}} F' - FK^{-\frac{1}{2}} H_2' H_2 K^{-\frac{1}{2}} F' \end{aligned}$$

But $H_2 K^{-\frac{1}{2}} F' = 0$; therefore $M'M = G$. Then $G^{-\frac{1}{2}} M'M G^{-\frac{1}{2}} = I_p$. This means that $MG^{-\frac{1}{2}}$ is orthogonal, hence $MG^{-1}M' = I_p$. In conclusion, we have $\hat{\alpha} \sim N(\alpha, I_p)$.

From lemma 2.2 we can write

$$\begin{aligned} \sum_{i=1}^N E(\tilde{Z}_{n+i} - Z_{n+i})^2 &= \sum_{i=1}^N E(Z_{n+i} - E(Z_{n+i} | \underline{Z}))^2 + \\ &\quad \sum_{i=1}^N E(\tilde{Z}_{n+i} - E(Z_{n+i} | \underline{Z}))^2 \end{aligned} \quad 3.13$$

the first term of the right hand side of 3.13 is

$$\sum_{i=1}^N (\sigma_i^2 - k_i' K^{-1} k_i). \quad \text{On the other hand, since}$$

$E(Z_{n+i} | \underline{Z}) = k_i' K^{-1} \underline{Z} - \phi_i' a$ the second term of the right hand side of 3.13 can be rewritten

$$\sum_{i=1}^N E(\phi_i' \hat{a} \left(1 - \frac{\eta}{\hat{a}' G \hat{a}}\right) - \phi_i' a)^2 \quad 3.14$$

Note that $\|\hat{\alpha}\|^2 = \hat{\alpha}' \hat{\alpha} = \hat{\alpha}' M' M \hat{\alpha} = \hat{\alpha}' G \hat{\alpha}$; and let C be the $N \times p$ matrix whose i th row is $\phi_i' M$. Then expression 3.14 becomes

$$\sum_{i=1}^N E(\phi_i' M \hat{\alpha} \left(1 - \frac{\eta}{\|\hat{\alpha}\|^2}\right) - \phi_i' M \alpha)^2 = E \|C \hat{\alpha} \left(1 - \frac{\eta}{\|\hat{\alpha}\|^2}\right) - C \alpha\|^2.$$

Since $\hat{\alpha} \sim N(\alpha, I_p)$, by Theorem 3.1 we know that $\exists \eta > 0$ such that for all α

$$E \|C \hat{\alpha} \left(1 - \frac{\eta}{\|\hat{\alpha}\|^2}\right) - C \alpha\|^2 < E \|C \hat{\alpha} - C \alpha\|^2$$

if and only if maxeigenvalue of $CC' < \frac{1}{2} \text{tr} CC'$.

To finish the proof let us observe on the one hand that $CC' = [\phi_i' G^{-1} \phi_j]$, and on the other hand (by going backward in the computation) that

$$E \|C \hat{\alpha} - C \alpha\|^2 = \sum_{i=1}^N E \{E(Z_{n+i} | \underline{Z}) - \hat{Z}_{n+i}\}^2$$

this, after another application of lemma 2.2, completes the proof of Theorem 3.2.

4. A non linear predictor of a single random variable z , based on $z_1, z_2, \dots, z_n$.

Coming back to the problem of predicting a single z from the observations $\underline{z}$, we show in this section another way to reduce the extra-term $\phi'G^{-1}\phi$, in the mean squared error of the estimator $\hat{z}$, that we have already discussed in Section 2.

The idea is to try to dispense with unbiasedness. This leads to a predictor $\hat{z}_a$, which depends on the unspecified vector parameter a . It is therefore of no use itself but the predictor denoted symbolically $\hat{z}_{\hat{a}}$, obtained by substituting $\hat{a}$ for a in $\hat{z}_a$, turns out to be interesting; we will denote it z^* .

Preliminary calculations show that the predictor $\hat{z}_a = \lambda_1 z_1 + \dots + \lambda_n z_n$ which minimizes $E(\hat{z}_a - z)^2$, with no unbiasedness restriction with respect to a , is

$$\hat{z}_a = k'K^{-1}\underline{z} - \frac{\phi'a}{1+a'Ga} a'FK^{-1}\underline{z} \quad 4.1$$

and its risk is

$$E(\hat{z}_a - z)^2 = \sigma^2 - k'K^{-1}k + \frac{(\phi'a)^2}{1+a'Ga} \quad 4.2$$

Note that $\sup_a \frac{(\phi'a)^2}{1+a'Ga} = \phi'G^{-1}\phi$ which is the third term in

the risk of $\hat{z}$.

So the predictor we want to study in this section is

$$z^* = k'K^{-1}\underline{z} - \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} \cdot \hat{a}'FK^{-1}\underline{z} \quad 4.3$$

as derived from formula 4.1 by substitution of $\hat{a}$ for a .

In the sequel we assume that z and $\underline{z}$ are jointly normally distributed.

Theorem 4.1: The predictor z^* can be reexpressed as

$$z^* = \hat{z} + \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} \quad 4.4$$

and its risk is

$$E(z^* - z)^2 = \sigma^2 - k'K^{-1}k + \text{Var}\left(\frac{\phi'\hat{a} \hat{a}'G\hat{a}}{1+\hat{a}'G\hat{a}}\right) + \left\{E\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right)\right\}^2 \quad 4.5$$

To prove Theorem 4.1 we need two lemmas.

Lemma 4.1: Consider $X \sim N(\mu, \Sigma)$ of size $n \times 1$; suppose Σ is positive definite. Let M be a $p \times n$ matrix ($p \leq n$) of full rank, and v a $p \times 1$ vector. Then

$$E(X|MX=v) = \mu - \Sigma M'(M\Sigma M')^{-1}M\mu + \Sigma M'(M\Sigma M')^{-1}v$$

$$\text{Cov}(X|MX=v) = \Sigma(I - M'(M\Sigma M')^{-1}M\Sigma)$$

For a proof of Lemma 4.1 the reader is referred to Cabannes (1979).

Lemma 4.2: $E(\underline{z}|\hat{a}) = F'\hat{a}$ and $E(\underline{z}|\hat{a}) = -\phi'a + k'K^{-1}F'\hat{a}$

Lemma 4.2 can be proved as an application of Lemma 4.1. Again we skip the calculations.

Proof of Theorem 4.1: To prove 4.4 we write

$$\begin{aligned} z^* - \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} &= k'K^{-1}\underline{z} - \frac{\phi'\hat{a}\hat{a}'FK^{-1}\underline{z}}{1+\hat{a}'G\hat{a}} - \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} \\ &= k'K^{-1}\underline{z} - \frac{\phi'\hat{a}\hat{a}'G\hat{a}}{1+\hat{a}'G\hat{a}} - \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} = k'K^{-1}\underline{z} - \phi'\hat{a} = \hat{z} \\ \text{Hence } z^* &= \hat{z} + \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} \end{aligned}$$

Next the risk of z^* is

$$\begin{aligned} E(z^* - z)^2 &= E\left(\hat{z} - z + \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right)^2 \\ &= E(\hat{z} - z)^2 + E\left[\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right)^2\right] + \\ &\quad 2E\left[(\hat{z} - z)\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right)\right] \end{aligned} \quad 4.6$$

To compute the cross-product term in 4.6 we condition on $\hat{a}$:

$$E\left[(\hat{z} - z)\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right) \mid \hat{a}\right] = \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}} E(\hat{z} - z \mid \hat{a})$$

and from Lemma 4.2 we have

$$\begin{aligned} E(\hat{Z}-Z|\hat{a}) &= \lambda'E(\underline{Z}|\hat{a}) - E(Z|\hat{a}) \\ &= (k'K^{-1}-\phi'G^{-1}FK^{-1})F'\hat{a} - (k'K^{-1}F'a - \phi'\hat{a}) \\ &= \phi'(a-\hat{a}) \end{aligned}$$

Therefore

$$\begin{aligned} E\left[(\hat{Z}-Z)\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right)\right] &= E\left[\phi'(a-\hat{a}) \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right] \\ &= \phi'a E\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right) - E\left[\frac{(\phi'\hat{a})^2}{1+\hat{a}'G\hat{a}}\right] \\ &= -\text{Cov}\left(\phi'\hat{a}, \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right) \end{aligned}$$

Then

$$\begin{aligned} E(Z^*-Z)^2 &= \sigma^2 - k'K^{-1}k + \phi'G^{-1}\phi + E\left[\left(\frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right)^2\right] - \\ &\quad 2\text{Cov}\left(\phi'\hat{a}, \frac{\phi'\hat{a}}{1+\hat{a}'G\hat{a}}\right) \end{aligned}$$

Formula 4.5 follows after some more manipulation, noting that $\phi'G^{-1}\phi = \text{Var}(\phi'\hat{a})$. ■

The next problem is to compare the risks $R(a, Z^*) = E(Z^*-Z)^2$ and $R(a, \hat{Z}) = E(\hat{Z}-Z)^2$. Let $R(\hat{Z}) \triangleq R(a, \hat{Z})$ because it does not depend on a . Since we showed in Section 2 the admissibility of $\hat{Z}$, Z^* cannot be uniformly better than $\hat{Z}$ when a varies.

However, we have the following result

Theorem 4.2: $R(0, Z^*) < R(\hat{Z})$ and $\lim_{\|a\| \rightarrow +\infty} R(a, Z^*) = R(\hat{Z})$

Proof: When $a = 0$, we have $E\left(\frac{\phi' \hat{a}}{1 + \hat{a}' G \hat{a}}\right) = 0$ because the distribution of $\hat{a}$ then is $N(0, G^{-1})$ and $\frac{\phi' \hat{a}}{1 + \hat{a}' G \hat{a}}$ has a distribution symmetric about zero. Therefore $R(0, Z^*) = \sigma^2 - k' K^{-1} k + \text{Var}\left(\phi' \hat{a} \frac{\hat{a}' G \hat{a}}{1 + \hat{a}' G \hat{a}}\right)$ and again since $\hat{a} \sim N(0, G^{-1})$ we have $\text{Var}\left(\phi' \hat{a} \cdot \frac{\hat{a}' G \hat{a}}{1 + \hat{a}' G \hat{a}}\right) < \text{Var}(\phi' \hat{a}) = \phi' G^{-1} \phi$ hence $R(0, Z^*) < R(\hat{Z})$.

Next note that when $\|a\| \rightarrow +\infty$ the term $\frac{\phi' \hat{a}}{1 + \hat{a}' G \hat{a}}$ tends to zero in probability, and therefore also in distribution. And since there exists a constant c such that $\forall b \in \mathbb{R}^p$, $\left| \frac{\phi' b}{1 + b' G b} \right| < c$ we deduce by a classical convergence theorem that all the moments of $\frac{\phi' \hat{a}}{1 + \hat{a}' G \hat{a}}$ tend to zero. Then because $\text{Var} \phi' \hat{a} = \phi' G^{-1} \phi$ we have

$$\lim_{\|a\| \rightarrow +\infty} \text{Var}\left(\phi' \hat{a} - \frac{\phi' \hat{a}}{1 + \hat{a}' G \hat{a}}\right) + E\left[\left(\frac{\phi' \hat{a}}{1 + \hat{a}' G \hat{a}}\right)^2\right] = \phi' G^{-1} \phi$$

This completes the proof of Theorem 4.2. ■

We state briefly some final results without proving them, referring the interested reader to the report of the author (1979).

The risk of Z^* can be reexpressed as follows

$$R(a, z^*) = R(\hat{z}) + \xi(a) \quad \text{where}$$

$$\xi(a) = \frac{1}{2} \phi' G^{-1} \phi \quad E \left[\frac{1}{1 + \chi_{p+2K}^2} - E \frac{1}{1 + \chi_{p+2K+2}^2} \right] +$$

$$\frac{5}{2} (\phi' a)^2 \left[E \frac{1}{1 + \chi_{p+2K+2}^2} - E \frac{1}{1 + \chi_{p+2K+4}^2} \right]$$

and $K \sim \text{Poisson} \left(\frac{a' G a}{2} \right)$. And this, combined with the following approximation (Stein 1966)

$$E \frac{1}{\chi_{p+2K}^2} \approx \frac{1}{p-2+t} \left\{ 1 - \frac{2t}{(p-2+t)^2} \right\}$$

where $t = a' G a$, can be used to compute approximate values of the risk function of z^* .

Illustration of the preceding method in the estimation of the mean of i.i.d. normal random variables with known variance:

Suppose that $Z, Z_1, Z_2, \dots, Z_n$ are i.i.d. $N(m, \sigma^2)$ then of course the predictors $\hat{z}$ and z^* truly estimate the mean m , so let's denote them $\hat{m}$ and m^* . Note that

$$E(\hat{m} - m)^2 = E(\hat{m} - Z)^2 - \sigma^2$$

$$\text{and } E(m^* - m)^2 = E(m^* - Z)^2 - \sigma^2$$

$$\text{let } R(m, \hat{m}) = E(\hat{m} - m)^2 \quad \text{and}$$

$$R(m, m^*) = E(m^* - m)^2$$

then if we define $\xi_0(m)$ by

$$\xi_0(m) = E \left[\left(\frac{X}{1+X^2} \right)^2 \right] - 2 \operatorname{cov} \left(X, \frac{X}{1+X^2} \right)$$

where $X \sim N(m,1)$, it is easy to see that

$$R(m, m^*) = R(m, \hat{m}) + \frac{\sigma^2}{n} \xi_0 \left(\frac{m\sqrt{n}}{\sigma} \right)$$

In the case $n = 3$ the figure below representing $R(m, m^*)$ and $R(m, \hat{m})$ shows that if we know a priori that $m \in [m_0 - .8\sigma, m_0 + .8\sigma]$ for some known constant m_0 we should subtract the constant m_0 from all our observations and use m^* instead of $\hat{m}$.

(see the figure next page)




Figure: Comparison of the risk functions of $\hat{m}$ and m^* .

REFERENCES:

- Cabannes, A. (1979): Estimation of Random Fields from from Network Observations. Stanford University, Statistics Dept., SIMS Technical Report #26.
- Cohen, A. (1965): Estimates of linear combinations of the parameters in the mean vector of a multivariate distribution. Annals of Mathematical Statistics 36, 78-87.
- Goldberger, A.S. (1962): Best linear unbiased prediction in the generalized linear regression model. J.A.S.A. 57, 369-375.
- James, W. and Stein, C. (1960): Estimation with quadratic loss. Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, 361-379.
- Stein, C. (1966): An approach to the recovery of inter-block information in balanced incomplete designs. In Research Papers in Statistics (Festschrift for J. Neyman) F.N. David, ed., John Wiley.

OFFICE OF NAVAL RESEARCH
STATISTICS AND PROBABILITY PROGRAM

BASIC DISTRIBUTION LIST
FOR
UNCLASSIFIED TECHNICAL REPORTS

JANUARY 1979

	Copies		Copies
Statistics and Probability Program (Code 436) Office of Naval Research Arlington, VA 22217	3	Office of Naval Research San Francisco Area Office One Hallidie Plaza - Suite 601 San Francisco, CA 94102	1
Defense Documentation Center Cameron Station Alexandria, VA 22314	12	Office of Naval Research Scientific Liaison Group Attn: Scientific Director American Embassy - Tokyo APO San Francisco 96503	1
Office of Naval Research New York Area Office 715 Broadway - 5th Floor New York, New York 10003	1	Applied Mathematics Laboratory David Taylor Naval Ship Research and Development Center Attn: Mr. G. H. Gleissner Bethesda, Maryland 20084	1
Commanding Officer Office of Naval Research Branch Office Attn: D. A.L. Powell Building 114, Section D 666 Summer Street Boston, MA 02210	1	Commandant of the Marine Corps (Code AX) Attn: Dr. A.L. Slafkosky Scientific Advisor Washington, DC 20380	1
Commanding Officer Office of Naval Research Branch Office Attn: Director for Science 536 South Clark Street Chicago, Illinois 60605	1	Director National Security Agency Attn: Mr. Stahly and Dr. Maar (R51) Fort Meade, MD 20755	2
Commanding Officer Office of Naval Research Branch Office Attn: Dr. Richard Lau 1030 East Green Street Pasadena, CA 91101	1	Navy Library National Space Technology Laboratory Attn: Navy Librarian Bay St. Louis, MS 39522	1

Copies

U.S. Army Research Office
P.O. Box 12211
Attn: Dr. J. Chandra
Research Triangle Park, NC 27706

1

Naval Sea Systems Command
(NSEA 03F)
Attn: Miss B. S. Orleans
Crystal Plaza #6
Arlington, VA 20360

1

Office of the Director
Bureau of The Census
Attn: Mr. H. Nisselson
Federal Building 3
Washington, DC 20233

1

OASD (I&L), Pentagon
Attn: Mr. Charles S. Smith
Washington, DC 20301

1

ARI Field Unit-USAREUR
Attn: Library
c/o ODCSPER
HQ USAREUR & 7th Army
APO New York 09403

1

Naval Underwater Systems Center
Attn: Dr. Derrill J. Bordelon
Code 21
Newport, Rhode Island 02840

1

Library, Code 1424
Naval Postgraduate School
Monterey, California 93940

1

Technical Information Division
Naval Research Laboratory
Washington, DC 20375

1

OFFICE OF NAVAL RESEARCH
STATISTICS AND PROBABILITY PROGRAM

MODELING AND ESTIMATION DISTRIBUTION LIST
FOR
UNCLASSIFIED TECHNICAL REPORTS

JANUARY 1979

	Copies		Copies
Technical Library Naval Ordnance Station Indian Head, Maryland 20640	1	Professor H. Robbins Department of Mathematics Columbia University New York, New York 10027	1
Bureau of Naval Personnel Department of the Navy Technical Library Washington, DC 20370	1	Professor W. M. Hirsch Courant Institute of Mathematical Sciences New York University New York, New York 10453	1
Library Naval Ocean Systems Center San Diego, CA 92152	1	Professor F. J. Anscombe Department of Statistics Yale University New Haven, CT 06520	1
Professor Robert Serfling Department of Statistics Florida State University Tallahassee, FL 32306	1	Professor S. S. Gupta Department of Statistics Purdue University Lafayette, Indiana 47907	1
Professor Ralph A. Bradley Department of Statistics Florida State University Tallahassee, FL 32306	1	Professor R. E. Bechhofer Department of Operations Research Cornell University Ithaca, New York 14850	1
Professor G. S. Watson Department of Statistics Princeton University Princeton, NJ 08540	1	Professor D. B. Owen Department of Statistics Southern Methodist University Dallas, Texas 75275	1
Professor P. J. Bickel Department of Statistics University of California Berkeley, CA 94720	1	Professor A.F. Veinott Department of Operations Research Stanford University Stanford, CA 94305	1

	Copies		Copies
Professor D. L. Iglehart Department of Operations Research Stanford University Stanford, CA 94305	1	Professor D. P. Gaver Department of Operations Research Naval Postgraduate School Monterey, CA 93940	1
Professor Herbert Solomon Department of Statistics Stanford University Stanford, CA 94305	1	Dr. M. J. Fischer Defense Communications Agency Defense Communications Engineering Center 1860 Wiehle Avenue Reston, Virginia 22090	1
Professor P.A.W. Lewis Department of Operations Research Naval Postgraduate School Monterey, CA 93940	1	Defense Logistics Studies Information Exchange Army Logistics Management Center Attn: Mr. J. Dowling Fort Lee, Virginia 23801	1
Professor R. L. Disney Dept. of Industrial Engineering and Operations Research Virginia Polytechnic Institute and State University Blacksburg, VA 24061	1	Professor D. O. Siegmund Department of Statistics Stanford University Stanford, CA 94305	1
Professor H. Chernoff Department of Mathematics Massachusetts Institute of Technology Cambridge, MA 02139	1	Professor M. L. Puri Department of Mathematics Indiana University Foundation P.O. Box F Bloomington, Indiana 47401	1
Dr. D. E. Smith Desmatics, Inc. P.O. Box 618 State College, PA 16801	1	Professor S. M. Ross College of Engineering University of California Berkeley, CA 94720	1
Professor J.A. Muckstadt Department of Operations Research Cornell University Ithaca, New York 14850	1	Professor J. E. Boyer, Jr. Department of Statistics Southern Methodist University Dallas, Texas 75275	1
Professor R. W. Madsen Department of Statistics University of Missouri Columbia, Missouri 65201	1	Professor Grace Wahba Department of Statistics University of Wisconsin Madison, Wisconsin 53706	1
Professor F. A. Tillman Department of Industrial Engineering Kansas State University Manhattan, Kansas 66506	1	Mr. David S. Siegel Code 210T Office of Naval Research Arlington, VA 22217	1

	Copies		Copies
Reliability Analysis Center (RAC)		Professor Franklin A. Graybill	
RADC/RBRAC		Department of Statistics	
Attn: I. L. Krulac		Colorado State University	
Data Coordinator/ Government Programs		Fort Collins, Colorado 80523	1
Griffiss AFB, New York 13441	1		
Mr. Jim Gates		Professor J. S. Rustagi	
Code 9211		Department of Statistics	
Fleet Material Support Office		Ohio State University Research	
U.S. Navy Supply Center		Foundation	
Mechanicsburg, PA 17055	1	Columbus, Ohio 43212	1
Mr. Ted Tupper		Mr. F. R. Del Priori	
Code 92		Code 224	
Fleet Material Support Office		Operational Test and Evaluation	
U.S. Navy Supply Center		Force (OPTEVFOR)	
Mechanicsburg, PA 17055	1	Norfolk, Virginia 23511	1
Mr. Barnard H. Bissinger			
Mathematical Sciences			
Capitol Campus			
Pennsylvania State University			
Middletown, PA 17057	1		
Professor Walter L. Smith			
Department of Statistics			
University of North Carolina			
Chapel Hill, NC 27514	1		
Professor S. E. Fienberg			
Department of Applied Statistics			
University of Minnesota			
St. Paul, Minnesota 55108	1		
Professor Gerald L. Sievers			
Department of Mathematics			
Western Michigan University			
Kalamazoo, Michigan 49008	1		
Professor Richard L. Dykstra			
Department of Statistics			
University of Missouri			
Columbia, Missouri 65201	1		

UNCLASSIFIED

an element
of R^2 -squared

The variation of temperature or of pollutant concentrations over a geographic area are adequately represented by random fields. Given a real-valued random field $\{Z(x), x \in R^2\}$ a basic problem is to interpolate Z over an area A from measurements taken at n stations $x_1, x_2, \dots, x_n$, when the distribution of Z is only partially specified. This is the motivation of the present paper,

It is shown that if the joint distributions are Gaussian the best linear unbiased predictor^(B.L.U.E.) is (among other properties) admissible when used to predict Z at a single point, but inadmissible in general when used to predict the values of the field at several points. A Stein-like predictor is produced which is uniformly better than the B.L.U.E. in the latter case. A non-linear predictor, based on relaxing the unbiasedness condition on the B.L.U.E., is also proposed and shown to be in some cases preferable.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 16	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9
4. TITLE (and Subtitle) Some properties of best linear unbiased predictors and related predictors.		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) 10 André Cabannes		6. PERFORMING ORG. REPORT NUMBER
8. CONTRACT OR GRANT NUMBER(s) 15 N00014-75-C-0555		
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Mathematics M.I.T. Cambridge, MA. 02139 14 TR-16		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (NR-C42-331)
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics & Probability Program Code 436 Arlington, Virginia 22217		12. REPORT DATE Nov. 30, 1979
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 11 30 Nov 79		13. NUMBER OF PAGES 29 12 38
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) best linear unbiased predictor; multivariate normal distribution; admissibility; James-Stein estimation.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See reverse side		

220 021

JCB