

AD-A080 463

BASEL UNIV (SWITZERLAND) MATHEMATICS INST
TOPICS IN ANALYSIS, (U)
SEP 79 A M OSTROWSKI

F/6 12/1

DA-ERO-78-6-068
NL

UNCLASSIFIED

| of |
42
3080462



END
DATE
FILMED
3-80
DRC

LEVEL

(12) JC

(6)

TOPICS IN
ANALYSIS.

A080447

A080463

DDO
RECEIVED
FEB 8 1980
E

(9)

FINAL TECHNICAL REPORT

1 Jul 78 - 30 Sep 79

Principal Investigator

(10)

Alexander M. Ostrowski

(12) 7/1

July, 1 1978 - September, 30 1979

EUROPEAN RESEARCH OFFICE

(11)

Sep 79

United States Army

(15)

Contract Number DAERC-78-G-068

Mathematical Institute, University of Basel

Switzerland

(16)

1T161102BH57

(11)

147

DDC FILE COPY

Approved for public release ; distribution unlimited

80 2

7 025

053 025

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

replacing the values of the functions considered at a and b with their values at a+0 and b-0. In this way the result of Marchand can be improved. Further improvements are pointed out for Sylvester's theorems and Newton's Rule.

The reversible transformation of space elements can be characterized, starting from ordinary one-to-one transformations between two convenient spaces and the complete sets of integrals of certain partial differential equations, which can be expressed by convenient determinants.

UNCLASSIFIED

Table of content

I. On Newton-Sylvester's theorems

II. Reversible transformations of space elements

page
2
4

Appendices:

- BMN 52, On a generalization of the Newton-Sylvester inequalities, pp. 1-52 ;
- BMN 53, On reversible transformations of space elements, pp. 1-90 .

I. The paper contains a simplified proof of a sharpening of Sylvester's theorems, obtained by replacing the values of the functions considered at a and b with their values at $a+0$ and $b-0$. In this way the result of Marchand can be improved. Further improvements are pointed out for Sylvester's theorems and Newton's Rule.

II. The reversible transformation of space elements can be characterized, starting from ordinary one to one transformations between two convenient spaces and the complete sets of integrals of certain partial differential equations, which can be expressed by convenient determinants.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist.	Avail and/or special
A	

I. On Newton-Sylvester's theorems

1. For

$$(1) \quad f_0(x) = a_0 x^n + a_1 x^{n-1} + \dots + a_n, \quad a_0 = 1$$

form the expressions

$$F_\nu(x) := r_\nu f^{(\nu)}(x) - r_{\nu-1} f^{(\nu-1)}(x) f^{(\nu+1)}(x) \quad (\nu=1, \dots, n-1),$$

$$(2) \quad F_0(x) := f_0(x)^2, \quad F_n(x) := 1,$$

assuming that the positive constants r_ν satisfy the relations

$$(3) \quad r_{\nu+1} = 2r_\nu - r_{\nu-1} \quad (\nu=1, \dots, n-2).$$

Denote by $VP(x_0)$ the number of indices ν with

$$\operatorname{sgn}\left(f^{(\nu-1)}(x_0) f^{(\nu)}(x_0)\right) = -1, \quad \operatorname{sgn}\left(F_{\nu-1}(x_0) F_\nu(x_0)\right) = +1.$$

These are variation permanences corresponding to ν and x_0 . Similarly denote by $PP(x_0)$ the number of indices ν with

$$\operatorname{sgn}\left(f^{(\nu-1)}(x_0) f^{(\nu)}(x_0)\right) = \operatorname{sgn}\left(F_{\nu-1}(x_0) F_\nu(x_0)\right) = +1.$$

Here we have permanence permanences corresponding to ν and x_0 .

Denoting by $N_m(a, b)$ the number of zeros of $f^{(m)}(x)$ in (a, b) , Sylvester's theorems say that the expressions

$$(4) \quad VP(a) - VP(b) - N_0(a, b)$$

$$(5) \quad PP(b) - PP(a) - N_0(a, b)$$

are non-negative and even.

2. The proofs given by Sylvester, Genocchi and de Jonquières were rather incomplete and only Marchand (1913) gave a complete discussion of the theorems in their original form. Difficulties arose mainly about the right interpretation of zeros in the double sequence of $f^{(v)}, F_v$ for a and b . In this respect the solution given by Marchand is not optimal since the optimal results can be only obtained for the open interval (a,b) considering the numbers of VP and PP at $a+0$ and $b-0$. Thus, asking for convenient attribution of the signs at a, b themselves is practically asking the wrong question.

3. In this paper a generalization of Sylvester's theorems is derived assuming only that $f_o^{(v)}(x)$ ($v=0, \dots, n$) are continuous functions of x in $\langle a, b \rangle$ with finite numbers of zeros in $\langle a, b \rangle$, while $f^{(n)}$ does not vanish in (a, b) . We further have to assume that each of the F_v formed accordingly to (2) for a fixed choice of the r_v either identically vanishes or has there only a finite number of zeros. Identically vanishing F_v are assumed to be provided with the plus sign.

Defining for $a < b$ the interval functions

$$(6) \quad \Delta'_m(a, b) := VP_m(a+0) - VP_m(b-0) ,$$

$$(7) \quad \Delta''_m(a, b) := PP_m(b-0) - PP_m(a+0) ,$$

we can formulate our main result as saying that the differences

$$(8) \quad \Delta'_m(a, b) - \left(N_o(a, b) - N_m(a, b) \right) , \quad \Delta''_m(a, b) - \left(N_o(a, b) - N_m(a, b) \right)$$

are non-negative even numbers.

We further give the explicit solution of the problem of obtaining the signs at $a+0$ and at $b-0$ for the values of functions

at a, b and derive as corollaries of our main result corresponding generalizations of Newton's Rule. The paper contains as compared with Marchand's presentation an essentially simplified presentation of the whole subject.

II. Reversible transformations of space elements

Consider, for $n > 1$, the coordinates, x_1, \dots, x_n , of a general point of the n -dimensional space, depending on and arbitrarily often differentiable with respect to m parameters T_1, \dots, T_m . Denote generally the derivatives $\frac{\partial x_\nu}{\partial T_\mu}$ by $p_{\nu\mu}$ ($\nu=1, \dots, n; \mu=1, \dots, m$).

In this paper we are going to consider the transformation

$$(1) \quad y_\nu = Y_\nu^*(x_\nu, p_{\nu\mu}) \quad (\nu=1, \dots, n)$$

where the Y_ν^* are homogeneous of dimension 0 in the $p_{\nu\mu}$ and have the further property:

Differentiating y_ν in (1) with respect to the T_μ and putting

$$q_{\nu\mu} := \frac{\partial y_\nu}{\partial T_\mu}$$

we can, eliminating the $p_{\nu\mu}$ and their derivatives, express the x_ν in function of y_ν and $q_{\nu\mu}$,

$$(2) \quad x_\nu = X_\nu^*(y_\nu, q_{\nu\mu}) \quad (\nu=1, \dots, n)$$

where the X_{ν}^{ξ} are homogeneous of dimension 0 in the $q_{\nu\mu}$; and inversely (1) can be deduced differentiating (2) and eliminating the $q_{\nu\mu}$. The functions X_{ν}^{ξ} , Y_{ν}^{ξ} are assumed arbitrarily often differentiable in their arguments.

Such transformations will be called reversible transformations.

We prove that there exist two sets of k functions

$$(3) \quad r_{\alpha} = r_{\alpha}^*(x_{\nu}, p_{\nu\mu}) \quad , \quad s_{\alpha} = s_{\alpha}^*(y_{\nu}, q_{\nu\mu}) \quad (\alpha=1, \dots, k) \quad ,$$

where each set is independent, and which have the property that the expressions Y_{ν}^{ξ} in (1) and X_{ν}^{ξ} in (2) can be written as

$$(4) \quad Y_{\nu}^{\xi} =: Y_{\nu}(x_{\nu}, r_{\alpha}) \quad , \quad X_{\nu}^{\xi} =: X_{\nu}(y_{\nu}, s_{\alpha}) \quad (\nu=1, \dots, n) \quad .$$

Hence, there exists a one to one transformation between two $(n+k)$ -dimensional spaces (x_{ν}, r_{α}) and (y_{ν}, s_{α}) ,

$$(5) \quad T \quad \begin{cases} y_{\nu} = Y_{\nu}(x_{\nu}, r_{\alpha}) \quad , \quad s_{\alpha} = S_{\alpha}(x_{\nu}, r_{\alpha}) \\ x_{\nu} = X_{\nu}(y_{\nu}, s_{\alpha}) \quad , \quad r_{\alpha} = R_{\alpha}(y_{\nu}, s_{\alpha}) \end{cases} \quad (\nu=1, \dots, n; \alpha=1, \dots, k) \quad .$$

The main problem of the paper is to find characteristic properties of the transformations (5).

The method of the paper consists in studying the system of partial differential equations

$$(6) \quad J_{\mu, \alpha} U := \sum_{\nu=1}^n X'_{\nu s_{\alpha}} U'_{p_{\nu\mu}} = 0 \quad (\mu=1, \dots, m; \alpha=1, \dots, k) \quad ,$$

$$(7) \quad \Delta_{\mu, \lambda} U := \sum_{\nu=1}^n p_{\nu\lambda} U'_{p_{\nu\mu}} = 0 \quad (\mu, \lambda=1, \dots, m) \quad ,$$

where U is a function of the $x_\nu, r_\lambda, p_{\nu\mu}$.

The integrals of these equations are the so called functions with the property U. The maximal number of such independent integrals is

$$N = m(n-k-m+d) .$$

Further, we prove that both systems (6) and (7) are complete as well as the system consisting of (6) and (7) taken together. Further, the system (6) and the system (7) are independent, while both systems taken together are not necessarily so. The discussion depends essentially on the number of independent relations existing between (6) and (7). This number can be denoted by dm , where d has the values $0, 1, \dots, m$. But the cases $d > 0$ are exceptional.

For $d=0$ we can write, using k indefinitely often differentiable arbitrary functions,

$$(8) \quad r_\lambda = \varphi_\lambda(U^{(k+1)}, \dots, U^{(k+N)}) \quad (\lambda=1, \dots, k) ,$$

where $U^{(\sigma)}$ are N independent integrals of (6) and (7). The $U^{(\sigma)}$ can be formed, using convenient determinants of the $X'_{\nu s_x}$ and $p_{\nu\mu}$, where the $X'_{\nu s_x}$ are expressed in x_ν, r_λ and $p_{\nu\mu}$. However, in order to obtain the r_λ solving the k equations (8), certain determinants must be assumed to be $\neq 0$.

As to the exceptional cases ($d > 0$), we give a complete solution for $d=1$ and $d=m$, obtained by some special methods. We obtain further, using such methods, the relation

$$(9) \quad k \leq \frac{n-m}{d+1} + \theta \quad , \quad 0 < \theta \leq \frac{d}{d+1} .$$

The author was no longer able to discuss the applications of the theory to the discussion of the partial differential equations solvable without integration.