

AD-A081 173

TEXAS A AND M UNIV COLLEGE STATION INST OF STATISTICS

F/G 12/1

EFFICIENT ESTIMATION OF MULTIVARIATE MOVING AVERAGE AUTOCOVARIA--ETC(U)

MAY 79 H J NEWTON

N00014-78-C-0599


UNCLASSIFIED TR-N-9

NL

| OF |
AD
A081173



END
DATE
FILMED
3-80
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

DDC FILE COPY

ADA 081173

North of Station
Phone 713-465-1141

TEXAS A&M UNIVERSITY
COLLEGE STATION, TEXAS 77843

EFFICIENT ESTIMATION OF MULTIVARIATE MOVING

AVERAGE AUTOCOVARIANCES

By H. J. Newton

Texas A&M University

Technical Report No. N-9

May 1979

Texas A & M Research Foundation
Project No. 3838

"Multiple Time Series Modeling and Time
Series Theoretic Statistical Methods"

Sponsored by the Office of Naval Research

Professor Emanuel Parzen, Principal Investigator

Approved for public release; distribution unlimited.

LEVER

DTIC
DELETED
FEB 28 1980
C

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)		REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. GOVT ACCESSION NO.	1. REPORT NUMBER	2. GOVT ACCESSION NO.	1. REPORT NUMBER	2. GOVT ACCESSION NO.
Technical Report No. N-9		1. REPORT NUMBER	2. GOVT ACCESSION NO.	1. REPORT NUMBER	2. GOVT ACCESSION NO.
4. TITLE (and Subtitle)		5. AUTHOR		6. PERFORMING ORG. REPORT NUMBER	
Efficient Estimation of Multivariate Moving Average Autocovariances		H. Joseph Newton			
7. AUTHOR		8. PERFORMING ORGANIZATION NAME AND ADDRESS		9. CONTRACT OR GRANT NUMBER(s)	
		Texas A&M University Institute of Statistics College Station, TX 77843			
		11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT PROJECT TASK AREA & WORK UNIT NUMBERS	
		Office of Naval Research Code 436 Arlington, VA 22217			
		14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)	
				Unclassified	
16. DISTRIBUTION STATEMENT (of this Report)		17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		18. SUPPLEMENTARY NOTES	
Approved for public release; distribution unlimited.		NA			
		19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		20. ABSTRACT (Continue on reverse side if necessary and identify by block number)	
		Moving average model; multiple time series; autocovariance; generalized least squares; spectral density		This paper proposes a method for estimating the autocovariances of a d-dimensional moving average process of order q. The estimators have the same asymptotic covariance matrix as those obtained by maximizing a Gaussian likelihood, and are obtained by performing a generalized least squares regression of the periodogram on the autocovariance, thus extending Parzen's (1971) estimators for d = 1.	

FORM 7-74 1473 EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-UF-014-6001

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

80 2 19 134 347380

yfb

$$Y(t) = \sum_{k=0}^q B(k) \varepsilon(t-k),$$

where the $B(k)$'s are $d \times d$ real matrices, and $B(0) = I_d$, the d -dimensional identity matrix.

Phadke and Kedem (1978) survey methods for obtaining estimators of $\hat{\varepsilon}$ and the $B(k)$'s from a sample realization $Y(1), \dots, Y(T)$. The methods generally consist of maximizing an exact or approximate Gaussian likelihood to obtain estimators $\hat{B}(1), \dots, \hat{B}(q)$, and $\hat{\varepsilon}$ that are called least squares estimators or a maximum likelihood identification.

Parzen (1971) proposed a method for estimating the moving average autocovariances $R(v) = E(Y(t)Y^T(t+v))$ for $d = 1$ by capitalizing on the linear relationship between the $R(v)$'s and the spectral density $f(\cdot)$ of $Y(t)$:

$$f(\omega) = \frac{1}{2\pi} \sum_{|v| \leq q} R(v) e^{-i v \omega}, \quad \omega \in [-\pi, \pi], \quad (1)$$

where

$$R(v) = R^T(-v) = \begin{cases} \sum_{k=0}^{q-v} B(k) \hat{\varepsilon}^T(k+v), & v = 0, \dots, q \\ 0 & , \quad v > q. \end{cases}$$

Newton, in a Buffalo PHD thesis showed that the Parzen estimators have the same asymptotic covariance matrix as those obtained by maximizing the approximate Gaussian likelihood.

The purpose of this paper is to provide an extension of the Parzen method for $d > 1$. The extension is given in section 2 while conclusions are given in section 3.

Efficient Estimation of Multivariate Moving

Average Autocovariances

By H. J. Newton

Institute of Statistics, Texas A&M University, College Station, Texas

Summary

This paper proposes a method for estimating the autocovariances of a d -dimensional moving average process of order q . The estimators have the same asymptotic covariance matrix as those obtained by maximizing a Gaussian likelihood, and are obtained by performing a generalized least squares regression of the periodogram on the autocovariances, thus extending Parzen's (1971) estimators for $d = 1$.

Some key words: Moving average model; multiple time series; autocovariance, generalized least squares; spectral density

1. INTRODUCTION

Let $\{\varepsilon(t), t \in \mathbb{Z}\}$ be a d -dimensional white noise series on the integers with mean zero and positive definite covariance matrix $\hat{\varepsilon}$. A d -dimensional moving average process $\{Y(t), t \in \mathbb{Z}\}$ of order q is defined by

2. GENERALIZED LEAST SQUARES ESTIMATORS OF $R(\cdot)$

Consider the sample spectral density or periodogram

$$f_T(\omega_j) = \frac{1}{2\pi T} \sum_{s,t=1}^T Y(s)Y^T(t) e^{-i(\omega-\omega_j)T} \quad \omega_j = \frac{2\pi j}{T}, \quad j=1, \dots, \left[\frac{T-1}{2}\right] = N,$$

where $[x]$ denotes the greatest integer less than or equal to x . By inspection of Hannan's Corollary 1 (see Hannan (1970), p. 249)

$$\lim_{T \rightarrow \infty} E(\text{vec}(f_T(\omega_j))) = \text{vec}(f(\omega_j)) \quad (2)$$

$$\lim_{T \rightarrow \infty} \text{Cov}(\text{vec}(f_T(\omega_j)), \text{vec}(f_T(\omega_k))) = \delta_{jk} f^T(\omega_j) \otimes f(\omega_j),$$

where δ is the Kronecker delta, $C \otimes D$ is the Kronecker product $(C_{jk}D)$

of the matrices C and D , and if A is an $n \times m$ matrix with columns

$a_1, \dots, a_m$, then $\text{vec}(A)$ is the $nm \times 1$ vector $(a_1^T, \dots, a_m^T)^T$.

Thus combining (1) and (2) we can write

$$\text{vec}(f_T(\omega_j)) = \text{vec} \left(\sum_{|v| \leq q} R(v) e^{-i\omega_j v} \right) + e_T(\omega_j), \quad j = 1, \dots, N, \quad (3)$$

where $e_T(\omega_1), \dots, e_T(\omega_N)$ are asymptotically uncorrelated d -dimensional random variables with asymptotic mean zero and variance $f^T(\omega_j) \otimes f(\omega_j)$.

To write (3) in a regression form we define the permutation

matrices P and G as follows. Let e_j denote a vector of zeros with

a 1 in column j . The k^{th} row of the $d^2 \times d^2$ matrix P is given by

e_{jd+L} where $L = \left[\frac{k-1}{d}\right]$ and $j = (k-1) - d \left[\frac{k-1}{d}\right] \equiv \text{mod } (k-1, d)$. Letting

$s = d(d+1)/2$, the k^{th} row of the $d^2 \times s$ matrix G is e_j^T where

$j = s - (d-m+1)(d-m+2)/2 - (L-m-1)$, with $L = \max(u, v)$, $m = \min(u, v)$,

and $u = \text{mod } (k-1, d) + 1$, $v = \left[\frac{k-1}{d}\right] + 1$.

Then

$$\begin{aligned} \text{vec}(R(-v)) &= \text{vec}(R^T(v)) = P \text{vec}(R(v)), \quad v=1, \dots, q \\ \text{vec}(R(0)) &= G \text{vec}(R(0)), \end{aligned} \quad (4)$$

where $\text{vec}(R(0))$ contains the distinct elements of the symmetric matrix $R(0)$ and is obtained by applying the vec operator to only the lower triangular portion of $R(0)$.

Using (4) we write

$$\text{vec} \left(\sum_{|v| \leq q} R(v) e^{-i\omega_j v} \right) = G \text{vec}(R(0)) + \sum_{v=1}^q (P e^{-i\omega_j v} e^{-i\omega_j v}) \text{vec}(R(v)) \quad (5)$$

$$= X(\omega_j) R, \quad j = 1, \dots, N,$$

where $R^T = ([\text{vec}(R(0))]^T, [\text{vec}(R(1))]^T, \dots, [\text{vec}(R(q))]^T)^T$ contains the distinct elements of $R(v)$, $|v| \leq q$, and

$$X(\omega_j) = (G, P e^{-i\omega_j} e^{-i\omega_j}, \dots, P e^{-i\omega_j} e^{-i\omega_j})$$

is a $d^2 \times (qd^2+s)$ matrix of known complex numbers.

Writing (3) for $j = 1, \dots, N$ and using (5), we have

$$f = XR + e$$

where $f = ([\text{vec}(f_T(\omega_1))]^T, \dots, [\text{vec}(f_T(\omega_N))]^T)^T$ is

an $Nd^2 \times 1$ observation vector, $X = (X^T(\omega_1), \dots, X^T(\omega_N))^T$ is an

$Nd^2 \times (qd^2+s)$ design matrix, and $e = (e_T^T(\omega_1), \dots, e_T^T(\omega_N))^T$ is an

$Nd^2 \times 1$ error vector with block diagonal asymptotic covariance matrix

$$V = \text{diag}(V(\omega_1), \dots, V(\omega_N)), \quad \text{where } V(\omega_k) = f^T(\omega_k) \otimes f(\omega_k).$$

A consistent estimator of $V(\omega_k)$ is given by

Accession For	
THIS GMA&I	☒
DOC TAB	☐
Unannounced	☐
Justification	☐
By	
Distribution/	
Availability Codes	
Dist	Avail and/or special

posed algorithm. Further the closeness to singularity of $\hat{f}$ provides an indication of whether the series generating the data is poorly behaved, i.e. whether the zeros of

$$|H(z)| = \left| \sum_{k=0}^q B(k)z^k \right|$$

are close to the unit circle. If this is the case, significantly improved estimators can be obtained by using the $\hat{R}$ as initial estimators in a direct maximization of the exact likelihood function.

Using the complex sweep operator (Parzen (1967)) one can find $\hat{f}^{-1}(\omega_j)$ by successively finding the inverse of various partitions of $\hat{f}(\omega_j)$, thus simultaneously finding estimators for various subsets of the components of $\hat{Y}(t)$.

Finally, the algorithm of Wilson (1972) can be used to find estimators $\hat{B}(1)$, ..., $\hat{B}(q)$, and $\hat{f}$ from $\hat{R}$.

$$\hat{V}(\omega_k) = \hat{f}^T(\omega_k) \otimes \hat{f}(\omega_k), \text{ where } \hat{f}(\omega_k) = \frac{1}{2\pi} \sum_{|v| \leq q} R_v(v) e^{-i v \omega_k},$$

and

$$R_v(v) = R_v^T(-v) = \frac{1}{T} \sum_{t=1}^{T-v} Y(t)Y^T(t+v), \quad v = 0, \dots, T-1.$$

Then the generalized least squares estimator $\hat{R}$ of R is the solution to

$$(X^* \hat{V}^{-1} X) \hat{R} = X^* \hat{V}^{-1} f,$$

or

$$\hat{R} = \left[\sum_{t=1}^N X^*(\omega_t) \hat{V}^{-1}(\omega_t) X(\omega_t) \right]^{-1} \sum_{t=1}^N X^*(\omega_t) \hat{V}^{-1}(\omega_t) \text{vec}(f(\omega_t)),$$

where A^* denotes the complex conjugate transpose of the complex matrix A .

Using arguments similar to the case $d = 1$ and results for information matrices of multiple time series given by Newton (1978), one can show that $\sqrt{T}(\hat{R} - R)$ is asymptotically normal with mean zero and covariance matrix $K_R = I_R^{-1}$ where the element of I_R corresponding to $R_{jk}(v)$, $R_{lm}(u)$ is given by

$$\int_{-\pi}^{\pi} f_{lm}^{-1}(\omega) f_{jk}^{-1}(\omega) e^{-i(v-u)\omega} d\omega$$

Further, K_R is consistently estimated by

$$\hat{K}_R = \left[\sum_{t=1}^N X^*(\omega_t) \hat{V}^{-1}(\omega_t) X(\omega_t) \right]^{-1}$$

3. CONCLUSIONS

In many applications, estimates of the spectral density are of interest, thus making available the quantities required by the pro-

REFERENCES

- Hannan, E. J. (1970). Multiple Time Series. New York: Wiley.
- Neudecker, H. (1969). Some theorems on matrix differentiation with special reference to Kronecker matrix products. J. Am. Statist. Assoc., 64, 953-964.
- Newton, H. J. (1978). The information matrices of the parameters of multiple mixed time series. J. Mult. Analysis, 8, 317-323.
- Parzen, E. (1967). Empirical multiple time series analysis. Proc. 5th Berk. Symp. on Math. Stat. and Prob., Eds. L. LeCam and J. Neyman, 305-340, Berkeley: U. California Press.
- Parzen, E. (1971). Efficient estimation of stationary time series mixed schemes. Bull. Inst. Int. Statist. 44, 315-319.
- Phadke, M.S. and Kedem, G. (1978). Computation of the exact likelihood function of multivariate moving average models. Biometrika, 65, 511-519.
- Wilson, G. T. (1972). The factorization of Matricial spectral densities. SIAM J. Appl. Math., 23, 420-426.