

AD-A082 105

COLORADO STATE UNIV FORT COLLINS DEPT OF MATHEMATICS F/G 12/1
ON THE EXISTENCE OF STRONG UNICITY OF ARBITRARILY SMALL ORDER.(U)
FEB 80 B L CHALMERS, G D TAYLOR F49620-79-C-0124

UNCLASSIFIED


AFOSR-TR-80-0151

NL

| OF |
/AD-
A082105



END
DATE
FILMED
4-80
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

2

LEVEL II

b. S.

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER 18 AFOSR-TR-80-0151	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON THE EXISTENCE OF STRONG UNICITY OF ARBITRARILY SMALL ORDER	5. TYPE OF REPORT & PERIOD COVERED 9 Interim / repty	
7. AUTHOR(s) B. L. Chalmers and G. D. Taylor	8. CONTRACT OR GRANT NUMBER(s) 15 F-49620-79-C-0124 VNSF-MCS 76-08518	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Colorado State University Department of Mathematics Fort Collins, CO 80523	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 16 2304 A3 17	
11. CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NM Bldg. 410 Bollin AFB, D.C. 20332	12. REPORT DATE 11 Feb 1980	13. NUMBER OF PAGES 6 pages
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 128	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DTIC ELECTE S MAR 18 1980 D		
18. SUPPLEMENTARY NOTES B		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Uniform approximation theory with constraints, strong uniqueness. $P_1(x) > \sigma = \emptyset$ with the constraints $f - P > \sigma = \text{to}$ $f - P_{\text{sub}} f + \delta \times P - P_{\text{sub}} f$ to the $1/2m$ power		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is shown that the strong uniqueness theorem need not hold in its standard form when constraints are imposed on a uniform approximation problem. In particular, it is shown that when approximation with polynomials subject to a monotone constraint ($p^1(x) \geq 0$) and Hermite-Birkhoff interpolating constraints, a best possible result is the inequality $\ f - p\ \geq \ f - p_f\ + \delta \ p - p_f\ ^{1/2m}$ where p is any approximating polynomial satisfying the constraints and the additional condition that (OVER)		

ADA082105

FILE COPY

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED 407344

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. Abstract cont.

norm of $P < \sigma = M$

$P_{sub} f$

$\|p\| \leq M$ (M fixed) and P_f is the unique best approximation to f from the given class of approximants.

ACCESSION for	
NTIS	White Section ☒
DDC	Buff Section ☐
UNANNOUNCED	☐
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. and/or SPECIAL
A.	

Unclassified

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release in accordance with
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

**ON THE EXISTENCE OF STRONG UNICITY
OF ARBITRARILY SMALL ORDER**

B. L. Chalmers¹

Department of Mathematics
University of California
Riverside, California

G. D. Taylor²

Department of Mathematics
Colorado State University
Fort Collins, Colorado

The strong unicity theorem, first given by Newman and Shapiro (4), may be described as follows: Given $C[a,b]$ and W an n -dimensional Haar subspace of $C[a,b]$. Let $f \in C[a,b]$ and $p_f \in W$ be the best approximation to f from W . Then there exists a positive constant γ , depending only on f , such that

$$\|f - p\| \geq \|f - p_f\| + \gamma \|p - p_f\| \quad (1.1)$$

for all $p \in W$ where $\|h\| = \max\{|h(t)| : t \in [a,b]\}$, $h \in C[a,b]$. The extension of this theorem to the setting of monotone approximation has recently been studied by Fletcher and Roulier (3) and Schmidt (5). Specifically, fix an interval $[a,b]$, integers $1 \leq r_0 < \dots < r_k$, signs $\epsilon_i = \pm 1$, $i=0, \dots, k$ and define

$$K = K(r_0, \dots, r_k; \epsilon_0, \dots, \epsilon_k) \text{ by} \\ K = \{p \in \Pi_n : \epsilon_j p^{(r_j)}(x) \geq 0, a \leq x \leq b, j=0, 1, \dots, k \text{ with } k \leq n\} \quad (1.2)$$

where Π_n denotes the class of all real algebraic polynomials of degree $\leq n$.

The study of approximation of $C[a,b]$ by K is called the monotone

¹Research supported in part by the National Science Foundation, under grant MCS-76-08518. Approved for public release
distribution unlimited.

²Research supported in part by the Air Force Office of Scientific Research, Air Force Systems Command, USAF, under contract F-49620-79-C-0124 and by the National Science Foundation, under grant MCS-78-05847.

910 71 3 08

approximation problem. Professor G.G.Lorentz has played a major role in the development of the theory for this problem. See (2) for a brief expository treatment of this problem and an extensive bibliography.

In (3), Fletcher and Roulier constructed an example in $K = \{p \in \Pi_3 : p'(x) \geq 0 \text{ on } [-1,1]\}$ which shows that the best result of form (1.1) that could hold in this setting would be where $\|p - p_f\|$ is replaced by $\|p - p_f\|^2$. Also, some positive results were given that were extended by Schmidt (5). In (5) it is proved that given $f \in C[a,b]$, K as defined in (1.2), $p_f \in K$ the best monotone approximation to f and a positive constant M , there exists $\gamma > 0$ depending only on f and M such that

$$\|f - p\| \geq \|f - p_f\| + \gamma \|p - p_f\|^2 \quad (1.3)$$

for all $p \in K$ satisfying $\|p\| \leq M$.

In (5) one has the following definition: If p_f is the best uniform approximation to $f \in C[a,b]$ from W a subset of $C[a,b]$, we say that p_f is strongly unique of order α ($0 < \alpha \leq 1$) if for each $M > 0$ there is a constant $\gamma > 0$ such that

$$\|f - p\| \geq \|f - p_f\| + \gamma \|p - p_f\|^{1/\alpha}$$

for all $p \in W$ satisfying $\|p\| \leq M$. Thus, these two papers taken together show that in monotone approximation strong unicity of order $1/2$ holds and this is a best possible result.

In this paper we shall show that by taking an appropriate combination of interpolatory constraints with a monotone constraint one obtains an approximation problem in which strong unicity of order $\frac{1}{2m}$, m a positive integer, holds and that this is also a best possible result.

Thus, fix m a positive integer and define $K \subset \Pi_n$ by

$$K = \{p \in \Pi_n : p^{(1)}(x) \geq 0, a \leq x \leq b \text{ and } p^{(2)}(x_0) = \dots = p^{(2m-1)}(x_0) = 0 \text{ for } x_0 \in (a,b) \text{ fixed, } n \geq 2m+1\}. \quad (1.4)$$

Now, by referring to the general theory of (1), one can prove that corresponding to each $f \in C[a,b]$, there exists a unique best approximation,

p_f from K to f . The basic tools of this theory are extreme linear functionals (extremals) of the dual of Π_n corresponding to f and a given $p \in K$. In this particular setting the extremals are as follows. Given $f \in C[a, b]$ and $p \in K$, define for $x \in [a, b]$, e_x^0 on $C[a, b]$ by $e_x^0(g) = g(x)$ for all $g \in C[a, b]$ (point evaluation) and for $x \in [a, b]$, and $1 \leq j \leq 2m$, e_x^j on Π_n by $e_x^j(q) = q^{(j)}(x)$ for all $q \in \Pi_n$. The linear functional e_x^0 , $x \in [a, b]$, is said to be an extremal for f and p provided $|e_x^0(f-p)| = \|f-p\|$. The linear functional e_x^1 , $x \in [a, b]$ is said to be extremal for f and p provided $e_x^1(p) = 0$. Whenever e_x^1 is an extremal for f and p and $x \notin \{a, x_0, b\}$ then an additional extremal called an augmented extremal is also present; namely, the extremal e_x^2 for which $e_x^2(p) = 0$ must also hold (since $p^{(1)}(x) \geq 0$). If $e_{x_0}^1$ is an extremal for f and p , then the linear functional $e_{x_0}^{2m}$ is an augmented extremal for f and p with $e_{x_0}^{2m}(p) = 0$ holding (since $p^{(1)}(x) \geq 0$). If one starts with an extremal set for f and p (which contains $e_{x_0}^2, \dots, e_{x_0}^{2m-1}$) and adds all possible augmented extremals (as described above) to this set, then one has the augmented set of extremals for f and p corresponding to the original extremal set. Observing that these augmented extremal sets always correspond to Hermite-Birkhoff interpolation problems in which every supported block is even, it is relatively straightforward to prove that the maximal augmented extremal set for f and its best approximation, p_f , from K must have $n+2$ elements which span the dual of Π_n . Thus, K is generalized Haar and uniqueness of best approximations holds (1). In addition, suppose p_f is the best approximation to f from K . Then there exists $k \leq n+2$ extremals (e.g. (2)), $E = \{e_i\}_{i=1}^k$, none of which are augmented extremals, for which 0 belongs to the convex hull of $\{\sigma(e)e : e \in E\}$ where $\sigma(e) = \text{sgn}(f(x) - p_f(x))$ if $e = e_x^0$ for some $x \in [a, b]$, $\sigma(e) = 1$ if $e = e_y^1$ for some $y \in [a, b]$ and $\sigma(e_{x_0}^j) = 1$, $j = 2, \dots, 2m-1$. Then, by adjoining to E the set $E^a = \{\text{all augmented extremals corresponding to elements of } E\}$ we must have that the set $E^{\text{aug}} = E \cup E^a$ contains at least $n+2$ elements of Π_n^* which will necessarily span Π_n^* by the fact that every

supported block in the corresponding Hermite-Birkhoff problem is even.

Likewise, we must have that there exists $e \in E$ for which $e = e_x^0$ some $x \in [a, b]$ as otherwise E is also an extremal set for f and $p_f + c$, c any constant, for which 0 is in the convex hull of $\{\sigma(e)e : e \in E\}$ violating uniqueness of best approximation. Using these observations we can now prove

THEOREM. Let $f \in C[a, b]$ and $p_f \in K$ be the best approximation to f from K .

Given $M > 0$ there exists $\gamma = \gamma(f, M) > 0$ such that for $p \in K$ satisfying $\|p\| \leq M$,

$$\|f - p\| \geq \|f - p_f\| + \gamma \|p - p_f\|^{2m}$$

(i.e. strong unicity of order $\frac{1}{2m}$) and this inequality is best possible.

Proof: The proof is an extension of the techniques of Fletcher and Roulier

and Schmidt. If $f \in K$ then $\gamma = (2M)^{1-2m}$ suffices. Thus, assume $f \notin K$. Let

$E = \{e_i\}_{i=1}^k$ be a set of k extremals, which contains $\{e_{x_0}^j\}_{j=2}^{2m-1}$ but contains no augmented extremals, for which 0 is in the convex hull of $\{\sigma(e)e : e \in E\}$. Set

$E^{\text{aug}} = E \cup E^a$. Further, define $E^0, E^1 \subset E$, where $e \in E$ is in E^0 if $e = e_x^0$ for some

$x \in [a, b]$ and $e \in E^1$ if $e = e_y^1$ for some $y \in [a, b]$. Define the semi-norm $\|\cdot\|'$ on

π_h by $\|q\|' = \max\{|\sigma(e)e(q)| : e \in E\}$. Set $Q = \{q = \frac{p_f - p}{\|p_f - p\|} : \|p_f - p\|' \neq 0 \text{ and } p \in K\}$. We

claim that $\inf_{q \in Q} \max_{e \in E} \sigma(e)e(q) = \tau > 0$. Indeed, if there exist $q \in Q$ with

$\max_{e \in E^0} \sigma(e)e(q) \leq 0$. Then from $q = \frac{p_f - p}{\|p_f - p\|}$ with $\|p_f - p\|' \neq 0$ and $p \in K$ we see that

$e(q) \neq 0$ for some $e \in E$ and $e(q) \leq 0$ for all $e \in E^1$. Thus, $\sigma(e)e(q) \leq 0$ for all

$e \in E$ with strict inequality holding at least once. This violates the fact

that 0 belongs to the convex hull of $\{\sigma(e)e : e \in E\}$. Using this lower bound,

we have for $p \in K$ with $\|p_f - p\|' \neq 0$ that there exists $e \in E^0$ for which

$\sigma(e)e(p_f - p) \geq \tau \|p_f - p\|'$. Now observe that (as usual)

$$\|f - p\| \leq \|f - p_f\| + \gamma \|p_f - p\|'$$

As this inequality holds for $\|p_f - p\|' = 0$, we have a strong uniqueness-type

result for the seminorm $\|\cdot\|'$. Next, the norm, $\|p\|_* = \max\{|\sigma(e)p| : e \in E^{\text{aug}}\}$, is

introduced. Thus, there exists a constant $\lambda > 0$ such that $\|p\|_* \geq \lambda \|p\| \forall p \in \pi_h$.

Finally, we claim that there exists $A > 0$ for which $\|p_f - p\|' \geq A (\|p_f - p\|_*)^{2m}$,

$\forall p \in K$ satisfying $\|p\| \leq M$. First observe that $\|p_f - p\|' = 0$ with $p \in K$ implies

$e(p_f - p) = 0 \forall a \in E^{\text{aug}}$ so that $\|p_f - p\|^* = 0$. Now, for $e \in E$, there exists a constant K_1 for which $|e(p_f - p)| \geq K_1 |e(p_f - p)|^{2m}$ as $\|p\| \leq M$. Let $e \in E^{\text{aug}} \setminus E$ and assume that $e = e_{x_0}^{2m}$ (the augmented extremal corresponding to $e_{x_0}^1$). We claim that there exists $K_2 > 0$ for which $|e_{x_0}^1(p_f - p)| \geq K_2 |e_{x_0}^{2m}(p_f - p)|^{2m} \forall p \in K$ satisfying $\|p\| \leq M$. If this is not the case, then corresponding to each integer $v > 0$ there exists $q_v \in K$ with $\|q_v\| \leq M$ for which $|q_v'(x_0)| < \frac{1}{v} |q_v^{(2m)}(x_0)|^{2m}$. Now we may assume that q_v converges uniformly to $q \in K$. Clearly, $q'(x_0) = 0$. We can write $q_v'(x) = q_v'(x_0) + \frac{q_v^{(2m)}(x_0)}{(2m-1)!} (x-x_0)^{2m-1} + s_v(x)(x-x_0)^{2m} = \beta_v + \alpha_v(x-x_0)^{2m-1} + s_v(x)(x-x_0)^{2m}$ where $\beta_v \rightarrow 0, \alpha_v \rightarrow 0$ (as $q^{(2m)}(x_0) = 0$ since $q \in K$), $|s_v(x)| \leq M_1$ for all $x \in [a, b]$, some M_1 independent of v and $q_v'(x) \geq 0 \forall x \in [a, b]$. Thus, $0 \leq \beta_v + \alpha_v(x-x_0)^{2m-1} + M_1(x-x_0)^{2m}$ for $x \in [a, b]$. For v sufficiently large (so that $x \in (a, b)$), set $x - x_0 = \frac{\alpha_v(2m-1)}{2mM_1}$. This gives $\frac{(2m-1)}{M_1} \left(\frac{2mM_1}{2m-1} \right)^{2m} \beta_v \leq \alpha_v^{2m}$ or that there exists a constant K_1 independent of v (sufficiently large) such that $|q_v'(x_0)| \geq K_1 |q_v^{(2m)}(x_0)|^{2m}$ which is our desired contradiction. Finally, if $e \in E^{\text{aug}} \setminus E$ is of the form $e = e_y^2$ some $y \in (a, b) \setminus \{x_0\}$, the above argument (modified) shows that there exists K_3 for which $|e_y^1(p_f - p)| \geq K_2 |e_y^2(p_f - p)|^2 \geq K_3 |e_y^2(p_f - p)|^{2m} \forall p \in K$ satisfying $\|p\| \leq M$ where K_3 is independent of p . By taking A to be the smallest of the constants produced above, we have that $\|p_f - p\|^* \geq A(\|p_f - p\|^*)^{2m}$ implying $\|f - p\| \geq \|f - p_f\| + \gamma \|p_f - p\|^{2m} \forall p \in K$ satisfying $\|p\| \leq M$ with $\gamma = \gamma(M, f) > 0$ independent of p .

To show this result is best possible we construct an example. Fix m a positive integer and let r_1, r_2, r_3 denote the three roots of $p_0(x) = x^{2m+1} + 2x^{2m-1}$ (note $-2 < r_1 < -1, r_2 = -1, 0 < r_3 < 1$). Define $K = \{p \in \Pi_{2m+1} : p'(x) \geq 0, x \in [r_1, r_3], 0 = p^{(2)}(0) = \dots = p^{(2m-1)}(0)\} = \{p(x) = a_0 x^{2m+1} + a_1 x^{2m} + a_2 x + a_3 : p'(x) \geq 0 \text{ on } [r_1, r_3]\}$. Define $g \in C[r_1, r_3]$ by $g(r_1) = \frac{1}{2}, g(-1) = \frac{1}{2}, g(r_3) = \frac{1}{2}$ and extend g linearly to all $[r_1, r_3]$. Set $f = g + 2x^{2m+1}$ and $p_f(x) = 2x^{2m+1}$. Note that $\{-e_{r_1}^0, e_{-1}^0, -e_{r_3}^0, e_1^1\}$ is an extremal set for f and p_f whose convex hull contains the zero of V^* , $V = \{a_0 x^{2m+1} + a_1 x^{2m} + a_2 x + a_3\}$. (Coefficients are: $\alpha_1 = 1, \alpha_2 = 1 + \alpha_3$,

$\alpha_3 = \frac{-r_1^{2m+1}-1}{r_3^{2m+1}+1}$, $\alpha_4 = r_1 + \alpha_2 + \alpha_3 r_3$, respectively.) Thus, p_f is the desired best approximation to f from $K^{(2)}$. Next, define $p_\alpha(x) = p_f(x) + \alpha p_0(x) + 4m\alpha^{2m}x$, for $0 < \alpha \leq \alpha_0$ where α_0 is chosen so small that $|f - p_\alpha| = |g - \alpha[p_0 + 4m\alpha^{2m-1}x]|$ decreases as x moves away from r_i in a neighborhood of $\{r_1, r_2, r_3\}$ for all α ($0 < \alpha \leq \alpha_0$). This can be done since $|g|$ decreases linearly as x moves away from r_i . Hence α_0 can be chosen so small that $\|f - p_\alpha\| = \max_{i=1,2,3} |(f - p_\alpha)(r_i)|$, $0 < \alpha \leq \alpha_0$ $= f(-1) - p_\alpha(-1) = \frac{1}{2} + 4m\alpha^{2m}$. Also, $\|f - p_f\| = \frac{1}{2}$, $\|p_f - p_\alpha\| \geq |p_f(0) - p_\alpha(0)| = \alpha$ and $p'_\alpha(x) = 2(2m+1)x^{2m} + \alpha((2m+1)x^{2m} + 4mx^{2m-1}) + 4m\alpha^{2m}$. Now, for $x > 0$, $p'_\alpha(x) > 0$; for $x \in [r_1, -\alpha]$, the term $2(2m+1)x^{2m}$ dominates showing that $p'_\alpha(x) > 0$ here; and for $x \in [-\alpha, 0]$ the term $4m\alpha^{2m} \geq |4\alpha mx^{2m-1}|$ again implying that $p'_\alpha(x) \geq 0$. Thus $p_\alpha \in K$ and $(\|f - p_\alpha\| - \|f - p_f\|) / \|p_f - p_\alpha\|^\beta \leq \frac{4m\alpha^{2m}}{\alpha^\beta}$. This implies that we must have $\beta \geq 2m$ in order for the strong unicity theorem to hold for this f and p_f .

By suitably selecting g , it can be shown that this weaker strong uniqueness result holds for an f which also satisfies all the constraints of K . Additional results on this topic will appear elsewhere.

REFERENCES

1. B.L.Chalmers, A unified approach to uniform real approximation by polynomials with linear restrictions, Trans.Amer.Math.Soc., 166(1972), 309-316.
2. B.L.Chalmers and G.D.Taylor, Uniform approximation with constraints, Iber.d.Dt.Math.-Verein., 81(1979), 49-86.
3. Y.Fletcher and J.A.Roulier, A counterexample to strong unicity in monotone approximation, preprint.
4. D.J.Newman and H.S.Shapiro, Some theorems on Chebyshev approximation, Duke Math.J., 30(1963), 673-682.
5. D.Schmidt, Strong unicity and Lipschitz conditions of order 1/2 for monotone approximations, preprint.