

U-4082 939

IRT CORP SAN DIEGO CA
MOLECULAR BEAM STUDIES OF LOW ENERGY REACTIONS.(U)
MAR 80 R H NEYNABER, S Y TANG
IRT-8103-010

F/G 7/4


N00014-74-C-0011
ML

UNCLASSIFIED

1 1
2 2
3 3
4 4
5 5
6 6
7 7
8 8
9 9
0 0



END
DATE
FILMED
5-80
DTIC



MICROCOPY RESOLUTION TEST CHART

NATIONAL BUREAU OF STANDARDS-1963-A

LEVEL

13

ADA 082939

IRT 8103-010

FINAL REPORT
OF
MOLECULAR BEAM STUDIES OF LOW ENERGY REACTIONS

PERIOD: 1 SEPTEMBER 1973-31 JANUARY 1980

ONR CONTRACT NO. N00014-74-C-0011

PRINCIPAL INVESTIGATOR: R. H. NEYNABER

MARCH 31, 1980

This document has been approved
for public release and sale; its
distribution is unlimited.

DTIC
SELECTED
APR 9 1980

IRT
Corporation

Instrumentation
Research
Technology

7650 Convoy Court • P.O. Box 80817
San Diego, California 92138

714 565-7171
Telex 69-5412

80 4 7 074

DOC FILE COPY

UNCLASSIFIED


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
		(7) t.
4. TITLE (and Subtitle) FINAL REPORT OF MOLECULAR BEAM STUDIES OF LOW ENERGY REACTIONS.		5. TYPE OF REPORT & PERIOD COVERED FINAL ✓ 1 Sept. 1973 - 31 Jan. 1980
7. AUTHOR(s) R. H./Neynaber S. Y./Tang		6. PERFORMING ORG. REPORT NUMBER IRT-8103-010
		8. CONTRACT OR GRANT NUMBER(s) N00014-74-C-0011
9. PERFORMING ORGANIZATION NAME AND ADDRESS IRT Corporation P.O. Box 80817 San Diego, CA 92138		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 012-115 NR 393-004
11. CONTROLLING OFFICE NAME AND ADDRESS ONR 800 N. Quincy St. Arlington, VA 22217		12. REPORT DATE 31 March 1980
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS (of this report) Unclassified
		15a. DECLASSIFICATION DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Chemi-ionization Ion-molecule reactions Cross sections Molecular beams Reaction rates		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A summary of research performed under ONR Contract No. N00014-74-C-0011 is given. The contract covered the period 1 Sept. 1973-31 Jan. 1980. The report describes merging and crossed-beams studies of chemi-ionization and/or ion-molecule reactions. Included are investigations of the Ne ⁺ -Ar, Ne ⁺ -Kr, Ne ⁺ -Xe, He ⁺ -D, Ar ⁺ -Na, He ⁺ -He ⁺ , He ⁺ -Ne ⁺ , Ar ⁺ -Kr ⁺ , and He ⁺ -Ne ⁺ systems.		

4-1388

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Final Report
of
Molecular Beam Studies of Low Energy Reactions
ONR Contract No. N00014-74-C-0011

1. Contract Description

Chemi-ionization and ion-molecule reactions involving metastable and ground-state atoms are studied by both crossed and merging beams at low relative energies (i.e., 0.01 to 10 or 20 eV).

2. Scientific Problem

Some theories exist for chemi-ionization involving collisions of metastable and ground-state rare gases. There are very little experimental data to test these theories over a range of relative kinetic energy from 0.01 to 10 or 20 eV. We will supply such data. Theoretical work for collisions between two metastables is almost non-existent, and experimental data are scant. We will supply experimental information such as absolute and relative cross sections and branching ratios for associative to Penning ionization. This information should establish patterns to test those calculations that do exist and will stimulate further theory. Our chemi-ionization data also will produce some information on unknown potentials for the systems A^*B and C^*D^* , where A, B, C, and D are atoms and asterisks denote metastables. This information includes well depths and the dependence of the long range potential on internuclear separation.

The composition of keV neutral rare gas beams formed by charge transfer of the rare gas parent ion beam in alkalis is unknown. The beams consist of rare gas metastables (generally in two states) and ground-state atoms. The technique for generating such beams is common, and information on the composition is needed in analyzing data obtained through their use. We have developed a method for obtaining the fraction of ground-state atoms in such beams by studying appropriate ion-molecule reactions. We will apply this

Accession For	NTIS Grant
	DDC TAB
	Unannounced
	Justification
By	
Distribution	
Availability Codes	
Dist	Avalland/or special
	A

method to determine unknown compositions.

No experimental information exists on low-energy resonant or near-resonant charge-transfer reactions between rare gas ions and metastables. Our experiments will supply such information. The data can be used to see if existing theories for charge transfer between ions and ground-state atoms can be extended to this case. We also will investigate energy distributions of product ions from which information on the reaction kinetics can be obtained.

3. Scientific and Technical Approach

Merging-beams techniques will be used for most of the studies. The two reactants of the process under investigation will be merged. Their velocities will be adjusted with respect to each other so that the desired relative energy in the center-of-mass system will be obtained. Product ions resulting from the reaction will be collected to give relative and absolute cross sections, and branching ratios will be obtained when appropriate.

Some crossed-beams measurements will be made of the ion-molecule reactions at relative energies above 1 eV. Again product ions will be measured to obtain cross sections.

4. Results

4.1 Chemi-ionization Studies

Our chemi-ionization studies have been largely of Penning ionization (PI) and associative ionization (AI) using merging-beams techniques. Most of the effort has been devoted to processes in which both reactants are atoms. Some of these reactions can be described as



where R^* is a metastable rare gas and A is a ground-state atom. These reactions are heteronuclear, and, hence, R is different from A. Although we have conducted some studies in which A was N or O, we will concentrate the discussion on those cases where A is a rare gas, atomic deuterium, or Na.

The remainder of the AI and PI reactions studied can be characterized as



where R'^* is a metastable rare gas, and R' can be the same as or different from R .

The R^* (R'^*) represents a composite of the two metastable states of the rare gas. For He the metastable states are 2^3S and 2^1S at energies of 19.8 and 20.6 eV, respectively. For Ne, Ar, and Kr the two metastable states are $3P_2$ and $3P_0$ at energies of 16.62 and 16.71 eV, 11.55 and 11.72 eV, and 9.92 and 10.56 eV, respectively.

Studies of Reactions (1)-(5) were generally made in the range $0.01 \leq W \leq 10$ eV (where W is the interaction energy, i.e., relative kinetic energy of the reactants) by measuring the product ionic currents. In addition, laboratory energy distributions of Penning ions were obtained. From these measurements the following quantities were determined: absolute and relative cross sections for AI and PI (Q_{AI} and Q_{PI}), total ionization cross sections $Q_T (\equiv Q_{AI} + Q_{PI})$, $r \equiv Q_{AI}/Q_{PI}$, and the branching ratio $R (\equiv Q_{AI}/Q_T)$. In addition, information was obtained on the dynamics of the collision process and on the long-range form and the well depth ϵ^* of the potential curve $V^*(R)$ of the reactants.

An effort has been made to observe patterns in Q_{AI} , Q_{PI} , Q_T , and R that can be related to the attractive or repulsive nature of the reactant system. (A system is defined as attractive if $W/\epsilon^* \ll 1$, where W is in the thermal energy range, i.e., near 0.05 eV, and repulsive if $W/\epsilon^* \gg 1$.) Such patterns could lead to some degree of predictability for the unmeasured Q and R of a given attractive or repulsive system.

The ideal situation would be to have a theory that could be applied to obtain dependable Q 's and R 's. A few theoretical models have been devised and have been applied to some systems. In their more sophisticated forms these have been difficult to use even for simple systems and, in general, require a priori knowledge of $V^*(R)$, the product potential curve $V^+(R)$, and the coupling width $\Gamma(R)$ for $V^-(R)$ and $V^+(R)$. These functions are difficult to obtain. When comparisons of Q and R can be made with experimental results over a wide energy range (e.g., $0.01 \leq W \leq 10$ eV), they are often disappointing. Because of these

theoretical difficulties, There is some merit in our empirical approach to estimate, at least crudely, the Q and R of a given system.

Our chemi-ionization results can best be summarized in tabular form. Shown below are Tables I and II¹. (References in these tables are given in Ref. 1 and reproduced in this report after the presentation of the tables.) Our rare gas beams not only consisted of metastables but also ground-state species. The measured fraction of ground-states species, f_g , is given in Table I. Theoretical fractions calculated using a statistical approach are designated as f_{gt} . The ground-state species do not contribute to chemi-ionization.

TABLE I. Fraction f_g of ground-state species in rare-gas beams formed by charge transfer of rare-gas ions at energy E in alkali vapor at pressure p.^a

Rare gas ^b	Alkali	E (eV)	p ^c (millitorr)	Range of p for const f_g (millitorr) ^d	f_{gr} ^e	f_g ^e	f_{gt} ^f
Ne	Cs	1.1	0.5	0.13-1.1	0.12	0.14	0.17
		1.6	0.6	0.5-1.8	0.14	0.16	0.13
		4.0	0.5	0.06-2.0	0.12	0.14	0.13
Ne	Na	1.1	0.8	---	0.05	0.06	<0.02
		1.6	0.8	---	0.05	0.06	<0.02
		4.0	0.8	0.2-1.5	0.05	0.06	0.03
³ Ne	Na	5.8	0.8	---	0.11	0.13	0.10
Ne	Ne	1.3 ^g	0.48 ^g	---	0.45	0.49 ± 0.04	0.42
		2.75	0.65	0.04-3.7	0.53	0.57 ± 0.05	0.50
		5.5	0.62	0.23-1.5	0.53	0.57 ± 0.05	0.50
Ar	Rb	1.0	0.15	---	0.56	0.61 ± 0.05	0.54
		2.0	0.15	---	0.58	0.62 ± 0.05	0.53
		4.0	0.6	---	0.62	0.66 ± 0.05	0.51
		5.0	0.6	0.12-1.0	0.60	0.63 ± 0.05	0.50
Kr	Cs	4.5	0.04	0.04-0.14	0.60	0.64 ± 0.07	0.60
		5.5	0.14	0.04-0.44	0.65	0.68 ± 0.06	0.56

^aThis table gives our most up-to-date values for f_g .

^bExcept when noted Ne denotes ⁴Ne.

^cValues are accurate to within a factor of 2. Length of charge-transfer cell is 10 cm.

^dPresumably f_g is constant for p less than minimum of range. Measurements were not made where a range is not specified except as noted by footnote g.

^eThe $\nu/\gamma = 1$ for f_{gr} and 1.16 for f_g . The ν 's are defined in Ref. 10. The f_{gr} and f_g for Ne are upper bounds, see Ref. 10 for the explanation.

^fThe f_{gt} contain undesignated errors due to the use of calculated charge-transfer O's.

^gThe p and E for this measurement result in multiple collisions and quenching of Ne⁺ in the cell. Thus, the associated f_{gr} and f_g are unduly large.

TABLE II. Data for various molecular systems

System ^a	Researchers ^b	s ^c	v ^c (eV)	W(eV) or equivalent ^d	P	$G_0 (10^{-16} \text{ cm}^2)^2$ or $a (10^{11} \text{ cm}^3/\text{sec})^2$	Ref.
He ⁺ -Ar	Neynaber & Magnuson	0.31±0.03	--	0.033	0.28±0.08	30.5 ⁺¹² ₋₁₀	8(a)
	Tang et al	0.31±0.007	--	0.033	--	9.1±0.4 ^h	14(a), 14(b)
	Graber et al	--	--	0.033	0.38±0.005	--	15
	West et al	--	--	435e	0.31±0.03	14.7±2.9	16
He ⁺ -Ne	Bran et al	--	--	300e	--	10	17
He ⁺ -Ar	Neynaber & Magnuson	0.30±0.04	--	0.05	0.30±0.05	12.6 ^{+6.2} _{-5.3}	18
	Tang et al	0.30±0.007	--	0.05	--	8.20±0.4 ^h	14(a), 14(b)
	Graber et al	--	--	330e	0.32±0.005	--	15
	Hickman	--	--	0.05	0.32±0.02	--	19
	West et al	--	--	435e	0.30±0.04	17.6±3.5	16
He ⁺ -Ne	Bran et al	--	--	300e	--	11	17
He ⁺ -Xe	Neynaber & Tang	0.46±0.05	--	0.05	0.24±0.04	14.0±5.2	13
	Tang et al	0.46±0.006	--	0.05	--	8.9±0.4 ^h	14(a), 14(b)
	Illenberger	--	--	0.05	0.20±0.07	--	20
	West et al	--	--	435e	0.24±0.03	14.0±5.2	16
	Bran et al	--	--	300e	--	12	17
He ⁺ -Kr	Bran et al	--	--	300e	--	8±7.4	17
He ⁺ -Rn	Huestis et al	--	--	300e	--	6±7.5 ⁱ	21
He ⁺ -C	Neynaber & Tang	0.34	2.1	0.05	0.23±0.04	24.0±2	8(b)
He ⁺ -N ₂	Hargner & Honeus	--	--	0.045	0.19±0.006	33.6±7	22
He ⁺ -O ₂	Fort et al	--	--	0.05	0.27±0.05	24.0±4	8(c), 8(d)
He ⁺ -CO	Hickman & Hargner/Tr	--	--	0.05	0.16	36.5	8(c)
He ⁺ -H ₂	Hickman & Hargner/Tr	--	--	0.05	0.20	34.0	8(c)
	Hickman et al	--	2.44	--	--	--	23
	Miller & Schoefer/Tr	--	1.9 ^j	--	--	--	24
	Hickman et al (Th)	--	2.07	--	--	--	25
Ar ⁺ -Ne	Neynaber & Magnuson	0.33	0.3±0.4	0.033	0.08±0.017	31.2±15.5	26
Ne ⁺ -Ne	Neynaber et al	0.38	0.6	0.033	0.04±0.006	112±30	12
He ⁺ -Ne	Garrett et al (Th)	--	0.56±0.6 ^k	0.033	0.74	92	7(n)
	Deloche et al	--	--	0.033	0.70±0.12	84±17	27
	Hyers & Cunningham	--	--	0.033	--	101±25	28
He ⁺ -Ne	Neynaber & Tang	--	0.5	0.033	<0.06±0.007 ^l	14 ^h ±44 ^f 39±39	29
Ar ⁺ -Ar	Neynaber & Tang	0.40	0.2±0.4	0.033	0.035±0.007 ^l	308±67	30

^aUnless specified He⁺ is ⁴He⁺
^bMore work is experimental unless (Th) for theoretical is specified. The list of researchers is not complete in all cases.
^cThe s is defined by $G_0 = v s^2$. The max v for validity is 0.03 eV for He⁺-Ar, He⁺-Ne, and He⁺-Xe, 0.04 eV for Ar⁺-Ne, 0.1 eV for He⁺-C and He⁺-N₂, and 10 eV for Ar⁺-Ar.
^dThis is the energy associated with the values of P and G₀ in the table.
^eUnless specified the value shown is G₀. Otherwise it is the rate coefficient k at 300K.
^fThese limits apply because P₀ of He⁺ was measured but not P₀ of Ne.
^gA recent measurement at 18° of G₀(=0.033 eV) for the He⁺-Ar system resulted in a value about 16% less than shown.
^hThis value is larger than that in Reference 14(a) to account for a more refined value of v as given in Reference 14(d).

REFERENCES FOR TABLES I AND II

[From R. H. Neynaber, "Merging Beams Experiments with Excited Atoms," *Electronic and Atomic Collisions*, ed. by N. Oda and K. Takayanagi (North-Holland Publishing Co., Amsterdam, 1980), pp. 287-300.

8. See, for example, (a) R. H. Neynaber and G. D. Magnuson, *Phys. Rev.* **A11**, 865 (1975); (b) R. H. Neynaber and S. Y. Tang, *J. Chem. Phys.* **69**, 4851 (1978); (c) A. P. Hickman and H. Morgner, *J. Chem. Phys.* **67**, 5484 (1977); (d) J. Fort, J. J. Laucagne, A. Pesnelle, and G. Watel, *Phys. Rev.* **A18**, 2065 (1978).
10. R. H. Neynaber and G. D. Magnuson, *J. Chem. Phys.* **65**, 5239 (1976) and R. H. Neynaber and S. Y. Tang, *Chem. Phys. Lett.* **65**, 150 (1979).

12. R. H. Neynaber, G. D. Magnuson, and S. Y. Tang, *J. Chem. Phys.* 68, 5112 (1978).
13. R. H. Neynaber and S. Y. Tang, *J. Chem. Phys.* 70, 4272 (1979).
14. (a) S. Y. Tang, A. B. Marcus, and E. E. Muschlitz, Jr., *J. Chem. Phys.* 56, 566 (1972); (b) M. R. Woodward, R. C. Sharp, M. Seeley, and E. E. Muschlitz, Jr., *J. Chem. Phys.* 69, 2978 (1978).
15. H. L. Kramer, J. A. Herce, and E. E. Muschlitz, Jr., *J. Chem. Phys.* 56, 4166 (1972).
16. W. P. West, T. B. Cook, F. B. Dunning, R. D. Rundel, and R. F. Stebbings, *J. Chem. Phys.* 63, 1237 (1975).
17. J. M. Brom, Jr., J. H. Kolts, and D. W. Setser, *Chem. Phys. Lett.* 55, 44 (1978).
18. R. H. Neynaber and G. D. Magnuson, *Phys. Rev.* A14, 961 (1976).
19. A. Niehaus, *Ber. Bunsenges. Phys. Chem.* 77, 632 (1973).
20. E. Illenberger, *Diplom-Arbeit, Universitat Freiburg*, 1971.
21. D. L. Huestis, R. M. Hill, H. H. Nakano, and D. C. Lorents, *J. Chem. Phys.* 69, 5133 (1978).
22. H. Morgner and A. Niehaus, *J. Phys.* B12, 1805 (1979).
23. H. Hotop, E. Illenberger, H. Morgner, and A. Niehaus, *Chem. Phys. Lett.* 10, 493 (1971).
24. W. H. Miller and H. F. Schaefer, *J. Chem. Phys.* 53, 1421 (1970).
25. A. P. Hickman, A. D. Isaacson, and W. H. Miller, *J. Chem. Phys.* 66, 1483 (1977).
26. R. H. Neynaber and G. D. Magnuson, *J. Chem. Phys.* 67, 430 (1977).
27. P. Deloche, P. Monchicourt, M. Cheret, and F. Lambert, *Phys. Rev.* A13, 1140 (1976).
28. G. Myers and A. J. Cunningham, *J. Chem. Phys.* 67, 247 (1977).
29. R. H. Neynaber and S. Y. Tang, *J. Chem. Phys.* 67, 5619 (1977).
30. R. H. Neynaber and S. Y. Tang, *J. Chem. Phys.* 71, 3608 (1979).

Some comments about the results in these tables are:

- a) The theoretical f_{gt} in Table I are in quite good agreement with the measured values indicating the validity of predicting the populations of the rare-gas states by a statistical approach.
- b) Our experimental values of ϵ^* in Table II are for attractive systems and are in good agreement with the ϵ^* of others when comparisons can be made. Those systems with no values of ϵ^* are repulsive systems.
- c) Except for He^*-D , the R (as seen in Table II) for attractive systems are quite small (generally $<10\%$) and, at thermal energies, considerably lower than the R for the repulsive systems.
- d) From Table II one notes that the Q_T for $\text{R}^*-\text{R}'^*$ systems [see Reactions (3), (4), and (5)] are considerably greater than those for R^*-A [see Reactions (1) and (2)] systems.

More details, additional comments, and explanations of some of the conclusions above have been presented.¹ Furthermore, comparisons of relative Q_T versus W and R versus W for the repulsive and attractive systems in Table II have been given. One interesting observation from such comparisons is that the Q_T and R curves are nearly the same for the repulsive Ne^*-Ar , Ne^*-Kr , and Ne^*-Xe systems, whereas for attractive systems such similarities do not exist.

Chemi-ionization studies under the contract have also been done on a few systems not shown in Table II. These are the He^*-H_2 ,² He^*-N ,³ and He^*-O^3 systems.

4.2 Ion-Molecule Studies

In addition to the chemi-ionization work the merging-beams apparatus has been used on this contract to study the charge transfer between $\text{He}^+(1S)$ and Ne^* in the energy range $0.1 \leq W \leq 500$ eV.⁴ The cross section monotonically increases with W . The threshold for the reaction is near 0.1 eV. A modified Demkov approach is used to calculate cross sections, which agree very roughly with the experimental values above $W \approx 3$ eV. At low W the agreement is poor.

A crossed-beams study was also done on reactions of C^+ and O^+ with CO_2 . The energy range of C^+ was from 1 to 500 eV.

5. Publications

Publications credited to the contract are listed below.

- a. R. H. Neynaber and G. D. Magnuson, "Associative and Rearrangement Ionization in Collisions of Metastable Helium with H₂," J. Chem. Phys. 61, 749 (1974).
- b. R. H. Neynaber and G. D. Magnuson, "Product Energies for N₂⁺ + O₂ → N₂ + O₂⁺," J. Chem. Phys. 61, 3490 (1974).
- c. R. H. Neynaber and G. D. Magnuson, "Chemi-ionization in Collisions of Metastable Neon with Argon," Phys. Rev. A11, 865 (1975).
- d. R. H. Neynaber and G. D. Magnuson, "Associative Ionization in Collisions Between Two Excited Reactants," Phys. Rev. A12, 891 (1975).
- e. R. H. Neynaber and G. D. Magnuson, "Associative Ionization in Collisions Between Metastable Helium and Atomic Nitrogen and Oxygen," J. Chem. Phys. 64, 2840 (1976).
- f. J. A. Rutherford and D. A. Vroom, "A Study of the Reactions of C⁺ and O⁺ with Carbon Dioxide," J. Chem. Phys. 64, 3057 (1976).
- g. R. H. Neynaber and G. D. Magnuson, "Penning and Associative Ionization in the Metastable Neon-Krypton System," Phys. Rev. A14, 961 (1976).
- h. R. H. Neynaber and G. D. Magnuson, "Composition of Partially Metastable Rare Gas Beams," J. Chem. Phys. 65, 5239 (1976).
- i. R. H. Neynaber and G. D. Magnuson, "Chemi-ionization in Collisions of Metastable Argon with Sodium," J. Chem. Phys. 67, 430 (1977).
- j. R. H. Neynaber and S. Y. Tang, "Chemi-ionization in Collisions of Metastable Helium with Metastable Neon," J. Chem. Phys. 67, 5619 (1977).
- k. R. H. Neynaber, G. D. Magnuson, and S. Y. Tang, "Chemi-ionization in Collisions of Metastable Helium with Metastable Helium," J. Chem. Phys. 68, 5112 (1978).
- l. S. Y. Tang and R. H. Neynaber, "Charge Transfer Between Helium Ions and Metastable Neon," Phys. Rev. A18, 1925 (1978).
- m. R. H. Neynaber and S. Y. Tang, "Penning and Associative Ionization in the Metastable Helium-Atomic Deuterium System," J. Chem. Phys. 69, 4851 (1978).
- n. R. H. Neynaber and S. Y. Tang, "Chemi-ionization in Collisions of Metastable Neon with Xenon," J. Chem. Phys. 70, 4272 (1979).
- o. R. H. Neynaber and S. Y. Tang, "Composition of Partially Metastable Kr Beams," Chem. Phys. Lett. 65, 150 (1979).
- p. R. H. Neynaber and S. Y. Tang, "Chemi-ionization in Collisions of Metastable Argon with Metastable Krypton," J. Chem. Phys. 71, 3608 (1979).

6. Extenuating Circumstances

None

7. There were no unspent funds remaining at the end of the contract.
8. No graduate students or postdoctoral personnel have been associated with the contract.
9. During the contract, R. H. Neynaber also received partial support from the General Physics Division of the Air Force Office of Scientific Research.

References

1. For more details see R. H. Neynaber, "Merging-Beams Experiments with Excited Atoms," Electronic and Atomic Collisions, ed. by N. Oda and K. Takayanagi (North-Holland Publishing Co., Amsterdam, 1980) pp 287-300.
2. R. H. Neynaber and G. D. Magnuson, J. Chem. Phys. 61, 749 (1974).
3. R. H. Neynaber and G. D. Magnuson, J. Chem. Phys. 64, 2840 (1976).
4. S. Y. Tang and R. H. Neynaber, Phys. Rev. A18, 1925 (1978).

June 1978

REPORTS DISTRIBUTION LIST FOR ONR PHYSICS PROGRAM OFFICE
UNCLASSIFIED CONTRACTS

Director Defense Advanced Research Projects Agency Attn: Technical Library 1400 Wilson Blvd. Arlington, Virginia 22209	3 copies
Office of Naval Research Physics Program Office (Code 421) 800 North Quincy Street Arlington, Virginia 22217	3 copies
Office of Naval Research Assistant Chief for Technology (Code 200) 800 North Quincy Street Arlington, Virginia 22217	1 copy
Naval Research Laboratory Department of the Navy Attn: Technical Library Washington, D. C. 20375	3 copies
Office of the Director of Defense Research and Engineering Information Office Library Branch The Pentagon Washington, D. C. 20301	3 copies
U. S. Army Research Office Box 12211 Research Triangle Park North Carolina 27709	2 copies
Defense Documentation Center Cameron Station (TC) Alexandria, Virginia 22314	12 copies
Director, National Bureau of Standards Attn: Technical Library Washington, DC 20234	1 copy
Commanding Officer Office of Naval Research Branch Office 536 South Clark Street Chicago, Illinois 60605	3 copies

<p> Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91101 </p>	<p>3 copies</p>
<p> San Francisco Area Office Office of Naval Research One Hallidie Plaza Suite 601 San Francisco, California 94102 </p>	<p>3 copies</p>
<p> Commanding Officer Office of Naval Research Branch Office 666 Summer Street Boston, Massachusetts 02210 </p>	<p>3 copies</p>
<p> New York Area Office Office of Naval Research 715 Broadway, 5th Floor New York, New York 10003 </p>	<p>1 copy</p>
<p> Director U. S. Army Engineering Research and Development Laboratories Attn: Technical Documents Center Fort Belvoir, Virginia 22060 </p>	<p>1 copy</p>
<p> ODDR&E Advisory Group on Electron Devices 201 Varick Street New York, New York 10014 </p>	<p>3 copies</p>
<p> Air Force Office of Scientific Research Department of the Air Force Bolling AFB, D. C. 22209 </p>	<p>1 copy</p>
<p> Air Force Weapons Laboratory Technical Library Kirtland Air Force Base Albuquerque, New Mexico 87117 </p>	<p>1 copy</p>
<p> Air Force Avionics Laboratory Air Force Systems Command Technical Library Wright-Patterson Air Force Base Dayton, Ohio 45433 </p>	<p>1 copy</p>
<p> Lawrence Livermore Laboratory Attn: Dr. W. F. Krupke University of California P. O. Box 808 Livermore, California 94550 </p>	<p>1 copy</p>

<p> Harry Diamond Laboratories Technical Library 2800 Powder Mill Road Adelphi, Maryland 20783 </p>	<p>1 copy</p>
<p> Naval Air Development Center Attn: Technical Library Johnsville Warminster, Pennsylvania 18974 </p>	<p>1 copy</p>
<p> Naval Weapons Center Technical Library (Code 753) China Lake, California 93555 </p>	<p>1 copy</p>
<p> Naval Training Equipment Center Technical Library Orlando, Florida 32813 </p>	<p>1 copy</p>
<p> Naval Underwater Systems Center Technical Library New London, Connecticut 06320 </p>	<p>1 copy</p>
<p> Commandant of the Marine Corps Scientific Advisor (Code RD-1) Washington, DC 20380 </p>	<p>1 copy</p>
<p> Naval Ordnance Station Technical Library Indian Head, Maryland 20640 </p>	<p>1 copy</p>
<p> Naval Postgraduate School Technical Library (Code 0212) Monterey, California 93940 </p>	<p>1 copy</p>
<p> Naval Missile Center Technical Library (Code 5632.2) Point Mugu, California 93010 </p>	<p>1 copy</p>
<p> Naval Ordnance Station Technical Library Louisville, Kentucky 40214 </p>	<p>1 copy</p>
<p> Commanding Officer Naval Ocean Research & Development Activity Technical Library NSTL Station, Mississippi 39529 </p>	<p>1 copy</p>
<p> Naval Explosive Ordnance Disposal Facility Technical Library Indian Head, Maryland 20640 </p>	<p>1 copy</p>

Naval Ocean Systems Center Technical Library San Diego, California 92152	1 copy
Naval Surface Weapons Center Technical Library Dahlgren, Virginia 22448	1 copy
Naval Surface Weapons Center (White Oak) Technical Library Silver Spring, Maryland 20910	1 copy
Naval Ship Research and Development Center Central Library (Code L42 and L43) Bethesda, Maryland 20084	1 copy
Naval Avionics Facility Technical Library Indianapolis, Indiana 46218	1 copy