

AD-A084 122


ARMY RESEARCH INST OF ENVIRONMENTAL MEDICINE NATICK MA F/G 6/14  
SUBJECTIVE REGULATION OF EXERCISE INTENSITY BY PERCEIVED EXERTI--ETC(U)  
FEB 80 M A SMUTOK, G S SKRINAR, K B PANDOLF

UNCLASSIFIED

USARIEM-M-8/80

NL

| 4 |  
AC  
A08-022



END  
DATE  
FILMED  
6 80  
DTIC

1

LEVEL II

CLASS 4  
DDC

UNCLAS

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ADA 084122

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER M 8/80 ✓	2. GOVT ACCESSION NO. AD-A084122	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Subjective regulation of exercise intensity by perceived exertion		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) Michael A. Smutok, Gary S. Skrinar, Kent B. Pandolf		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Army Research Institute of Environmental Medicine, Natick, MA 01760		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Same as 9. above (12) 302		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 3E162777A845 (16)
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (14) USARIE M-8/80		12. REPORT DATE 7 Feb 1980 (11)
		13. NUMBER OF PAGES 28
		15. SECURITY CLASS. (of this report) UNCLAS
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DTIC ELECTE S MAY 13 1980 D E		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Exercise prescription; heart rate; perceived exertion; speed; subjective regulation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The primary purpose of this investigation was to determine if a safe and effective physical conditioning heart rate (HR) could be prescribed by perception of exertion. Ratings of perceived exertion (RPE) were requested from ten normal adult males during treadmill exercise trials at 4.7, 6.5, 9.7, 11.3, and 12.9 km/hr (T <sub>1</sub> ). Subjects were then requested to subjectively regulate their own treadmill speed during two separate trials (T <sub>2</sub> and T <sub>3</sub> ) at the RPE reported for each speed during T <sub>1</sub> . Speed and HR at equivalent RPE were compared during		

DDC FILE COPY

DD FORM 1473 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE  
80 5 12 133

UNCLAS 040850  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLAS

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>. Regression analyses revealed that there was no difference in speed across all RPE between the three trials, however, HR was seen to become progressively higher during T<sub>2</sub> and T<sub>3</sub> than during T<sub>1</sub> as speed and RPE decreased. HR reliability was significant ( $p < .05$ ) during running but not significant ( $p > .05$ ) during walking. It is concluded that prescription of exercise by RPE can produce safe, effective and reliable conditioning HR above 150 beats/min (80% HR max) and running speeds above 9 km/hr (5.6 mph). Use of RPE for exercise prescription below these levels can result in inaccurate and unreliable conditioning heart rates. This method of exercise prescription has limitations and could result in dangerously high HR if used in cardiac rehabilitation programs for patients in which strict adherence to target HR is essential.

UNCLAS

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

SUBJECTIVE REGULATION OF EXERCISE INTENSITY  
BY PERCEIVED EXERTION

Michael A. Smutok, M.S., Gary S. Skrinar, Ph.D.,  
and Kent B. Pandolf, Ph.D.

From Sargent College of Allied Health Professions, Boston University, Boston, MA 02215; and Walter Reed Army Medical Center, Washington, D.C. 20012 and U.S. Army Research Institute of Environmental Medicine, Natick, MA 01760. CPT Smutok is currently at Walter Reed Army Medical Center. Dr. Skrinar is employed by Boston University while Dr. Pandolf works at the U.S. Army Research Institute of Environmental Medicine.

Running Head: Rated Perceived Exertion and Exercise Regulation

Key Words: Exercise prescription; heart rate; perceived exertion; speed;  
subjective regulation

Mailing Address: Dr. Kent B. Pandolf  
Military Ergonomics Division  
U.S. Army Research Institute of  
Environmental Medicine  
Natick, MA 01760

Accession For	
NTIS GAMA	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

### Abstract

The primary purpose of this investigation was to determine if a safe and effective physical conditioning heart rate (HR) could be prescribed by perception of exertion. Ratings of perceived exertion (RPE) were requested from ten normal adult males during treadmill exercise trials at 4.7, 6.5, 9.7, 11.3, and 12.9 km/hr ( $T_1$ ). Subjects were then requested to subjectively regulate their own treadmill speed during two separate trials ( $T_2$  and  $T_3$ ) at the RPE reported for each speed during  $T_1$ . Speed and HR at equivalent RPE were compared during  $T_1$ ,  $T_2$  and  $T_3$ . Regression analyses revealed that there was no difference in speed across all RPE between the three trials, however, HR was seen to become progressively higher during  $T_2$  and  $T_3$  than during  $T_1$  as speed and RPE decreased. HR reliability was significant ( $p < .05$ ) during running but not significant ( $p > .05$ ) during walking. It is concluded that prescription of exercise by RPE can produce safe, effective and reliable conditioning HR above 150 beats/min (80% HR max) and running speeds above 9 km/hr (5.6 mph). Use of RPE for exercise prescription below these levels can result in inaccurate and unreliable conditioning heart rates. This method of exercise prescription has limitations and could result in dangerously high HR if used in cardiac rehabilitation programs for patients in which strict adherence to target HR is essential.

KEY WORDS: Exercise prescription; heart rate; perceived exertion; speed; subjective regulation.

The recent popularity of physical conditioning for preventative as well as rehabilitative purposes necessitates an increased awareness and concern for safe and effective exercise prescription. The prescription of conditioning programs for individuals unaccustomed to regular physical exercise involves consideration of the type of physical activity along with the duration, frequency and intensity of that activity. Exercise intensity is often prescribed at approximately 70% of an individual's maximum functional aerobic capacity (maximum oxygen uptake). There is a linear relationship, except when approaching maximum, between oxygen uptake and heart rate during dynamic exercise involving large muscle groups.<sup>1,2</sup> Consequently, the intensity is often prescribed as a percent of maximum heart rate<sup>3,4</sup> or as a percent of maximum heart rate minus resting heart rate in the form of a target heart rate.<sup>5</sup> These methods are widely used today and necessitate the monitoring of heart rate either by some mechanized device (ECG or pulse counter) or by the exerciser counting his own pulse. Although this last method is popular and has proven satisfactory, errors in determining heart rate have been shown to occur due to pulse counting errors,<sup>6</sup> effects of palpation site<sup>7</sup> and also from the time delay encountered during post exercise pulse location.<sup>8,9</sup>

It has been suggested by Borg that some people have difficulty counting heart rate.<sup>10</sup> On the other hand, if no trouble is encountered they tend to become "pulse counters". Thus they focus too much attention on this activity becoming mentally preoccupied with it. Borg's psychophysical scale for ratings of perceived exertion (RPE) has been shown to have a high correlation with heart rate.<sup>10,11,12</sup>

The Borg psychophysical scale has been used almost exclusively to elicit a perceptual response to an exercise stimulus. Little work has been done using the

perceived exertion scale to set and adjust exercise intensity. If subjects are able to set and regulate their exercise intensity by perceptual feelings it may be possible to supplement the exercise prescription with subjective estimates of exertion. Once a subject's perceptual response to a particular standardized exercise (i.e. walking and running) is known it may be feasible to prescribe the intensity by perceptual feelings alone, under the same standard conditions. The primary purpose of this study was to investigate the difference in exercise intensity (speed of walking and running and heart rate) at the same RPE during standardized exercise of predetermined speed and exercise in which speed was subjectively regulated.

## Methods

### SUBJECTS AND DESIGN

Ten healthy male volunteers were selected as experimental subjects. These subjects had an average age (mean  $\pm$  SD) of  $25.5 \pm 2.7$  years (range 20 to 29); height,  $175.6 \pm 7.7$  cm; weight,  $76.4 \pm 9.5$  kg; maximal oxygen uptake,  $55.4 \pm 8.0$  ml/kg $\cdot$ min; maximal heart rate,  $189 \pm 9.1$  beats/min; and maximal rated perceived exertion,  $19.5 \pm 0.5$ . There were no cardiovascular, respiratory, orthopedic or metabolic conditions contraindicating exercise in any of the subjects. No one was currently under the care of a physician nor was anyone taking prescribed medications. All subjects were totally informed as to the plan and risk in the experiment and signed an informed consent form. Maximal oxygen uptake ( $\dot{V}O_2$  max) was measured for each subject using a modified version of the treadmill test described by Taylor et al.<sup>13</sup>

Pre-recorded instructions for the Borg Scale of rated perceived exertion were played while the subject read a written copy. The scale for eliciting RPE is a bipolar scale with numbers from 6 to 20 (see Figure 1). Every odd number has an attached verbal expression ranging from "very, very light" at 7 to "very, very hard" at 19.

---

INSERT FIGURE 1 ABOUT HERE

---

Exercise trials ( $T_1$ ) at 4.7, 6.5, 9.7, 11.3, and 12.9 km/hr (3, 4, 6, 7, and 8 mph) on a level treadmill were administered in order to measure heart rate, oxygen uptake, minute ventilation, and RPE at each speed. Subjects were exercised at each speed, presented in progressive order, for a five minute period. Heart rate, oxygen uptake and rated perceived exertion were measured during the last minute of exercise. An appropriate rest period, determined by the recovery heart rate, followed each trial. If the heart rate during the previous trial did not exceed 110 beats/min the heart rate was allowed to recover to resting before the next trial was started. When the heart rate exceeded 110 beats/min during the previous trial the heart rate was allowed to recover to 100 beats/min before the start of the next trial.

Exercise trials were then administered at several subjective intensities ( $T_2$ ). The subjective intensities were the same RPE values the subject reported during the trials of  $T_1$ . These RPE values were presented in random order. The subjects first listened to pre-recorded instruction explaining the adjustment of treadmill speed according to perceived exertion. They then began walking at 3.5 km/hr (2.2 mph) on the level treadmill and were allowed to adjust their own speed, having access to the treadmill controls; the speed indicator dial was blinded from view. Four minutes were allowed for adjustment at the assigned RPE value. Warnings were given at 2, 3, and 3.5 minutes during the trial so that final adjustments could be made. At the end of the fourth minute the controls were removed from the subject's reach. Measurement of speed, heart rate and oxygen uptake were taken a minimum of one minute following the last speed

adjustment (or between the fifth and sixth minute of each trial). Rest periods as outlined in the previous section followed each trial.

Exercise trials ( $T_3$ ) were administered at the same subjective intensities as  $T_2$  for determining reproducibility. The RPE values were the same; however, the order in which they were presented was randomized once again. All other procedures were identical to those of  $T_2$ . The time period between trials ( $T_1$ ,  $T_2$  or  $T_3$ ) ranged from a minimum of 3 days to a maximum of 7 days.

#### PROCEDURE

Oxygen uptake (STPD) during each trial of  $T_1$ ,  $T_2$ , and  $T_3$  were calculated using the same formulas presented by Consolazio et al.<sup>14</sup> Expired air was collected in a 150 liter meteorological balloon. Oxygen and carbon dioxide concentrations were determined with a Beckman E-2 oxygen analyzer and an LB-2 carbon dioxide analyzer, respectively. Expired volume was measured by drawing the air into a Tissot chain compensated spirometer. Heart rate was determined from a modified  $V_5$  lead (i.e. negative on mid-sternum, positive on  $V_5$ , and ground on right chest) using a Hewlett Packard 1500B electrocardiograph. Overall ratings of perceived exertion were requested by the standardized statement, "rate your exertion".

#### STATISTICAL TREATMENT

Analysis of variance with repeated measures was used to test for differences in speed, heart rate and oxygen uptake of each mean RPE during  $T_1$ ,  $T_2$  and  $T_3$ . Pearson product-moment correlation coefficients and regression equations were computed between RPE and speed, heart rate, minute ventilation and oxygen uptake for  $T_1$ ,  $T_2$  and  $T_3$ . Reliability coefficients between  $T_2$  and  $T_3$

for each variable at corresponding RPE were computed using the Pearson product-moment correlation formula. Mean percent error in heart rate between  $T_1$  and the average of  $T_2$  and  $T_3$  at each RPE was calculated. Mean percent error in heart rate between  $T_1$  and the average of  $T_2$  and  $T_3$  across all RPE was calculated for each individual subject.

### Results

The descriptive data presented in the methods indicate the group to be young adults of average height and weight while maximum  $\dot{V}O_2$  suggests the group is of above average fitness level. The mean RPE reported at maximum confirms an appropriate rating response at maximum exercise levels.

Table 1 illustrates the means of RPE reported at each predetermined speed trial during  $T_1$ . The individual RPE for each subject ranged from 7 at the slowest speed to 18 at the fastest. The RPE values suggest the subjects were able to rate their perceptions of exertion appropriately during low and high intensity exercise. Mean RPE can be seen to increase as speed increased. A walking speed increment of approximately 1.7 km/hr (1 mph) elicited an RPE increment of about 2 rating units. Increments in running speed on the order of 1.7 km/hr (1 mph) resulted in increases in RPE of approximately 1 rating unit. All subjects were able to complete 5 minutes of exercise at each of the speeds with the exception of the 12.9 km/hr (8.0 mph) trial. Three of the subjects were stopped during this trial because their exertion rating reached the predetermined end point of greater than 18.

Each subject walked at 4.7 and 6.5 km/hr (3 and 4 mph) and ran at 9.7, 11.3, and 12.9 km/hr (6, 7 and 8 mph) during  $T_1$ . Subjects chose the same mode of exercise (walking or running) during  $T_2$  and  $T_3$  when regulating their own speed according to RPE reported for the 4.7, 9.7, 11.3 and 12.9 km/hr (3, 6, 7, and 8 mph)  $T_1$  trials. In contrast, six subjects preferred to run during  $T_2$  and  $T_3$  at the RPE level they reported for the 6.5 km/hr (4 mph)  $T_1$  trial. This represented a change in mode of exercise (from walking to running) for 60% of the subjects at mean RPE 10.3 between the standardized ( $T_1$ ) and the subjectively regulated ( $T_2$  and  $T_3$ ) exercise trials.

---

INSERT TABLE 1 ABOUT HERE

---

Figure 2 graphically demonstrates the relationship of the regression lines comparing speed to RPE during  $T_1$ ,  $T_2$  and  $T_3$ . Correlation coefficients ( $r$ ) of .83 and above in  $T_1$ ,  $T_2$  and  $T_3$  demonstrate a significant positive relationship between RPE and speed in each trial. The slopes of individual regression equations are very near .80 in each trial; y-intercepts for each trial are close to each other. The individual linear regression lines relating speed to RPE in  $T_1$ ,  $T_2$  and  $T_3$  can be seen to be nearly parallel and fall in close proximity to one another. The regression lines exhibit this very close relationship at all speeds and RPE.

---

INSERT FIGURE 2 ABOUT HERE

---

Figure 3 displays the relationship of the regression lines relating HR to RPE during  $T_1$ ,  $T_2$ , and  $T_3$ . Correlation coefficients of .85 and above in  $T_1$ ,  $T_2$  and  $T_3$  indicate a strong positive relationship between HR and RPE during each trial. During the two subjectively regulated trials ( $T_2$  and  $T_3$ ) the slopes (.093 and .088) and y-intercepts (-1.96 and -1.39) of the regression equations did not differ appreciably. The similarity of the regression lines for the  $T_2$  and  $T_3$  trials is evident at all RPE and HR. The slope of the individual regression equation for  $T_1$  is not as steep as the slopes for  $T_2$  and  $T_3$ . The y-intercept of  $T_1$  is higher than those for  $T_2$  and  $T_3$ . The individual regression line relating HR to RPE in  $T_1$  diverges from the regression lines in  $T_2$  and  $T_3$  as HR and RPE decline. The standard ( $T_1$ ) and subjectively regulated trials ( $T_2$  and  $T_3$ ) manifest similarities only near maximum HR (> 180 beats/min). A progressive difference in HR at the same RPE occurs between standard and subjectively regulated exercise as HR declines from maximum.

---

INSERT FIGURE 3 ABOUT HERE

---

Table 2 lists the Pearson product-moment correlation coefficients (r) between RPE and speed, HR,  $\dot{V}_E$ , and  $\dot{V}O_2$  during  $T_1$ ,  $T_2$  and  $T_3$ . The r values range from .81 to .90 and all are significant at the .05 level. Each variable

demonstrates a strong positive relationship to RPE during standard as well as subjectively regulated exercise.

Test - retest reliability coefficients for speed, HR and  $\dot{V}O_2$  between  $T_2$  and  $T_3$  are listed in Table 3. Reliability of the subjective regulation of speed was significant above RPE 10. At RPE below 10, HR and  $\dot{V}O_2$  reliability was not significant at the .05 level. At lower RPE ( $< 10$ ), HR and oxygen uptakes at the same RPE are not reliable even though speed is reliably regulated.

---

INSERT TABLE 2 ABOUT HERE

---

---

INSERT TABLE 3 ABOUT HERE

---

Table 4 presents the percent difference in HR between standardized ( $T_1$ ) and the mean of the subjectively regulated trials ( $T_2$  and  $T_3$ ) at the same RPE

level,  $\frac{\overline{HR}_{T_{2,3}} - HR_{T_1}}{HR_{T_1}}$ , for each individual at each RPE. Mean values for

accuracy at each mean RPE reveal 22.3 and 34.7% differences in HR occurring at RPE of 7.9 and 10.3, respectively. HR differences were below 5.5% for RPE above 12. A general trend of improved accuracy with increasing RPE can be seen. Mean percent differences in HR between  $T_1$  and the mean of  $T_2$  and  $T_3$  for each individual across all RPE ranged from 41% for the least accurate

subject (# 6) to 1.2% for the most accurate (# 7). A wide range of intra-individual HR differences across all RPE can be seen. Interindividual HR differences ranged from 100 to 3% for the RPE with the greatest mean difference (10.3) and from 7 to 0.3% for the RPE with the least mean difference (14.6). Above RPE 7.9 interindividual differences decrease as RPE ascends. There was also less variability between subjects as speed increased.

---

INSERT TABLE 4 ABOUT HERE

---

#### Discussion

#### COMPARISON OF SPEED AND HR REGRESSION DURING STANDARD AND SUBJECTIVELY REGULATED EXERCISE

This investigation has demonstrated that at equivalent RPE the subjectively regulated speed during treadmill exercise is the same as the predetermined treadmill speed of a standardized exercise test (Figure 2). On the other hand, when heart rates during standard exercise are compared to subjectively regulated exercise at the same RPE the differences become progressively larger as HR declines (Figure 3). The HR regression lines indicate that at the same RPE, heart rates were higher during subjectively regulated exercise than during the standard exercise test. The HR difference between standard and subjectively regulated exercise is accentuated at lower RPE, speeds and absolute HR levels.

The point at which the HR divergence between standard and subjectively regulated exercise becomes significant has been established statistically. Analysis of mean HR variance at each mean RPE illustrates significant ( $p < .05$ )

differences in HR occur between standard and subjectively regulated exercise below approximately 150 beats/min (80% HR max) or RPE 12. At RPE 12, Figure 3 displays the speed (at which HR become significantly different) to be approximately 9 km/hr (5.6 mph).

Although the regression lines for speed are nearly identical during standard and subjectively regulated exercise the HR regressions demonstrate significant divergence at speeds below 9 km/hr. The discrepancy in HR in the fast walking speed range (6 to 8 km/hr or 4 to 5 mph) may be explained by the fact that during the standard exercise trial all subjects walked when presented with the 6.4 km/hr speed. During the subjective regulation trials six subjects adjusted their speed to running at the same RPE. Noble and associates have shown that at the same RPE, heart rate for running is greater than walking.<sup>15</sup> Åstrand and Rodahl report oxygen uptake to be higher for running than walking at all speeds below 8 km/hr.<sup>16</sup> Assuming linearity with oxygen consumption, HR will also be higher. Since 60% of the subjects in this study preferred to change their mode of exercise (from walking to running) between the standard and subjectively regulated trials, the mode change may be responsible for the higher HR seen in T<sub>2</sub> and T<sub>3</sub>.

Discrepancies in HR at the slower walking speed range (5 to 6 km/hr or 3 to 4 mph) were also noted. The mean speed at the lowest RPE was 2.9 mph during T<sub>1</sub> and 3.3 mph during T<sub>2</sub> and T<sub>3</sub>. The increment of .40 mph between the trials resulted in a significant increase in HR. LeBlanc has reported that in young military men a minimum increment of only .25 mph in walking speed is necessary to evoke a significant increase in HR.<sup>17</sup> The speed increment (demonstrated in the present investigation) of approximately twice the minimum requirement to elicit a significant HR increase is shown not to be perceptually different when using RPE to adjust speed.

Comparison of speed and HR regression during exercise intensities reported and regulated by perceived exertion suggests that this group of subjects relied on sensations of speed (arising from a combination of stride length and leg motion) to subjectively adjust intensity, especially at low exercise intensities. Ekblom and Goldbarg first suggested the possible importance of feelings of strain in the working muscles as important cues in determining RPE.<sup>18</sup> They hypothesized that an individual will evaluate his perceived exertion during exercise on the basis of at least two factors: a local factor, i.e. the feeling of strain in the working muscles, and a central factor, i.e. perceived tachycardia or tachypnea. Others support the importance of this two factor theory in the regulation of effort sensations.<sup>19</sup> The present finding lends support to the involvement of local factors in rating perceived exertion and also in regulating exercise by perceived exertion.

#### RELATIONSHIP BETWEEN SPEED AND EXERCISE MODE DURING STANDARD AND SUBJECTIVELY REGULATED EXERCISE

Noble et al. have described a perceived exertion intersection point (PEIP) as the point at which walking and running are perceived to be the same.<sup>15</sup> This point is suggested to occur at approximately 6.9 km/hr (4.3 mph). In the present study the subjects were made to walk at 6.4 km/hr (4 mph) during the predetermined speed trial. The fact that 60% of the subjects changed mode when adjusting their own speed at this RPE lends support to a PEIP within the 4 mph range.

## CORRELATIONS DURING STANDARD AND SUBJECTIVELY REGULATED EXERCISE

The correlation between RPE and HR ( $r = .87$ ) during the standard treadmill exercise trial is in close agreement with previous investigators;  $r = .85$ , reported by Borg<sup>11</sup> and  $r = .87$  reported by Edwards et al.<sup>12</sup> A correlation of .86 between RPE and  $\dot{V}_E$  obtained in this study approximates the correlations of .90 obtained by Edwards et al.<sup>12</sup> The  $r$  value of .87 for RPE versus  $\dot{V}O_2$  closely approaches values of  $r = .88$  and  $r = .92$ .<sup>12,20</sup> During the subjectively regulated trials the coefficients of correlation between RPE and the above mentioned variables did not differ significantly from the coefficients during the standard exercise. These findings reflect a similar magnitude of relationship or strong association between variables when RPE is used to regulate exercise intensity.

## RELIABILITY OF SUBJECTIVE EXERCISE REGULATION

Coefficients of reliability during the subjectively regulated exercise trials are significant for speed across all RPE studied. HR and  $\dot{V}O_2$  are reliable at RPE of 10.3 and above in this group of subjects. Failure of HR to be reliable at speeds in the 5 km/hr or 3 mph (RPE 7.9) range may be due to a greater HR variability within subjects at this low exercise intensity. The previously mentioned study demonstrated that minimum change in walking speed (.25 mph) can significantly change HR.<sup>17</sup> This study has shown that a walking speed change (.40 mph) of nearly twice the minimum is not perceptually different when adjusting exercise intensity by RPE. A significant variability in HR during walking has been demonstrated even though speed adjustments were not significantly different. In essence, the lack of HR reliability during walking supports the finding of HR regression line divergence at low exercise intensities.

## ACCURACY IN ATTAINMENT OF PRESCRIBED HR DURING SUBJECTIVE EXERCISE REGULATION

The standardized exercise test ( $T_1$ ) of this experiment was designed in order to 'calibrate' each subject's perceptual response with his HR. The subjects then regulated their own exercise intensity ( $T_2$  and  $T_3$ ) at each of the 'calibration points' (RPE). In effect, exercise was prescribed at each of the subjective

RPE levels. The percent differences in HR,  $\frac{\overline{HR}_{T_{2,3}} - HR_{T_1}}{HR_{T_1}}$ , between the standard and subjectively regulated exercise trials was computed at each subject's RPE (Table 4). This percent difference is a measure of percent error in exercise HR that occurred when subjectively regulated exercise was compared to the standard exercise test of the same RPE.

An accurate determination of HR during exercise is necessary in conditioning programs that prescribe exercise intensity by HR. The post exercise 10 second pulse count has been shown to estimate exercise heart rate with a mean error of 2 to 5.7% in strenuous exercise (80-90% HR max).<sup>8,9</sup> During exercise intensities of 70% and 62% maximum HR, McArdle et al., report mean errors of 7.6 and 13.5%, respectively.<sup>8</sup> The present study demonstrates mean errors in exercise HR of 1.8 to 5.5% during running in the 78 to 92% maximum HR range (RPE 12.5 to 14.6) when subjects regulated intensity by perceived exertion alone. Mean errors of 22.3 and 34.7% were found during walking in the 49 to 73% HR max range (RPE 7.9 to 10.3). A comparison of the previously reported exercise HR errors estimated by pulse counting with the subjectively regulated HR errors computed in this study demonstrates no difference in accuracy during running at high (> 78% max) heart rates.

Table 4 also includes mean percent differences in HR across all RPE for each subject individually. The large range of intraindividual HR error across all RPE suggests that some individuals are far more accurate in subjectively regulating their exercise intensity by RPE. Although the intraindividual HR accuracy improves at higher speeds and heart rates, these individual differences must be taken into account if this method of prescription is to be used in instances where limits of prescribed target HR must be strictly enforced (i.e. cardiac rehabilitation programs).

#### CLINICAL CONSIDERATIONS

The subjective regulation of exercise intensity by perceived exertion investigated in this study shows potential for its use as a clinical method of exercise prescription in normals above intensities of 80% HR max, and 9 km/hr (5.6 mph). For individuals with restricted maximal heart rates (lower age adjusted max HR) this method may have limited value. A group of this nature would also include cardiac patients in whom accurate determinations of exercise HR may be a matter of life and death. Secondly, many patients, have relatively low symptom - limited maximal heart rates requiring their conditioning programs to consist of only low level walking. The subjective regulation of walking exercise in this investigation has been shown to be inaccurate when considering HR as the measure of exercise intensity.

ADDRESS REPRINT REQUESTS TO:

Kent B. Pandolf, Ph.D.

Military Ergonomics Division

US Army Research Institute of Environmental Medicine

Natick, MA 01760

### References

1. Åstrand PO, Caddy TE, Saltin B, Stenberg J: Cardiac output during submaximal and maximal work. *J Appl Physiol* 19:268, 1964
2. Hellerstein HK, Ader R: Relationship between percent maximal heart rate and percent maximal oxygen uptake in normals and cardiacs. *Circulation* 44:4 Suppl II:76, 1971
3. Hellerstein HK, Franklin BA: Exercise testing and prescription. Wenger NK, Hellerstein HK (eds): *Rehabilitation of the coronary patient*. New York, Wiley & Sons, 1978
4. Hoskins TA, Habasevich RA: Cardiac rehabilitation: an overview. *J Am Phy Ther Assoc* 58: 1183-1190, 1978
5. Karvonen M: Some effects of long-term exercise on health and aging. Larson A (ed): *Health and fitness in the modern world*. Chicago, Athletic Institute, 1961
6. Slater-Hammel AT, Butler LK: Accuracy in securing pulse rates by palpation. *Res Quart* 11:18-21, 1940
7. White JR: EKG changes using carotid artery for heart rate monitoring. *Med Sci Sports* 5:88-94, 1977
8. McArdle W, Zwiren L, Magel J: Validity of the post exercise heart rate as a means of estimating heart rate during work. *Res Quart* 40:523-528, 1969
9. Pollock M, Broida J, Kendrick Z: Validity of the palpation technique of heart rate determination and its estimation of training heart rate. *Res Quart* 43:77-81, 1972
10. Borg G: Perceived exertion as an indicator of somatic stress. *Scand J Rehab Med* 2:92-98, 1970

11. Borg GAV: Physical performance and perceived exertion. Lund, Gleerups, 1962
12. Edwards RHT, Melcher A, Hesser CM, Wigertz O, Ekelund LG: Physiological correlates of perceived exertion in continuous and intermittent exercise with the same average power output. *Eur J Clin Invest* 2:108-114, 1972
13. Taylor HL, Buskirk E, Henschel A: Maximal oxygen uptake as objective measure of cardio-respiratory performance. *J Appl Physiol* 8:73-80, 1955
14. Consolazio CF, Johnson RE, Pecora LJ: Physiological measurements of metabolic functions in man. New York, McGraw-Hill, 1962
15. Noble BJ, Metz KF, Pandolf KB, Bell CW, Cafarelli E, Sime WE: Perceived exertion during walking and running-II. *Med Sci Sports* 5:116-120, 1973
16. Åstrand PO, Rodahl K: Textbook of work physiology: physiological bases of exercise. New York, McGraw-Hill, 1977, p 584
17. LeBlanc JA: Use of heart rate as an index of work output. *J Appl Physiol* 10:275-280, 1957
18. Ekblom B, Goldbarg AN: The influence of physical training and other factors on the subjective rating of perceived exertion. *Acta Physiol Scand* 83:399-406, 1971
19. Pandolf KB: Influence of local and central factors in dominating rated perceived exertion during physical work. *Percept Mot Skills* 46:683-698, 1978
20. Sargeant AJ, Davies CTM: Perceived exertion during rhythmic exercise involving different muscle masses. *J Hum Ergol* 2:3-11, 1973

1. The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation.

2. Human subjects participated in these studies after giving their free and informed voluntary consent.

## FIGURE LEGENDS

Fig. 1 - The Borg category rating scale for ratings of perceived exertion (RPE).

Fig. 2 - Linear regression lines comparing speed to rated perceived exertion during  $T_1$ ,  $T_2$  and  $T_3$  ( $T_1$  = standardized exercise test;  $T_2$  = subjectively regulated exercise test; and  $T_3$  = retest of  $T_2$ ).

Fig. 3 - Linear regression lines comparing heart rate to rated perceived exertion during  $T_1$ ,  $T_2$  and  $T_3$ . Abbreviations the same as previous figure.

Table 1: Mean RPE Reported during Predetermined Speed Trials ( $T_1$ ).

---

Speed		RPE		
km/hr	mph	mean	$\pm$ SD	N
4.7	3	7.9	1.1	10
6.5	4	10.3	1.6	10
9.7	6	12.5	1.4	10
11.3	7	13.9	2.1	10
12.9	8	14.6	2.1	7

---

Table 2: Pearson Product - Moment Correlation Coefficients between RPE  
and Speed, HR,  $\dot{V}_E$ ,  $\dot{V}O_2$ .

---

	Speed	HR	$\dot{V}_E$	$\dot{V}O_2$
T <sub>1</sub>	.83	.87	.86	.87
T <sub>2</sub>	.85	.87	.90	.81
T <sub>3</sub>	.85	.85	.90	.82

---

p > .05 = non-significant (\*)

Table 3: Test - Retest Reliability Coefficients  
 at Mean RPE between T<sub>2</sub> and T<sub>3</sub>.

	RPE				
	7.9	10.3	12.5	13.9	14.6
Speed	.82	.94	.92	.74	.93
HR	.52*	.92	.91	.86	.91
$\dot{V}O_2$	.48*	.93	.92	.83	.94

p > .05 = non-significant(\*)

Table 4: Individual Percent Differences in HR between

$$T_1 \text{ and the Mean of } T_2 \text{ and } T_3 \frac{\overline{HR}_{T_{2,3}} - HR_{T_1}}{HR_{T_1}} .$$

Negative Value indicates Lower HR in  $T_{2,3}$  Compared to  $T_1$  at the Same RPE.

S	RPE					MEAN	± SD
	7.9	10.3	12.5	13.9	14.6		
1	-3.5	-2.7	-18.7	-1.2	----	-6.5	8.2
2	16.5	41.4	1.8	-0-	2.3	12.4	17.5
3	13.6	59.3	4.2	4.1	----	20.3	26.4
4	12.7	22.3	8.7	-3.6	-4.3	7.2	11.3
5	25.5	27.1	5.4	3.6	----	15.4	12.6
6	71.6	99.6	28.5	12.0	-6.8	41.0	43.7
7	13.0	13.6	-6.1	-8.2	-6.4	1.2	11.1
8	38.1	37.6	-2.4	- .3	-1.9	14.2	21.6
9	17.0	26.4	14.7	8.2	.3	13.3	10.0
10	18.8	22.4	18.6	6.5	4.4	14.4	8.1
MEAN	22.3	34.7	5.5	2.1	-1.8		
± SD	20.2	28.2	13.3	6.0	4.3		

- 6
- 7 Very, Very Light
- 8
- 9 Very Light
- 10
- 11 Fairly Light
- 12
- 13 Somewhat Hard
- 14
- 15 Hard
- 16
- 17 Very Hard
- 18
- 19 Very, Very Hard
- 20




Fig. 2 Smolov et al




FIG. 3 Smith et al.