

AD-A084 780

ARMY MEDICAL RESEARCH INST OF INFECTIOUS DISEASES FR--ETC F/6 6/5
PLASMID DETECTION IN LEGIONELLA PNEUMOPHILA AND LEGIONELLA-LIKE--ETC(U)
APR 80 P MIKESELL, J W EZZELL, G B KNUDSON

UNCLASSIFIED

ML

for
AD A
S-14700



END
DATE
FILMED
6-80
DTIC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION	3. RECIPIENT'S CATALOG NUMBER
	AD-A084 700	
4. TITLE (and Subtitle) PLASMID DETECTION IN <u>LEGIONELLA PNEUMOPHILA</u> AND <u>LEGIONELLA-LIKE</u> ORGANISMS		5. TYPE OF REPORT & PERIOD COVERED Interim
7. AUTHOR(s) PERRY MIKESSELL,* J. W. EZZELL AND GREGORY B. KNUDSON		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Army Medical Research Institute of Infectious Disease SGRD-UIB-A Fort Detrick, Frederick, MD 21701		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Medical Research and Development Command, Office of the Surgeon General Department of the Army, Washington, D.C. 20314		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS LFW USAMRIID A91C-00-133
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 30 April 1980
		13. NUMBER OF PAGES 13 pages
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Information regarding reprints not available at this time. It has been submitted for publication in <u>Infection and Immunity</u> .		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Plasmid, <u>Legionella pneumophila</u> , <u>Legionella-like</u> organisms		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Five strains of bacteria, designated <u>Legionella-like</u> organisms, and one strain of <u>Legionella pneumophila</u> were examined for the presence of extrachromosomal deoxyribonucleic acid. Cryptic plasmids were found in three of the isolates.		

ADA084700


DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

6) Plasmid Detection in Legionella pneumophila
and Legionella-like Organisms

16) PERRY MIKESELL, J. W. EZZELL AND GREGORY B. KNUDSON

United States Army Medical Research Institute of Infectious Diseases
Fort Detrick, Frederick, Maryland 21701

The views of the authors do not purport to reflect the positions of
the Department of the Army or the Department of Defense.

Dr. Mikesell's phone number: (301) 663-7341

11) 30 Apr 1988

Approved for public release; distribution unlimited

402031
80 5 22 042

J-L

Plasmid Detection in Legionella pneumophila
and Legionella-like Organisms

PERRY MIKESELL,^{*} J. W. EZZELL AND GREGORY B. KNUDSON

United States Army Medical Research Institute of Infectious Diseases
Fort Detrick, Frederick, Maryland 21701

The views of the authors do not purport to reflect the positions of
the Department of the Army or the Department of Defense.

Dr. Mikesell's phone number: (301) 663-7341

ABSTRACT

Five strains of bacteria, designated Legionella-like organisms, and one strain of Legionella pneumophila were examined for the presence of extrachromosomal deoxyribonucleic acid. Cryptic plasmids were found in three of the isolates.

Accession for	
NTIS	General
DOC	TAB
Unannounced	
Justification	
By	
Date	
App	
Dist	
A	

Approximately one third of all diagnosed cases of pneumonia in the United States each year cannot be attributed to known viral or bacterial agents (7). Recent evidence has demonstrated that certain strains of bacteria, designated Legionella-like organisms, can cause pneumonia in man (3, 14, 17). These bacteria, like many other medically important bacteria, may contain plasmid-mediated drug resistance or virulence factors (20). The demonstration of either of such factors could alter the chemotherapeutic management of patients with diseases caused by these organisms. This paper describes the isolation of plasmid deoxyribonucleic acid (DNA) from three of the six isolates tested.

All strains examined, WIGA, HEBA, TATLOCK, OLDA, PPA and TEX-KL, possess varying degrees of phenotypic or genotypic relatedness to L. pneumophila. The "Pittsburgh Pneumonia Agent" (PPA), (18) and the HEBA and TATLOCK bacteria (10) are believed to be members of a bacterial species separate from L. pneumophila (11). The OLDA bacterium, originally isolated in 1947, is now considered to be a strain of L. pneumophila, serogroup 1 (15). Two additional agents, TEX-KL and WIGA, have been proposed to represent a bacterial species, phenotypically similar but genetically unrelated to Legionella (14). Hereafter, all isolates will be referred to as Legionella-like organisms, recognizing that the OLDA isolate is a strain of L. pneumophila.

Bacteria were cultured on charcoal yeast extract agar (6) or in yeast extract broth (19) according to established parameters of growth for Legionella. Plasmids were detected utilizing agarose gel electrophoresis (16). While there are numerous modifications of this technique, most require some measure of plasmid purification prior to electrophoresis. One variation of the procedure is to lyse intact cells directly in the agarose wells (1). We have used a similar method to

screen Legionella-like bacteria for plasmids (5). One or two colonies from plates are resuspended directly in an agarose well containing 15 μ l of lysozyme mixture [lysozyme, 0.2 mg/ml; ribonuclease I, 0.1 μ g/ml; 0.05% bromphenol blue and 20% Ficoll 400,000 in Tris-borate buffer (89mM Tris base, 2.5 mM disodium EDTA and 8.9 mM boric acid, pH 8.2)]. This cell suspension was allowed to stand at room temperature for at least 15 min to assure spheroplast formation. Thirty μ l of lysis mixture [0.2% sodium dodecyl sulfate (SDS) and 10% Ficoll 400,000 in Tris-borate buffer) were added to each well, mixed gently with a toothpick and overlaid with 100 μ l of 0.2% SDS and 5% Ficoll 400,000 in same buffer and sealed with molten agarose. The definition of plasmid bands was improved if cells from broth cultures or plates were first resuspended in 20% Ficoll 400,000 in Tris-borate buffer before being added to the lysozyme mixture in the well. This observation may reflect inadequate mixing of cells in the well when samples are taken directly from solid media. Gel concentrations were varied from 0.7 to 1% agarose to enhance plasmid separation and define covalently closed circular and open circle forms of plasmid species. The samples were subjected to electrophoresis in Tris-borate buffer at 2 mA for 60 min followed by 50 mA for 100 to 180 min using a BioRad Model 220 electrophoresis cell and Model 500 power source. Following electrophoresis gels were stained with ethidium bromide (0.5 g/ml in distilled water) for 30 min followed by two 15-min rinses in distilled water. Plasmid bands were visualized using an ultraviolet transilluminator (Model C-63, Ultraviolet Products, San Gabriel, Calif).

Three of the six isolates examined, PPA, HEBA and TATLOCK, did not exhibit plasmid DNA. A single isolation or detection procedure may not be suitable for isolating plasmids from a given bacterial strain (13)

and alternate methodologies may eventually establish the presence of plasmid DNA in these bacteria. Pseudomonas aeruginosa was used as a molecular weight marker for all runs (12) (Fig. 1A). Plasmid DNA was detected in the WIGA, OLDA and TEX-KL isolates (Fig. 1B-D). These bands appear to represent separate plasmid species based on band migration patterns under different electrophoretic conditions. There also appear to be two bands of material migrating in the chromosome region (Fig. 1B-D). Resolution of these bands was obtained with the OLDA isolate by changing electrophoretic conditions (Fig. 2). Ostensibly, there are four distinct plasmids in the OLDA bacterium. A similar observation has not been made in either WIGA or TEX-KL and it is not yet known whether these two isolates also contain smaller molecular weight plasmids.

The method described above is economical, rapid and well suited to epidemiological survey work, although there are limitations. This is a screening procedure, and as such, definitive molecular and genetic characterizations of plasmid material are not possible. We believe the plasmids are large; therefore, estimates of plasmid molecular weight are not possible using this electrophoretic system (9). Attempts to purify the plasmids from these bacteria by established methods have been unsuccessful (2, 4, 8). The plasmids may be highly susceptible to shear by virtue of their size or they may exhibit some degree of nonintegrated association with the host chromosome, both factors complicating purification. It should also be emphasized that the cell concentration can greatly affect the detection of plasmid material particularly in the 12 to 19 megadalton range. We apply 15-, 30- and 40- μ l amounts of sample material to the gel to preclude masking of plasmids which migrate near the chromosomal DNA. This is particularly advantageous if optimal growth conditions have not been defined for a given isolate.

We have established the presence of extrachromosomal genetic elements in two Legionella-like organisms, WIGA and TEX-KL, and in the OLDA strain of L. pneumophila. The results indicate that Legionella organisms are able to maintain plasmid elements as part of their genetic composition. The ability of these organisms to exchange genetic material with other bacteria has, to our knowledge, not been reported. However in view of the narrow spectrum of antibiotics effective in the treatment of legionellosis and Legionella-like diseases, the acquisition of drug resistance factors by the bacteria could have serious clinical ramifications.

ACKNOWLEDGMENT

We wish to thank G. A. Jacoby for supplying us with Pseudomonas aeruginosa, PU21. We are also indebted to Earnest L. Cook and Helen Hargett for their excellent technical assistance.

LITERATURE CITED

1. Barnes, W. M. 1977. Plasmid detection and sizing in single colony lysates. *Science* 195:393-394.
2. Clewell, D. B., and D. R. Helinski. 1969. Supercoiled circular DNA-protein complex in Escherichia coli: purification and induced conversion to an open circular DNA form. *Proc. Natl. Acad. Sci. U.S.A.* 62:1159-1166.
3. Cordes, L. G., H. W. Wilkinson, G. W. Gorman, B. J. Fikes, and D. W. Fraser. 1979. Atypical Legionella-like organisms: fastidious water-associated bacteria pathogenic for man. *Lancet* 2:927-930.
4. Currier, T. C., and E. W. Nester. 1976. Isolation of covalently closed circular DNA of high molecular weight from bacteria. *Anal. Biochem.* 76:431-441.
5. Eckhardt, T. 1978. A rapid method for the identification of plasmid desoxyribonucleic acid in bacteria. *Plasmid* 1:584-588.
6. Feeley, J. C., R. J. Gibson, G. W. Gorman, N. C. Langford, J. K. Rasheed, D. C. Mackel, and W. B. Baine. 1979. Charcoal-yeast agar: primary isolation medium for Legionella pneumophila. *J. Clin. Microbiol.* 10:437-441.
7. Fraser, D. W., and J. E. McDade. 1979. Legionellosis. *Scient. Am.* 241(4):82-99.
8. Guerry, P., D. J. LeBlanc, and S. Falkow. 1973. General method for the isolation of plasmid deoxyribonucleic acid. *J. Bacteriol.* 116:1064-1066.
9. Hansen, J. B., and R. H. Olsen. 1978. Isolation of large bacterial plasmids and characterization of the P2 histocompatibility group plasmids pMG1 and pMG5. *J. Bacteriol.* 135:227-238.

10. Hebert, G. A., C. W. Moss, L. K. McDougal, F. M. Bozeman, R. M. McKinney, and D. J. Brenner. 1980. The rickettsia-like organisms TATLOCK (1943) and HEBA (1959): bacteria phenotypically similar to but genetically distinct from Legionella pneumophila and the WIGA bacterium. Ann. Intern. Med. 92:45-52.
11. Hebert, G. A., B. M. Thomason, P. P. Harris, M. D. Hicklin, and R. M. McKinney. 1980. "Pittsburgh Pneumonia Agent": a bacterium phenotypically similar to Legionella pneumophila and identical to the TATLOCK bacterium. Ann. Intern. Med. 92:53-54.
12. Jacoby, G. A. 1977. Classification of plasmids in Pseudomonas aeruginosa, p. 119-126. In D. Schlesinger (ed.), Microbiology - 1977. American Society for Microbiology, Washington, D.C.
13. LeBlanc, D. J., and L. N. Lee. 1979. Rapid screening procedure for detection of plasmids in streptococci. J. Bacteriol. 140:1112-1115.
14. Lewallen, K. R., R. M. McKinney, D. J. Brenner, C. W. Moss, D. H. Dail, B. M. Thomason, and R. A. Bright. 1979. A newly identified bacterium phenotypically resembling, but genetically distinct from Legionella pneumophila: an isolate in a case of pneumonia. Ann. Intern. Med. 91:831-834.
15. McDade, J. E., D. J. Brenner, and F. M. Bozeman. 1979. Legionnaires' disease bacterium isolated in 1947. Ann. Intern. Med. 90:659-661.
16. Meyers, J. A., D. Sanchez, L. P. Elwell, and S. Falkow. 1976. Simple agarose gel electrophoretic method for the identification and characterization of plasmid deoxyribonucleic acid. J. Bacteriol. 127:1529-1537.

17. Myerowitz, R. L., A. W. Pasculle, J. N. Dowling, G. J. Pazin, M. Puerzer, R. B. Yee, C. R. Rinaldo, Jr., and T. R. Hakala. 1979. Opportunistic lung infection due to "Pittsburgh Pneumonia Agent." N. Engl. J. Med. 301:953-958.
18. Pasculle, A. W., R. L. Myerowitz, and C. R. Rinaldo, Jr. 1979. New bacterial agent of pneumonia isolated from renal-transplant recipients. Lancet 2:58-61.
19. Ristroph, J. D., K. W. Hedlund, and R. G. Allen. Liquid medium for growth of Legionella pneumophila. J. Clin. Microbiol. 11:19-21.
20. Shipley, P. L., W. S. Dallas, G. Dougan, and S. Falkow. 1979. Expression of plasmid genes in pathogenic bacteria, p. 176-180. In D. Schlessinger (ed.) Microbiology - 1979. American Society for Microbiology, Washington, D.C.

FIGURE LEGENDS

FIG. 1. 1.0% agarose gel of plasmid DNA. Gels were run at 2 mA for 60 min followed by 50 mA for 150 min. Migration is from top to bottom. (A) P. aeruginosa, PU21 (12) (plasmid molecular weight 312 megadaltons). Smaller 20-megadalton plasmid is not evident. (B) TEX-KL, two distinct plasmid bands representing separate molecular species. (C) OLDA a single plasmid band is evident migrating in the same region as top band of TEX-KL. (D) WIGA, very faint band (arrow) migrates in the same region as lower band of TEX-KL. The large diffuse band in all wells is chromosome. Note double band appearance in chromosome region (B, C, and D).

FIG. 2. 0.7% agarose gel run of OLDA. Electrophoresis was carried out at 2 mA for 60 min followed by 50 mA for 105 min. There appear to be four separate plasmid bands. The two bands immediately below and the single band immediately above the chromosomal band are masked by the chromosomal DNA shown in Fig. 1C.

