

AD-A084 985

NAVAL RESEARCH LAB WASHINGTON DC
NUMERICAL EXPERIMENTS ON THE STRATOSPHERIC-TROPOSPHERIC DYNAMIC--ETC(U)
APR 80 W C CHAO, M R SCHOEBERL, D F STROBEL
NRL-MR-4209

F/6 4/2

SBIE -AD-E000 419


NL

UNCLASSIFIED

[OF]
AE 6
7-2-88



END
DATE
FILMED
6-80
DTIC



MICROCOPY RESOLUTION TEST CHART

NATIONAL BUREAU OF STANDARDS-1963-A

(12.5)

LEVEL III

AD-E000 419

NRL Memorandum Report 4209

Numerical Experiments on the Stratospheric-Tropospheric Dynamical Interaction

WINSTON C. CHAO

*Science Applications, Inc.
McLean, Virginia 22102*

AND

MARK R. SCHOEBERL AND DARRELL F. STROBEL

*Geophysical and Plasma Dynamics Branch
Plasma Physics Division*

April 11, 1980

DTIC
ELECTE
JUN 4 1980
S B D



NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

80 4 17 00 5

ADA 084985

DOC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 4209	2. GOVT ACCESSION NO. AD-A084985	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NUMERICAL EXPERIMENTS ON THE STRATOSPHERIC- TROPOSPHERIC DYNAMICAL INTERACTION		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem.
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Winston C. Chao*, Mark R. Schoeberl and Darrell F. Strobel		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D.C. 20375		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62759N; 9F52-551-792; 9F52-551; 67-0885-0-0
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Environmental Prediction Research Facility Washington, D.C. 20361		12. REPORT DATE April 11, 1980
		13. NUMBER OF PAGES 20
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES *Present address: Science Applications, Inc., McLean, Virginia 22102. This work was sponsored by the Naval Environmental Prediction Research Facility under Project No. ZF52551001, work unit title NRL-1 Stratospheric Global Prediction Model.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Stratospheric-tropospheric interaction		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Numerical experiments with the UCLA-NEPRF tropospheric weather forecast model were performed to test the influence of stratospheric conditions which are representative of sudden warmings on tropospheric weather. Stratospheric conditions were represented by imposed vertical velocities at the upper boundary. An externally forced stratospheric disturbance as large as a sudden warming was found to generate an observable response in sea level pressure with a time constant of the order of a day. The numerical results are interpreted with the aid of an analytical model based on the linearized shallow water equation.		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

i
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

PRECEDING PAGE BLANK-NOT FILMED

CONTENTS

I. INTRODUCTION	1
II. NUMERICAL EXPERIMENTS	3
III. NUMERICAL RESULTS	5
IV. ANALYTIC STUDY OF THE ATMOSPHERIC RESPONSE TO THE IMPOSED VERTICAL VELOCITY	6
V. DISCUSSION	10
ACKNOWLEDGMENTS	11
REFERENCES	12

DTIC
ELECTE
S JUN 4 1980 D
B

ACCESSION for		
NTIS	White Section	☒
DDC	Buff Section	☐
UNANNOUNCED		☐
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	A/AIL. and/or	SPECIAL
A		-

NUMERICAL EXPERIMENTS ON THE STRATOSPHERIC-TROPOSPHERIC DYNAMICAL INTERACTION

I. INTRODUCTION

Most tropospheric weather prediction models impose an upper boundary condition of $\omega = \frac{dp}{dt} = 0$ at some finite height in the lower stratosphere. Lindzen et al. (1968) pointed out that this "rigid lid" boundary condition produces spurious reflections and introduces oscillations and standing waves whose amplitudes can mask the true solutions. In a more recent study Kirkwood and Derome (1977) investigated the effect of this boundary condition on forced stationary planetary waves. They used realistic mean zonal winds in their quasi-geostrophic study rather than a quiescent, isothermal atmosphere as Lindzen et al. (1968) adopted. The reflection and transmission of these ultralong waves are strongly dependent on the structure of the polar night jet centered near the stratopause (Charney and Drazin, 1961; Schoeberl and Geller, 1977). Kirkwood and Derome found that the stationary planetary wave structure could only be accurately modeled when the vertical resolution in the stratosphere adequately represented the mean zonal winds and Newtonian cooling (≥ 21 levels or $\Delta p \leq 10$ mb). Insufficient stratospheric resolution resulted in a poorly defined polar night jet and Newtonian cooling profile and spurious reflection at the upper boundary. They also suggested that misrepresentation of the structure of stationary planetary waves can lead to the generation of spurious external, free waves of which the 5 day wave (Geisler and Dickinson, 1976) has been found to be dominant in some forecast models (Lambert and Merilees, 1978).

The movement of tropospheric weather systems at middle and high latitudes is greatly influenced by these stationary, ultralong

Manuscript submitted February 6, 1980.

waves, particularly wave numbers 1 and 2. Therefore a good forecast of the former requires a good forecast of the latter. These planetary scale eddies are formed in the troposphere through baroclinic, diabatic, and orographic processes. During winter these waves propagate vertically, perturb the stratospheric circulation, transport heat and momentum from mid-latitudes into the polar regions, and generate sudden stratospheric warmings. Thus it is imperative to incorporate a sufficiently resolved stratosphere for tropospheric weather prediction.

In this report we document an effort to determine the influence of stratospheric conditions which are representative of a sudden warming on tropospheric weather. Quiroz (1977), McGuirk (1978), and O'Neill and Taylor (1979) showed that the 1976/77 sudden warming definitely affected tropospheric weather patterns. In addition Rosmond (private communications, 1978) has noted that Navy forecasts have lower skill during sudden warmings. The goals of this study were to ascertain the importance of stratospheric-tropospheric dynamical coupling and the time scale for stratospheric processes to affect tropospheric weather.

II. NUMERICAL EXPERIMENTS

The UCLA atmospheric general circulation model (Arakawa and Lamb, 1977) has been developed as a forecast model at the Naval Environmental Prediction Research Facility (NEPRF) (Payne, 1979) and was used for the numerical experiments discussed below. Two forecasts of 5-days duration each were performed with the same initialized data for January 19, 1977, generated from NEPRF's global model initialization program. During that week, record cold temperatures were experienced over most of the continental United States and an intense sudden warming had developed in the stratosphere. The reference forecast had the usual $\omega = 0$ upper boundary condition at 50 mb. The other forecast was performed with an imposed vertical velocity, ω_I , at 50 mb based on the observed geopotential heights of wave number 1 and 2 during that intense sudden stratospheric warming.

The vertical velocity boundary condition has two wave components, $\omega_I = \omega_I^1 + \omega_I^2$, where the superscripts represent wavenumber, and are defined as:

$$\omega_I^1 = Ag(\theta)f_1(\phi)$$

$$\omega_I^2 = Ag(\theta)f_2(\phi),$$

Here θ is the latitude and ϕ is the longitude, $g(\theta)$ is defined to be zero between equator and 30°N and to be a sine curve between 30°N and 90°N and is symmetric with respect to the equator. Also

$$f_1(\phi) = \cos(\phi - \phi_1)$$

$$f_2(\phi) = \cos [(\phi - \phi_2)/2]$$

where ϕ_1 corresponds to 120°W and ϕ_2 to 0° . A is the observed amplitude and equal to -4.1×10^{-5} mb/sec, or -3.54 mb/day. These values are crude estimates of the kind of vertical velocities which may be associated with sudden warming events. The values were obtained from the mechanistic model of Schoeberl and Strobel (1980). ϕ_1 and ϕ_2 are chosen such that ω_1^1 and ω_1^2 are 180° out of phase with their respective components of the observed geopotential height to insure that the vertical propagation of wave energy is upward. Momentum and heat are also transported through the model top by the imposed vertical velocity.

III. NUMERICAL RESULTS

From the detailed results of these two integrations, we have selected three grid points to illustrate the salient features. The general weather pattern is shown in Fig. 1 which is day 5 of the integration with imposed upper boundary vertical velocity. Point A is the location of the maximum negative value of ω_I ($=6 \text{ mb day}^{-1}$) at the upper boundary, whereas point C is the maximum positive value of ω_I ($=7.7 \text{ mb day}^{-1}$). Point B, which coincides with a deep low in the Gulf of Alaska, has $\omega_I \sim 0$ (actually -0.5 mb day^{-1}). To emphasize stratospheric influence deep in the troposphere, sea level pressure differences from 1000 mb for the two experiments are shown in Fig. 2 to 4 for a 5 day period. Perceptible differences are evident after only 1 day at points A and C. At point A, after 4 days, the pressure difference between the two runs reaches 3 mb. At point B, a large change between day 3 and 4 reflects the movements and deepening of the low the pressure difference. A maximum difference of $\sim 6 \text{ mb}$ is reached after 3.5 days at point C.

IV. ANALYTIC STUDY OF THE ATMOSPHERIC RESPONSE TO THE IMPOSED VERTICAL VELOCITY.

Before interpreting the meaning of these results, it is instructive to investigate with an analytic model the expected planetary scale response of the atmosphere to the imposed vertical velocity. We hypothesize that the essential features of the response can be modeled with the linearized shallow water equations with a mean depth, H , equal to the equivalent depth of an isothermal atmosphere (~ 10 km):

$$\frac{\partial \vec{V}}{\partial t} + f \vec{k} \times \vec{V} + \nabla \phi' = 0, \quad (1)$$

$$\frac{\partial \phi'}{\partial t} + g H \nabla \cdot \vec{V} = Q, \quad (2)$$

where $\phi' = gh' = g(h-H)$, h is the height of the water surface and is analogous to the sea level pressure, $\vec{V}$ is the horizontal velocity and is constant with height, and (Q/g) is the mass source and represents the effect of imposed vertical velocity, ω_I .

From (1) we obtain the divergence and vorticity equations:

$$\frac{\partial}{\partial t} \nabla \cdot \vec{V} - f \vec{k} \cdot \nabla \times \vec{V} + \nabla^2 \phi' = 0, \quad (3)$$

$$\frac{\partial}{\partial t} \vec{k} \cdot \nabla \times \vec{V} + f \nabla \cdot \vec{V} = 0. \quad (4)$$

Substitution of (4) and (2) into $\frac{\partial}{\partial t}$ (3) gives

$$\frac{\partial^2}{\partial t^2} \nabla \cdot \vec{V} + f^2 \nabla \cdot \vec{V} - gH \nabla^2 (\nabla \cdot \vec{V}) + \nabla^2 Q = 0. \quad (5)$$

If Q has wave numbers k and l in x and y directions, respectively, i.e.,

$$Q = \hat{Q} \exp [i(kx + ly)],$$

the Laplacian can be written as

$$\nabla^2 = -(k^2 + l^2) \equiv -K^2.$$

If we use the notation

$$D \equiv \nabla \cdot \vec{V} = \hat{D} e^{i(kx+ly)}, \text{ Eq. (5) becomes} \quad (6)$$

$$\frac{\partial^2}{\partial t^2} \hat{D} + (gHK^2 + f^2) \hat{D} = K^2 \hat{Q}.$$

For the initial conditions we choose $\vec{V} = \vec{\phi}' = 0$ at $t = 0$, which are consistent with the data initialization for the numerical experiments; thus,

$$D = 0 \text{ (since } \vec{V} = 0 \text{ at } t = 0),$$

$$\frac{\partial D}{\partial t} = 0 \text{ (from (3))}. \quad (7)$$

The solution is

$$\hat{D} = A \cos(\nu t) - A, \quad (8)$$

where

$$A = \frac{-K^2 \hat{Q}}{\nu^2},$$

and

$$\nu = (gHK^2 + f^2)^{1/2}.$$

Eq. (2) can now be written

$$\frac{\partial \hat{\phi}'}{\partial t} + gH[A \cos(\nu t) - A] = \hat{Q},$$

$$\hat{\phi}' = 0 \text{ at } t = 0.$$

The solution for ϕ' is

$$\begin{aligned} \phi' &= \left[\frac{-gHA}{\nu} \sin(\nu t) + (gHA + \hat{Q})t \right] e^{i(kx+ly)} \\ &= \left[\frac{gHK^2 \hat{Q}}{3\nu} \sin(\nu t) + \frac{f^2}{2\nu} \hat{Q}t \right] e^{i(kx+ly)} \end{aligned} \quad (9)$$

Since the surface height variation is analogous to the sea level pressure variation in the atmospheric model and Q represents the imposed vertical velocity at model top, Eq. (9) suggests the sea level pressure has a standing oscillation with frequency, ν , and a linear variation with time, which vanishes as $f = 0$. The standing oscillation is a direct consequence of the $D = 0$ initial condition and the imposed vertical velocity ($Q \neq 0$). If we choose $\hat{D} = -A$ instead for the initial condition along with $\frac{\partial \hat{D}}{\partial t} = 0$, then Eq. (8) is just $\hat{D} = -A$ and the standing oscillation term in Eq. (9) vanishes. Thus the mismatch of initial and upper boundary conditions generates the oscillatory term which corresponds to the creation of two free modes traveling in opposite directions.

A numerical estimate for ϕ' is given below:

$$g = 9.8 \text{ m/s}^2, H = 10^4 \text{ m}, f = 2\Omega \sin \theta = 1.3 \times 10^{-4} \text{ s}^{-1},$$

$$\Omega = 7.3 \times 10^{-5} \text{ s}^{-1}, \theta = \frac{\pi}{3}, a = 6.4 \times 10^6 \text{ m, earth's radius,}$$

$$k = \frac{2\pi}{L_x}, \ell = \frac{2\pi}{L_y}, L_x = 2\pi a \cos \theta = \pi a, L_y = \frac{2\pi a}{3},$$

Thus ,

$$k = \frac{2}{a}, \ell = \frac{3}{a},$$

$$K^2 \equiv k^2 + \ell^2 = \frac{13}{a^2} = 3.2 \times 10^{-13} \text{ m}^{-2},$$

$$\nu = (gHK^2 + f^2)^{1/2} = 2.2 \times 10^{-4} \text{ s}^{-1},$$

$$\frac{gHK^2}{\nu} = 3.1 \times 10^3 \text{ s}, \quad \frac{f^2}{\nu} = 0.32, T \equiv \frac{2\pi}{\nu} = 2.9 \times 10^4 \text{ s},$$

$$\hat{Q} = \max |\omega_I| = 4.1 \times 10^{-5} \text{ mb s}^{-1} \text{ for each wave number.}$$

Thus the amplitude of the standing oscillation is 0.17 mb and its period is 8 hr. The growth rate of the second term in Eq. (9) is 1.2 mb day⁻¹.

Fig. 5 shows the sea level pressure difference at three locations in the first one and half days between the two runs, $P_u - P_r$, where P_u is the sea level pressure of the run with imposed upper boundary vertical velocity, and P_r , that of the reference run. Since $P_u - P_r$ is analogous to the ϕ' in the shallow water equation, the time variation of $P_u - P_r$, shown in the figure as an oscillation superimposed on a linear change, corresponds well with that predicted by the shallow water model.

V. DISCUSSION

The study shows that the difference between the two simulations can be qualitatively explained by Eq. (9). In the shallow water equation case, the oscillatory part of the solution in Eq. (9) can be eliminated by using a different initial condition, but not the linear term.

However in a model with topographic forcing and differential heating due to sea-land contrast the imposed vertical velocity can be initialized as part of the stationary waves and then both terms in Eq. (9) are eliminated. In future proposed experiments the imposed vertical velocity upper boundary condition will be initialized as part of the stationary waves because it is kept constant in time. It is anticipated that the difference in sea level pressure between such a proposed experiment and the reference experiment will not be dominated by the direct effect of the imposed vertical velocity and will probably be smaller in amplitude than the results presented in Fig. 2-4. However the present experiment does demonstrate that an externally generated stratospheric disturbance as large as a sudden warming can generate an observable response in sea level pressure with a time constant of the order of a day.

Acknowledgments

This work was supported by the NAVAIR block fund under subtask number 9F52-551-792.

REFERENCES

- Arakawa, A. and V. R. Lamb, 1977: Computational design of the basic dynamical processes of the UCLA general circulation model, Methods in Computational Physics, Vol. 17, Academic Press, New York, pp. 174-207.
- Charney, J. G. and P. G. Drazin, 1961: Propagation of planetary scale disturbances from the lower into the upper atmosphere. J. Geophys. Res., 66, 83-109.
- Geisler, J. E. and R. E. Dickinson, 1976: The five-day wave on a sphere with realistic zonal winds. J. Atmos. Sci., 33, 63-641.
- Kirkwood, E. and J. Derome, 1977: Some effects of the upper boundary condition and vertical resolution on modeling forced stationary planetary waves. Mon. Wea. Rev. 105, 1239-1251.
- Lambert, S. J. and P. E. Merilees, 1978: A study of planetary wave errors in a spectral numerical weather prediction model. Atmosphere-Ocean, 16, 197-211.
- Lindzen, R. S., E. S. Batten and J. W. Kim, 1968: Oscillations in atmospheres with tops. Mon. Wea. Rev., 96, 133-140.
- McGuirk, J. P., 1978: Planetary scale forcing of January 1977 weather. Science, 199, 293-295.
- O'Neill, A. and B. F. Taylor, 1979: A study of the major stratospheric warming of 1976/77. Quart. J. R. Met. Soc., 105, pp. 71-92.

- Payne, S. W., 1979: A verification case study of the NEPRF/UCLA global forecast system, Fourth Conf. on Numerical Weather Prediction, AMS, Silver Spring, Md. Oct. 29-Nov. 1, 1979, 155-160.
- Quiroz, R. S.: 1977, The tropospheric-stratospheric polar vortex breakdown of January 1977. Geophys. Res. Lett., 4, 151-154.
- Schoeberl, M. R. and M. A. Geller, 1977: A calculation of the structure of stationary planetary waves in winter. J. Atmos. Sci., 34, 1235-1255.
- Schoeberl, M. R. and D. F. Strobel: 1978a, The zonally averaged circulation of the middle atmosphere. J. Atmos. Sci., 35, 577-591.




Fig. 1 - General weather pattern on day 5




Fig. 2 - Sea level pressure - 1000 mb at point A




Fig. 3 - Sea level pressure - 1000 mb at point B




Fig. 4 - Sea level pressure - 1000 mb at point C




Fig. 5 - Detailed sea level pressure difference between the two runs at point A, B, and C for the first 1.5 days