

AD-A085 440

NAVAL OCEANOGRAPHIC OFFICE-NSTL STATION MS

F/S 8/3

SURFACE CURRENTS, SOUTHEAST NORTH PACIFIC OCEAN INCLUDING THE W--ETC (11)

AUG 77


UNCLASSIFIED

NOO-SP-1402-NP-13

NL

1 OF 1
AL-000000

END
DATE
FILMED
7-80
DTIC




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

✓ NAVAL OCEANOGRAPHIC OFFICE SPECIAL PUBLICATION T402-NP-13

SURFACE CURRENTS.

(12) LEVEL II

SOUTHEAST NORTH PACIFIC OCEAN INCLUDING THE
WEST COAST OF CENTRAL AMERICA



AUGUST 1977

REFRESHED 1978

DTIC

SELECTED

JAN 15 1980



THIS PUBLICATION IS THE PROPERTY OF THE
NAVY DEPARTMENT AND IS LOANED TO YOUR
OFFICE. IT IS TO BE RETURNED TO THE
NAVY DEPARTMENT ON REQUEST.

ABSTRACT

THIS ATLAS, AND THE SERIES OF WHICH IT IS A PART, IS COMPUTER GENERATED AND AUTOMATICALLY PLOTTED. IT MAKES AVAILABLE TO THE USER THE MOST RECENT SURFACE CURRENT DATA COLLECTED AND WILL BE UPDATED WHENEVER SUFFICIENT AMOUNTS OF DATA ARE ADDED TO THE DATA FILE. THIS AND THE OTHER ATLASES ARE BASED ON A VAST QUANTITY OF DATA AS COMPARED TO THE PREVIOUS MANUALLY-COMPILED EDITIONS PRINTED IN THE MID-THIRTIES.

THE SURFACE CURRENT INFORMATION IS BASED MAINLY ON SHIP DRIFT, WHICH IS THE DIFFERENCE BETWEEN THE DEAD RECKONING POSITION AND THE POSITION DETERMINED BY ANY TYPE OF NAVIGATIONAL FIX. THIS DIFFERENCE DESCRIBES THE DIRECTION AND SPEED OF THE CURRENT.

DISCLAIMER NOTICE

**THIS DOCUMENT IS BEST QUALITY
PRACTICABLE. THE COPY FURNISHED
TO DTIC CONTAINED A SIGNIFICANT
NUMBER OF PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

(14) N

(6)

SURFACE CURRENTS.

SOUTHEAST NORTH PACIFIC OCEAN INCLUDING
WEST COAST OF CENTRAL AMERICA.

(9) Final rept.



(12)

(11) AUG [REDACTED] 77
Reprinted 1979

APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.

NAVAL OCEANOGRAPHIC OFFICE
NSTL STATION, MISSISSIPPI 39522

1

250450

14 NOO-SP-1402-NP-13

FACE CURRENTS.

RTH PACIFIC OCEAN INCLUDING THE
OAST OF CENTRAL AMERICA.



12 50

11 AUG 1977
Reprinted 1979

FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED

VAL OCEANOGRAPHIC OFFICE
L STATION, MISSISSIPPI 39522

250450

2
LW

ACKNOWLEDGMENTS


**Messrs. Raymond J. Beauchesne* and William E. Boisvert
made major contributions to this atlas.**

***Mr. Beauchesne presently is employed by the Bureau of
Naval Personnel.**

FOREWORD

THIS ATLAS, ONE IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLAS, IS PRODUCED TO FULFILL A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND INDUSTRIAL COMMUNITIES FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. THESE ATLASES ADD TO THE WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERATIONAL PLANNING, NAVIGATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID PRODUCTION AND WIDE DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE LATEST COMPUTER TECHNIQUES.

THE CONSTANT IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA RECEIVED OVER THE YEARS IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH OBSERVATIONS OF VOLUNTARY OBSERVERS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE OCEANOGRAPHIC OFFICE, AND THE USER OF THE ATLASES RELY ON THE PERSONAL OBSERVATIONS OF THE MAN WHO HAS "BEEN THERE." MARINERS, IN REPORTING THEIR OBSERVATIONS, RENDER A SERVICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL WHO GO DOWN TO THE SEA IN SHIPS." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC NAVIGATION AIDS, AND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED ENVIRONMENTAL AND NAVIGATIONAL INFORMATION HAS BECOME INCREASINGLY IMP


JOHN R. McDONNELL
Captain, U.S. Navy
Commander

FOREWORD

IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES, A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. A WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE ATLAS.

IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE USER OF THE ATLASES RELY ON THE PERSONAL OBSERVATION HAS "BEEN THERE." MARINERS, IN REPORTING THEIR SERVICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO USE THEM." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC SURVEILLANCE SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED NAUTICAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.

John R. McDonnell
JOHN R. McDONNELL
Captain, U.S. Navy
Commander

ACCESSION for		
NTIS	White Section	☒
DDC	Buff Section	☐
UNANNOUNCED		☐
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist. A/RIL and/or SPECIAL		
A	23	20-


2

SURFACE CURRENT ATLASES

THIS SERIES OF COMPUTERIZED ATLASES REPLACES THE OLD HYDROGRAPHIC OFFICE ATLASES OF SURFACE CURRENTS (HOP 566, 568, 569, 570) WHICH WERE MANUALLY COMPILED FROM DATA OBTAINED DURING THE PERIOD 1903 - 1934. THESE NEW ATLASES CONFORM TO THE STANDARD NAVY OCEAN AREA AND REGION INDEX LIMITS SHOWN BELOW: e.g., NOO SP 1402-NP 10 COVERS NORTH PACIFIC REGION 10 EAST OF THE PHILIPPINES.

THESE ATLASES
AREAS AS THE NORTH
CURRENTS ARE STRONGLY
PREDICTABLE HOURLY

RECENT IMPROVEMENTS IN THE DATA FILE ASSURE THE INCLUSION OF THE LATEST, HIGH QUALITY SURFACE CURRENT DATA AVAILABLE. THE FILE NOW CONTAINS MORE THAN 4,200,000 OBSERVATIONS AND A GENERAL UPDATE OF THE FILE WILL BE MADE



Introduction

The Surface Current Data File, from which these atlases are derived, consists primarily of over four million ship set and drift observations. These data were collected by the Netherlands, Japan, Britain, France, and the United States. The file is supplemented by several thousand Geomagnetic Electrokinetograph (GEK) observations, mostly Japanese. The file spans the period from the early 1850's to the present. The earliest observations were collected by the Netherlands and Great Britain; those of the 1960's through the present are primarily United States data.


General Quality

The quality of this data file is considered high for this type of derived value. The data have been carefully screened for duplication; observations taken under adverse conditions (i.e. high winds and waves, time between observations greater than 12 hours) have been eliminated when warranted. Consideration was given to the reliability of the observer; doubtful shipboard computations of set and drift were edited; and observations with erroneous locations (mostly observations on land) have been eliminated. The accepted data are considered most useful when used collectively as in summaries where a number of observations show trends.

General Observation Technique

The set (direction) and drift (speed) are computed by the navigator from the difference between the dead reckoning (DR) position and the position determined by any type of navigational fix. The drift can be determined along any straight line track and includes all factors which cause changes in the DR position. When a fix is obtained, the current set (direction) is FROM the DR position TO the fix; the drift (speed) is equal to the distance in nautical miles between the DR and the fix, divided by the number of hours since the last fix. For successive observations, the TO POSITION of one observation becomes the FROM POSITION of the next observation.


Because the influence of current may vary along a ship's track, the MEAN POSITION of the track is assigned as the geographic location of the current observation. An example of a current computation is shown in the figure below.




EXAMPLE OF A SURFACE CURRENT (SHIP'S DRIFT) OBSERVATION

Data Presentation


The following legend shows two types of surface current presentations by 1° quadrangle, type 1 with 12 or more observations and type 2 with fewer than 12 observations. Where there are 11 or fewer observations within a 1° quadrangle, the total number of observations is shown within the 90° quadrant containing the observations.




If there are 12 or more obs.
 by vector resultants as fol:






- (1) Persistent Current - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.



- (4) Bizonal Flow - Practically all obs are concentrated in opposite pairs sectors, and one pair contains at 80 percent as many observations as pair. This generally indicates a that occurs in zones of entrainment opposing currents (see examples A quadrangles 1, 2, and 3).



If there are 12 or more observations in a 1° quadrangle, the surface current is depicted by vector resultants as follows:





(1) Persistent Current - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.

(2) Prevailing Current - 70 percent or more of all observations fall within two adjacent 45° sectors.


(3) Primary Current with Secondary Direction
(a) Primary Current - 50 percent or more of all observations fall within three adjacent 45° sectors.


(b) Secondary Direction - 20 percent or more of all observations fall within a 45° sector, and the two resultant vector directions are separated by more than 90° of arc.




(4) Bizonal Flow - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).


(5) Variable Current - The 45° sector with most observations has less than 25 percent of all observations; direction is indeterminate.





JANUARY





100 W

95 W

90 W

85 W

80 W

75 W

29 N

25 N

20 N

15 N

10 N

5 N

0

100 W

95 W

90 W


85 W


80 W

75 W


FEBRUARY

2






2




APRIL




APRIL


2




MAY




2





JUNE




2












2




SEPTEMBER





2







2







2











AL-JAN THROUGH DEC

2



WINTER-JAN, FEB, MAR



100 W 95 W
WINTER-JAN, FEB, MAR


90 W


85 W


80 W

75 W


A







SUMMER-JUL, AUG, SEP



2

120 W 115 W 110 W 105 W 100 W 95 W

35 N

30 N


20 N

15 N

10 N


5 N

0



120 W 115 W 110 W 105 W 100 W 95 W

AUTUMN-OCT, NOV, DEC



AUTUMN-OCT, NOV, DEC

2

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER N00 SP1402-NP13	2. GOVT ACCESSION NO. ADA085440	3. RECIPIENT'S CATALOG NUMBER	
4. TITLE (and Subtitle) Surface Currents Southeast North Pacific Ocean including the West Coast of Central America		5. TYPE OF REPORT & PERIOD COVERED Final	
6. AUTHOR(s) Naval Oceanographic Office		7. PERFORMING ORG. REPORT NUMBER	
8. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Naval Oceanographic Office NSTL Station, MS 39522		9. CONTRACT OR GRANT NUMBER(s)	
10. CONTROLLING OFFICE NAME AND ADDRESS Naval Oceanographic Office		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. REPORT DATE August 1977	
		14. NUMBER OF PAGES 24	
		15. SECURITY CLASS. (of this report)	
		16. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.			
18. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
19. SUPPLEMENTARY NOTES			
20. KEY WORDS (Continue on reverse side if necessary and identify by block number) Surface currents, Southeast North Pacific Ocean, Central America			
21. ABSTRACT (Continue on reverse side if necessary and identify by block number) This atlas, and the series of which it is a part, is computer generated and automatically plotted. It makes available to the user the most recent surface current data collected and will be updated whenever sufficient amounts of data are added to the data file. This and the other atlases are based on a vast quantity of data as compared to the previous manually compiled editions printed in the mid-thirties.			

FORM 1473 EDITION OF 1 NOV 65 IS OBSOLETE
1 JAN 73 S/N 0102-014-6001

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. The surface current information is based mainly on ship drift, which is the difference between the dead reckoning position and the position determined by any type of navigational fix. This difference describes the direction and speed of the current.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DISTRIBUTION LIST

NAVY

CINCPACFLT (02M)
COMTHIRDFLT
COMSEVENTHFLT
COMSUBPAC
COMNAVAIRPAC
COMPATWINGSPAC
PATWINGSPAC DET ADAK
PATWING 1
COMNAVSURFPAC
DIRNAVOCEANMET
FLENUMWEACEN
FLEWEACEN GUAM
FLEWEACEN PEARL
NAVWEASERVFAC SAN DIEGO
NAVWEASERVFAC YOKOSUKA
NWS D ASHEVILLE
NWS D ADAK
NWS D AGANA
NWS D ATSUGI
NWS D KADENA
NWS D MISAWA

OTHER GOVT.

NOAA/NODC
NOAA/NCC

PRIVATE

AMAX EXPLORATION INC.

FOREIGN

HYDROGRAPHER/R.A.N.
DEPT. TRANSPORTATION/AUSTRALIA

MED
8