

AD-A085 955

TEXAS A AND M UNIV COLLEGE STATION DEPT OF CHEMISTRY F/G 7/4
THE NUCLEAR MAGNETIC RESONANCE SPECTRA OF BISPHthalocyaninatOLA--ETC(U)
JUN 80 M TSUTSUI, K KASUGA N00014-75-C-0417

UNCLASSIFIED


TR-36

NL

[or]
AD
20

END
DATE
FILMED

8-80
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 085955

LEVEL II

9

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 36	2. GOVT ACCESSION NO. AD-A085955	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The Nuclear Magnetic Resonance Spectra of Bisphthalocyaninatolanthanide(III).		5. TYPE OF REPORT & PERIOD COVERED (9) Interim rept.
7. AUTHOR(s) M./Tsutsui and K./Kasuga		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Texas A&M University College Station, Texas 77843		8. CONTRACT OR GRANT NUMBER(s) NO 0014-75-C-0417 N00014-75-C-0417
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-559
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1980
		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release; Distribution Unlimited (14) TR-367		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) (13) N00014-75-C-0417		
18. SUPPLEMENTARY NOTES Submitted for publication in J. Coord. Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) bisphthalocyaninatolanthanide(III) complexes, NMR studies of phthalocyanine compounds		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The NMR spectra of bisphthalocyaninatolanthanide(III) complexes (La, Nd, Sm, and Eu) have been examined. An effect of a macrocyclic-ring current and induced shift caused by lanthanide(III) ions were discussed.		

DTIC
ELECTE
S JUN 25 1980 D
E


DDC FILE COPY

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 68 IS OBSOLETE S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

80 6 23 054

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

The Nuclear Magnetic Resonance Spectra of Bisphthalocyaninatolanthanide(III)

K. Kasuga and M. Tsutsui*

Department of Chemistry, Texas A&M University, College Station, TX 77843

(Received , 1980)

The NMR spectra of bisphthalocyaninatolanthanide(III) complexes (La, Nd, Sm, and Eu) have been examined. An effect of a macrocyclic-ring current and an induced shift caused by lanthanide(III) ions were discussed.

INTRODUCTION

NMR studies of phthalocyanine compounds have been difficult for a long time because of their poor solubilities.¹ Recently, using Fourier transform techniques, Marks et al. reported some NMR results of lithium, zinc, and uranyl phthalocyanines.² Lanthanide complexes such as acetylacetonatolanthanide(III) and its derivatives are also known to be useful for NMR shift reagents.³ We report here NMR results of bisphthalocyaninatolanthanide (La(III), Nd(III), Sm(III), and Eu(III)) to study the effects of a macrocyclic ring current of phthalocyanine ligands and induced shifts caused by lanthanide(III) ions on peaks of benzene ring protons of the complexes.

EXPERIMENTAL

Samples used for NMR measurements were prepared by the methods of Kirin et al. and Misumi et al.⁴ The purity of samples was verified by elemental

Accession For	
NTIS GRA&I	☒
DDC TAB	☐
Library Codes	
Dist	Avail and/or special
A	

analyses and visible spectra. A deuteriodimethylsulfoxide (DMSO-d_6) solution of the samples was refluxed for ca. 30 minutes under argon gas, and the filtrate was transferred into a NMR tube under argon atmosphere. A Varian FT-80 spectrometer was used to obtain spectra at 60°C .

RESULTS AND DISCUSSION

Since the synthesis of bisphthalocyaninatolanthanide(III) complexes was published,⁴ many studies on their properties have been reported.⁵ For instance, the complexes show blue in polar solvents such as methanol, N,N-dimethylformamide, and dimethylsulfoxide, while they show green in dichloromethane and chloroform.⁴ A few ideas have been reported to explain the color difference.^{5c,d,e} Previously, we reported that green bisphthalocyaninatoneodymium(II) recrystallized from dichloromethane is a radical species, $\text{PcNdPc}^\cdot$ which is generated by abstracting imine hydrogen of a blue Pc_2NdH complex (Pc = phthalocyanine dianion).⁶ Furthermore, a molecular structure of the green bisphthalocyaninatoneodymium(III) complex was analyzed; a neodymium(III) ion occupies a central position between parallel, but staggered phthalocyanine ligands.⁷ An X-ray photoelectron spectrum was also obtained to verify the location of acidic hydrogen-imine hydrogen in bisphthalocyaninatoneodymium(III).⁷ The data imply that eight central nitrogen atoms are chemically equivalent to each other, and thereby acidic hydrogen does not bind strongly to any of the nitrogen atom in the complex, consistent with Kirin's model which has a weak intermolecular hydrogen bond, $\text{PcNdPc} \begin{smallmatrix} \text{H} \\ \text{H} \end{smallmatrix} \text{PcNdPc}$.^{5e} We studied a FT-NMR of bisphthalocyaninatolanthanide(III) to verify the behavior of acidic proton of the complex, but it could not be detected. Figure 1

Figure 1

shows the FT-NMR spectrum of bisphthalocyaninatone odymium(III). On the basis of works of Lever and Marks, the lower-field multiplet is assigned to α proton of a benzene ring and the higher field one is assigned to β proton.^{1,2} FT-NMR results of bisphthalocyaninatolanthanide(III) (La, Nd, Sm, and Eu) and lithium and zinc phthalocyanine complexes are shown in Table 1.

Table 1

Comparing of sandwich-type bisphthalocyaninatolanthanum(III) complex with 1:1 type metallophthalocyanine complexes, it can be shown that the position of a β proton signal in the lanthanum(III) complex is almost the same as those of normal-type complexes, while the position of α proton of the lanthanum(III) complex shifts to the higher magnetic field (Table 1). This shift is the result of shielding of α protons belonging to one macrocycle by the benzene ring current associated with the other macrocycle. Since the two multiplets have the same general outline as in the 1:1 type compound, it is apparent that both α protons of the benzene ring are affected by equal shielding, consistent with the fact of the staggered arrangement of the two phthalocyanine ligands. The induced shifts of α and β protons of the complexes increase in the order Nd<Eu<<Sm (Figure 2).

Figure 2

It is interesting that the induced shift of bisphthalocyaninatosamarium(II) is much larger than those of acetylacetonatosamarium(III) and its derivatives.³ Further studies on this point may be performed.

ADKNOWLEDGEMENTS

We are grateful for support of this work by the Office of Naval Research (Grant No. 3210) and the Robert A. Welch Foundation (Grant No. A420).

REFERENCES

1. A. B. P. Lever, Advances in Inorganic Chemistry and Radiochemistry, 7, 27 (1965).
2. T. J. Marks and D. R. Stojakovic, J. Am. Chem. Soc., 100, 1695 (1978).
3. A. F. Cockerill, G. L. O. Davies, R. C. Harden, and D. M. Rackham, Chemical Reviews, 73, 553 (1973).
4. a) I. S. Kirin, P. N. Moskalev, and Yu. A. Makashev, Russ. J. Inorg. Chem., 10, 1065 (1965). b) I. S. Kirin, R. N. Moskalev, and N. V. Ivannikova, Russ. J. Inorg. Chem., 12, 497 (1967). c) S. Misumi and K. Kasuga, Nippon Kagaku Zasshi, 92, 335 (1971).
5. a) I. S. Kirin, P. N. Moskalev, and Yu. A. Makashev, Russ. J. Inorg. Chem., 12, 369 (1967). b) I. S. Kirin, Yu. L. Shelekhin, and P. N. Moskalev, Soviet Physics - Solid State, 9, 1461 (1967). c) P. N. Moskalev and I. S. Kirin, Russ. J. Phys. Chem., 46, 1019 (1972). d) A. G. Mackay, J. F. Boas, and G. J. Troup, Aust. J. Chem., 27, 955 (1974). e) P. N. Moskalev, G. N. Shapkin, and A. N. Daronskikh, Russ. J. Inorg. Chem., 24, 188 (1979).
6. K. Kasuga and M. Tsutsui, J. Am. Chem. Soc., (submitted).
7. K. Kasuga, M. Tsutsui, R. G. Pettersen, K. Tatsumi, N. V. Opdenbosh, G. Pepe, and E. F. Meyer, Jr., J. Am. Chem. Soc., (in press).

TABLE 1

¹H NMR Data for Bisphthalocyaninatolanthanide(III) and Lithium and Zinc Phthalocyanine Complexes*,**

Compound	¹ H NMR	
	α proton	β proton
Pc ₂ LaH	8.75 (16H)	8.10 (16H)
Pc ₂ NdH	6.90 (16H)	6.30 (16H)
Pc ₂ SmH	18.67 (16H)	18.34 (16H)
Pc ₂ EuH	10.84 (16H)	9.03 (16H)
PcLi ₂ ***, 2	9.38 (8H)	8.04 (8H)
PcZn****, 2	9.50 (8H)	7.94 (8H)

* FT data in δ (ppm). ** All data in DMSO_{d6} at 60 °C. *** C₆D₆ added 2% DMSO_{d6}. **** C₆D₆. All peaks show multiplets.

Figure Captions

Figure 1. NMR spectrum of bisphthalocyaninatoneodymium(III) in DMSO_{d6} at 60°C .

Figure 2. Induced chemical shifts of α and β protons. $\bigcirc$, α proton;
 β proton.




Figure 1. NMR spectrum of bisphthalocyaninatoneodymium(III) in DMSO-d_6 at 60°C .




FIGURE 2. Induced chemical shifts of α and β protons. o, α proton; β proton.

