

AD-A086 178

NAVAL OCEANOGRAPHIC OFFICE NSTL STATION NS
SURFACE CURRENTS NORTHWEST NORTH ATLANTIC OCEAN AND LABRADOR SE--ETC(U)
JAN 78

F/8 8/3

UNCLASSIFIED

N00-SP-1400-NA-4

NL

REF ID

A086178



END
DATE
FILMED
8-80
DTIC


52

ADA 086178

NAVAL OCEANOGRAPHIC OFFICE SPECIAL P

SURFACE CURRENTS

NORTHWEST NORTH ATLANTIC OCEAN AND LABRADOR SEA



JANUARY 1978
REPRINTED 1980


NO. 1000-1000-1000-1000
NO. 1000-1000-1000-1000
NO. 1000-1000-1000-1000

NAVAL OCEANOGRAPHIC OFFICE SPECIAL PUBLICATION 1400—NA 4

SURFACE CURRENTS

12 LEVEL II

NORTHWEST NORTH ATLANTIC OCEAN AND LABRADOR SEA



JANUARY 1973
REVISED 1981

DTIC
ELECTE
S D

ABSTRACT

THIS ATLAS, AND THE SERIES OF WHICH IT IS A PART, IS COMPUTER GENERATED AND AUTOMATICALLY PLOTTED. IT MAKES AVAILABLE TO THE USER THE MOST RECENT SURFACE CURRENT DATA COLLECTED AND WILL BE UPDATED WHENEVER SUFFICIENT AMOUNTS OF DATA ARE ADDED TO THE DATA FILE. THIS AND THE OTHER ATLASES ARE BASED ON A VAST QUANTITY OF DATA AS COMPARED TO THE PREVIOUS MANUALLY-COMPILED EDITIONS PRINTED IN THE MID-1970S.

THE SURFACE CURRENT INFORMATION IS BASED MAINLY ON DRIFT NETS, WHICH IS THE DIFFERENCE BETWEEN THE DRIFT NECKLINE POSITION AND THE POSITION DETERMINED BY ADRIFT BY NAVIGATIONAL FIX. THIS DIFFERENCE DETERMINES THE DIRECTION AND SPEED OF THE CURRENT.

14

NOO-SP-1400-NA-4

SURFACE CURRENTS

NORTHWEST NORTH ATLANTIC OCEAN AND LABRADOR SEA



12 21

11

JAN ~~1978~~ 1978

REPRINTED 1980

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

NAVAL OCEANOGRAPHIC OFFICE
NSTL STATION, MISSISSIPPI 39522

250450

7/6

ACKNOWLEDGMENTS

Messrs. Raymond J. Beauchesne* and William E. Boisvert made major contributions to this atlas.

*Mr. Beauchesne presently is employed by the Bureau of Naval Personnel.

FOREWORD

↓

THIS ATLAS, ONE IN A SERIES OF 43 REGIONAL SURFACE CURRENT ATLASES, IS PRODUCED TO FULFILL A NEED OF NAVY PLANNING STAFFS AND THE SCIENTIFIC AND INDUSTRIAL COMMUNITIES FOR THE LATEST AVAILABLE OCEAN SURFACE CURRENT DATA. THESE ATLASES ADD TO THE WEALTH OF NAUTICAL INFORMATION UPON WHICH OPERATIONAL PLANNING, NAVIGATIONAL SAFETY, AND SHIPPING ECONOMY DEPEND. RAPID PRODUCTION AND WIDE DISSEMINATION OF THIS ATLAS ARE MADE POSSIBLE BY THE LATEST COMPUTER TECHNIQUES.

↑

THE CONSTANT IMPROVEMENT IN THE QUALITY OF SURFACE CURRENT DATA RECEIVED OVER THE YEARS IS MADE POSSIBLE LARGELY BY THE MORE THOROUGH REPORTS OF VOLUNTARY OBSERVERS IN RECENT YEARS. THE DEFENSE MAPPING AGENCY, THE OCEANOGRAPHIC OFFICE, AND THE USER OF THE ATLASES RELY ON THE PERSONAL OBSERVATIONS OF THE MAN WHO HAS "BEEN THERE." MARINERS, IN REPORTING THEIR OBSERVATIONS, RENDER A SERVICE NOT ONLY TO THEMSELVES BUT ALSO TO ALL "WHO GO DOWN TO THE SEA IN SHIPS." WITH THE ADVENT OF NUCLEAR POWER, ELECTRONIC NAVIGATION AIDS, AND 300,000-TON SHIPS, UP-TO-DATE, RAPIDLY DISSEMINATED ENVIRONMENTAL AND NAVIGATIONAL INFORMATION HAS BECOME INCREASINGLY IMPORTANT.

John R. McDonnell
 JOHN R. McDONNELL
 Captain, U.S. Navy
 Commander

iii

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED <input type="checkbox"/>		
JUSTIFICATION _____		
BY _____		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL and/or	SPECIAL
A	23	P

SURFACE CURRENT ATLASES


THIS SERIES OF COMPUTERIZED ATLASES REPLACES THE OLD HYDROGRAPHIC OFFICE ATLASES OF SURFACE CURRENTS (HOP 566, 568, 569, 570) WHICH WERE MANUALLY COMPILED FROM DATA OBTAINED DURING THE PERIOD 1903 - 1934. THESE NEW ATLASES CONFORM TO THE STANDARD NAVY OCEAN AREA AND REGION INDEX LIMITS SHOWN BELOW: e.g., NOO SP 1402-NP 10 COVERS NORTH PACIFIC REGION 10 EAST OF THE PHILIPPINES.

AS AMOUNTS OF NEW DATA AVAILABLE.

THESE GRAPHICS MAY NOT COVER ALL OCEANIC AREAS AS THE NORTH SEA, PERSIAN GULF, AND BALTIC SEA CURRENTS ARE STRONGLY TIDAL AND SUBJECT TO PREDICTABLE HOURLY CHANGES OF

RECENT IMPROVEMENTS IN THE DATA FILE ASSURE THE INCLUSION OF THE LATEST, HIGH QUALITY SURFACE CURRENT DATA AVAILABLE. THE FILE NOW CONTAINS MORE THAN 4,200,000 OBSERVATIONS AND A GENERAL UPDATE OF THE FILE WILL BE MADE

INDEX



SURFACE CURRENT ATLASES


PHIC OFFICE
 QUALITY
 NEW ATLASES
 OWN BELOW:
 PHILIPPINES.

AS AMOUNTS OF NEW DATA WARRANT, MOST LIKELY EVERY 12 - 18 MONTHS.

THESE GRAPHICS MAY NOT BE TRULY REPRESENTATIVE OF THE ACTUAL FLOW IN SUCH AREAS AS THE NORTH SEA, PERSIAN GULF, GULF OF THAILAND, AND YELLOW SEA WHERE CURRENTS ARE STRONGLY TIDAL. FOR SUCH AREAS, OTHER SOURCES DESCRIBING PREDICTABLE HOURLY CHANGES OF TIDAL CURRENTS SHOULD BE CONSULTED.

IF LATEST,
 MORE
 MADE

INDEX



Introduction

The Surface Current Data File, from which these atlases are derived, consists primarily of over four million ship set and drift observations. These data were collected by the Netherlands, Japan, Britain, France, and the United States. The file is supplemented by several thousand Geomagnetic Electrokinetograph (GEK) observations, mostly Japanese. The file spans the period from the early 1850's to the present. The earliest observations were collected by the Netherlands and Great Britain; those of the 1960's through the present are primarily United States data.


General Quality

The quality of this data file is considered high for this type of derived value. The data have been carefully screened for duplication; observations taken under adverse conditions (i.e. high winds and waves, time between observations greater than 12 hours) have been eliminated when warranted. Consideration was given to the reliability of the observer; doubtful shipboard computations of set and drift were edited; and observations with erroneous locations (mostly observations on land) have been eliminated. The accepted data are considered most useful when used collectively as in summaries where a number of observations show trends.

General Observation Technique

The set (direction) and drift (speed) are computed by the navigator from the difference between the dead reckoning (DR) position and the position determined by any type of navigational fix. The drift can be determined along any straight line track and includes all factors which cause changes in the DR position. When a fix is obtained, the current set (direction) is FROM the DR position TO the fix; the drift (speed) is equal to the distance in nautical miles between the DR and the fix, divided by the number of hours since the last fix. For successive observations, the TO POSITION of one observation becomes the FROM POSITION of the next observation.


Because the influence of current may vary along a ship's track, the MEAN POSITION of the track is assigned as the geographic location of the current observation. An example of a current computation is shown in the figure below.




EXAMPLE OF A SURFACE CURRENT (SHIP'S DRIFT) OBSERVATION

Data Presentation


The following legend shows two types of surface current presentations by 1° quadrangle, type 1 with 12 or more observations and type 2 with fewer than 12 observations. Where there are 11 or fewer observations within a 1° quadrangle, the total number of observations is shown within the 90° quadrant containing the observations.




If there are 12 or more observations in a 1° quadrangle, the vector resultant is shown as follows:






- (1) Persistent Current - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.
- (2) Prevailing Current - all observations fall within 45° sectors.



- (4) Bisonal Flow - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).



If there are 12 or more observations in a 1° quadrangle, the surface current is depicted by vector resultants as follows:





- (1) Persistent Current - 60 percent or more of all observations fall within a 45° sector of the 8-point compass.
- (2) Prevailing Current - 70 percent or more of all observations fall within two adjacent 45° sectors.
- (3) Primary Current with Secondary Direction -
 - (a) Primary Current - 50 percent or more of all observations fall within three adjacent 45° sectors.
 - (b) Secondary Direction - 20 percent or more of all observations fall within a 45° sector, and the two resultant vector directions are separated by more than 90° of arc.

consists primarily of observations collected by the Japanese, mostly Japanese. The earliest observations are from the 1960's through 1965. The value is under adverse conditions (more than 12 hours) and is not reliable. The observations have been edited and observations have been eliminated from summaries.

from the difference between any type of track and includes the current is equal to the number of hours of observation.

POSITION of observation. An example


2 KN
RED.
N
085
1°12'
1°06'N
1°00'
N



- (4) Bizonal Flow - Practically all observations are concentrated in opposite pairs of 45° sectors, and one pair contains at least 80 percent as many observations as the opposite pair. This generally indicates variability that occurs in zones of entrainment between opposing currents (see examples A and B, quadrangles 1, 2, and 3).
- (5) Variable Current - The 45° sector with most observations has less than 25 percent of all observations; direction is indeterminate.


by 1° quadrangle, observations. Where number of observa-

N TOTAL OBSERVATIONS.
D USED FOR 1:1000 OR
LL OBSERVATIONS.
(*T) FOR ALL OBSERVATIONS.
IMARY DIRECTION, 238
BY DIRECTION, 0.6 KNOT
ADRAWN.



1

2



WINTER-JAN, FEB, MAR

1

55 W

50 W

45 W

40 W

35 W

30 W

64 N

60 N

55 N

50 N

49 N

55 W

50 W

45 W

40 W

35 W


30 W


WINTER-JAN, FEB, MAR

1

1

2





1 SPRING-APR, MAY, JUN 1

55 W

50 W

45 W

40 W

35 W

30 W

64 N

60 N

55 N

50 N

49 N

55 W

50 W

45 W

40 W

35 W


30 W


1 SPRING-APR, MAY, JUN

1

1

2






SUMMER-JUL, AUG, SEP

1

55 W 50 W 45 W 40 W 35 W 30 W

64 N



60 N

55 N

50 N

49 N


55 W 50 W 45 W 40 W 35 W 30 W

SUMMER-JUL, AUG, SEP

1

1

2



AUTUMN-OCT, NOV, DEC

1

55 W

50 W

45 W

40 W

35 W

30 W

64 N

60 N

55 N

50 N

49 N

55 W

50 W

45 W

40 W

35 W


30 W

AUTUMN-OCT, NOV, DEC

1


1

2



ANNUAL-JAN THROUGH DEC

1



ANNUAL-JAN THROUGH DEC

1

1

2