



AD-A087 252

VIRGINIA UNIV CHARLOTTESVILLE DEPT OF CHEMISTRY F/G 7/2
TRANSITION METAL-PROMOTED OXIDATIVE FUSION WITH CARBORANE AND B--ETC(U)
JUL 80 N S HOSMANE, R N GRIMES N00014-75-C-0305
UNCLASSIFIED TR-33 NL

1 of 1
AD
A087252



END
DATE
FILMED
9-80
DTIC



134
LEVEL

OFFICE OF NAVAL RESEARCH

15
CONTRACT NO. N00014-75-C-0305

9
TECHNICAL REPORT NO. 33

12
6
Transition Metal-Promoted Oxidative Fusion With
Carborane and Borane Anions. Synthesis of $C_nB_n^{n-}$ ($n = 8-11$)
Carborane Systems in the Presence of $B_5H_8^{n-}$ or $B_3H_8^{n-}$ Ions.

10
Narayan S. Hosmane and Russell N. Grimes

Department of Chemistry
University of Virginia
Charlottesville, Va. 22901

11) Jul 1980
14) TR-33
Prepared for Publication in

Inorganic Chemistry

July, 1980

Reproduction in whole or in part is permitted
for any purpose of the United States Government

Approved for public release; distribution unlimited

ADA 087252

DDC FILE COPY.

462144

80

7

25

186

Transition Metal-Promoted Oxidative Fusion With
 Carborane and Borane Anions. Synthesis of C_4B_n ($n = 8-11$)
 Carborane Systems in the Presence of $B_5H_8^-$ or $B_3H_8^-$ Ions

Narayan S. Hosmane and Russell N. Grimes*

Department of Chemistry
 University of Virginia
 Charlottesville, VA 22901

Abstract. Simultaneous treatment of $CoCl_2$ or $FeCl_2$ with the $(2,3-(CH_3)_2C_2B_4H_5)^-$ and $(B_5H_8)^-$ ions in THF at room temperature gave mixed-ligand metallocarborane complexes. With $CoCl_2$ the major isolable product was $[(CH_3)_2C_2B_4H_4]CoH[(CH_3)_2C_2B_7H_7]$, a yellow air-stable crystalline solid. The reaction with $FeCl_2$ gave several products including the known carborane $(CH_3)_4C_4B_8H_8$ and the new species $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$ and $(CH_3)_4C_4B_{11}H_{11}$; the last of these is believed to form by oxidation of an unstable precursor complex, $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_7H_7]$, which was observed mass spectroscopically. The red $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$ complex is moderately air-stable, but slowly reacted with O_2 to form $(CH_3)_4C_4B_9H_{11}$. From electron-counting arguments the new carboranes $(CH_3)_4C_4B_{11}H_{11}$ and $(CH_3)_4C_4B_9H_{11}$ are proposed to have 15-vertex nido and 13-vertex arachno cage geometries, respectively, but linked-cage structures are not ruled out. In the iron-borane-carborane reaction the order of addition of reagents is important; thus, treatment of $FeCl_2$ initially with $(B_5H_8)^-$ followed later by $((CH_3)_2C_2B_4H_5)^-$ gave species identified mass spectroscopically as $((CH_3)_2C_2B_5H_5)_2FeH_2$ and $(CH_3)_4C_4B_{10}H_{10}$, neither of which was observed in the previously described 'simultaneous'


reaction. Surprisingly, the reaction of FeCl_2 , $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$, and B_3H_8^- gave products and yields almost identical to those obtained in the corresponding $\text{FeCl}_2/(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-/\text{B}_5\text{H}_8^-$ reaction. However, the $\text{FeCl}_2/(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-/\text{BH}_4^-$ reaction gave a virtually quantitative yield of the known complex $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{FeH}_2$. Oxidative fusion of $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ ions to $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ via complexation with iron was found to occur in the absence of O_2 or other external oxidants, with metallic iron formed as a byproduct. A sequence of events is proposed to account for the observations in the iron-borane-carborane reactions.

Accession For	
NTIS GRA&I	☒
DDC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

Transition metal-facilitated dimerizations, cyclizations, and polymerizations of hydrocarbons are well known and constitute a growing area of interest in organic and organometallic synthetic chemistry.¹ One might anticipate that roughly analogous processes should be possible in transition metal-polyhedral borane systems, and indeed, the "oxidative fusion" reactions² that are exhibited by certain types of metal-carborane complexes provide real cases in point. In oxidative fusion, two open-cage (e.g., pyramidal) ligands are face-coordinated to a transition metal atom which is subsequently eliminated, the ligands merging face-to-face to form a single large polyhedral system (Figure 1). Thus far, the most clearly defined examples involve the stable, isolable bis(carboranyl) complexes^{3,4} $[2,3-(CH_3)_2C_2B_4H_4]_2Co^{III}H$ and $[2,3-(CH_3)_2C_2B_4H_4]_2Fe^{II}H_2$; each of these undergoes nearly quantitative air-oxidation at room temperature, forming $(CH_3)_4C_4B_8H_8$, a 12-vertex distorted icosahedron⁵ which is formally a $(2n + 4)$ -electron nido⁶ system. These complexes and the carborane product have been fully characterized by spectroscopic and X-ray methods. Other carborane ligands, e.g., $C_2B_4H_7^-$ and $CH_3C_2B_4H_6^-$, can also be oxidatively fused,⁷ and other transition metal ions such as Ni^{2+} can be employed, but in these cases the metal sandwich intermediates have not been isolated.

Oxidative fusion has also been observed⁸ with small metallacarboranes such as $(\eta^5-C_5H_5)CoC_2B_3H_7$, which undergoes dimerization to several $(\eta^5-C_5H_5)_2C_4B_6H_{10}$ isomers belonging to different structural classes; thus far, two of these fused complexes have been crystallographically characterized.^{8,9} As part of a long-range study of the scope

of the oxidative fusion process, we have attempted to prepare mixed-ligand borane-carborane metal complexes, whose ligands might then combine to produce large carboranes, e.g.





If successful, reactions of this type could open the way to low-energy designed syntheses of a variety of large cage systems, perhaps including polyhedra larger than any presently known.

In the study reported herein, our attempts at mixed-ligand complex formation and fusion led not to the expected products but to new four-carbon carboranes which we believe to be the first examples of supra-icosahedral carboranes without metal atoms.

Results and Discussion

Reaction of $B_5H_8^-$, $(CH_3)_2C_2B_4H_5^-$ and $CoCl_2$. Previous work in our laboratory had established the following reactions:^{4,10,11}





In the present investigation, we attempted to coordinate $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4^{2-}$ and a borane anion ligand to a common cobalt ion and thereby form a mixed-ligand borane-carborane complex, via the reaction of CoCl_2 with $\text{Na}^+\text{B}_5\text{H}_8^-$ and $\text{Na}^+(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ simultaneously. However, the major isolable product was a mixed-ligand bis(carboranyl) species, $[2,3-(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{CoH}[2,3-(\text{CH}_3)_2\text{C}_2\text{B}_7\text{H}_7]$, I, obtained in 21% yield as a yellow air-stable solid, m.p. 79° , and purified by sublimation at 30°C onto a cold finger. The mass spectrum of this complex exhibits a cutoff at m/e 300 corresponding to the parent ion, and the profile of intensities in the parent region is consistent with the formula given above; a high-resolution mass measurement (Experimental Section) confirmed the elemental composition.

The 32.1-MHz ^{11}B FT NMR spectrum of I (Table I) is not particularly informative, but the range of chemical shifts is compatible with the ^{11}B spectra of the related cobaltacarborane complexes $(6,9\text{-C}_2\text{B}_7\text{H}_9)_2\text{Co}^-$ (formerly numbered 6,7)¹² and $[2,3-(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_6]_2\text{CoH}$.⁴ The 100-MHz ^1H FT NMR spectrum (Table II) contains two methyl peaks of equal area, indicating that each of the carborane ligands, viewed separately, has equivalent C-CH₃ groups and hence a local mirror plane bisecting the framework C-C bond. From the mild conditions of synthesis it can be assumed that the cage carbon atoms in each ligand remain adjacent and are in the metal-bonded face as shown in Figure 2. The Figure as

drawn suggests the presence of an overall molecular mirror plane, but this cannot be confirmed from the NMR data since magnetic interaction between the carborane ligands may be so weak that different rotamers cannot be distinguished.¹³ The "extra" proton bound to cobalt exhibits a broad peak at δ -12.3 ppm [rel. to $(\text{CH}_3)_4\text{Si}$] in the region typical of metal-hydrogen resonances.

The proposed 6,9- $(\text{CH}_3)_2\text{C}_2\text{B}_7\text{H}_7^{2-}$ ligand in I is a C,C'-dimethyl derivative of the 6,9- $\text{C}_2\text{B}_7\text{H}_9^{2-}$ ("dicarbazapide") ligand found in several previously reported metallacarboranes,¹² and the 2,3- $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4^{2-}$ moiety in I occurs in numerous complexes also.^{3,4,10,14} However, mixed-ligand complexes involving small carborane ligands have not previously been reported. Since I appeared unreactive toward oxygen and failed to undergo oxidative fusion (in contrast to $[\text{2,3-}(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{CoH}$),⁴ we turned our attention to the $\text{B}_5\text{H}_8^{--}(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^{--}\text{FeCl}_2$ reaction system.

Reaction of $\text{B}_5\text{H}_8^{--}$, $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^{--}$, and FeCl_2 (Simultaneous). This system proved much more complex than the corresponding cobalt reaction and was found to be highly sensitive to experimental details, particularly the order of addition of reagents.²⁶ Treatment of an equimolar THF solution of the $\text{B}_5\text{H}_8^{--}$ and $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^{--}$ ions with FeCl_2 at room temperature generated H_2 , B_2H_6 , and the volatile complex $\text{BH}_3\cdot\text{THF}$, the solution turning dark purple. Removal of the solvent and workup of the residue in carbon tetrachloride (primarily to liberate monomeric species from the resin-like matrix) gave several products which were sublimed together onto a cold finger and subsequently separated by

high-pressure liquid chromatography (HPLC). In addition to the known³ carborane $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ (64% yield based on starting carborane), the major isolated products were red $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{FeH}_2[(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5]$ (II), m.p. 71°C , 11% yield, and colorless $(\text{CH}_3)_4\text{C}_4\text{B}_{11}\text{H}_{11}$ (III), m.p. 68°C , obtained in 2.3% yield. In addition, mass spectra of the sublimed product mixture revealed the presence of $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{FeH}_2-[(\text{CH}_3)_2\text{C}_2\text{B}_7\text{H}_7]$ IV, the probable precursor of III via oxidative fusion; complex IV proved extremely air-sensitive (in curious contrast to its cobalt analogue I, described above) and was not isolated.

Very small amounts of white crystalline carboranes, identified as $[(\text{CH}_3)_2\text{C}_2\text{B}_8\text{H}_8]_2$ and $[(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_7]_2$ from their mass spectra, were recovered in the HPLC separation of the products but could not be further characterized. Spectroscopic data on III, however, were obtained, as described in a later section.

Figure 3 depicts the proposed structures of II and IV. The mass spectrum of II contains a cutoff at m/e 274 corresponding to the parent ion (^{56}Fe isotope), as well as an intense peak group having a local cutoff at m/e 262, indicating facile loss of BH in the mass spectrometer to form the known³ complex $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{FeH}_2$. The proton-decoupled ^{11}B NMR spectrum (Table I) contains four identifiable resonances whose range of chemical shifts is consistent with the presence of $(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5^{2-}$ and $2,3-(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4^{2-}$ ligands¹⁵ and with the structure postulated in Figure 3, but which cannot be unequivocally assigned. The ^1H spectrum (Table II) contains only two methyl peaks, demonstrating that each of the ligands per se contains equivalent C- CH_3 units related by a mirror plane; as in the case of the cobalt

species I, discussed above, the NMR data do not establish the orientation of the ligands relative to each other. Also present in the ^1H spectrum is a broad resonance at δ -12.4 ppm arising from the FeH_2 protons.

The formal $(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5^{2-}$ ligand in II has not previously been observed, the only other examples of $\text{C}_2\text{B}_5\text{M}$ -type metallocarboranes being 3,1,7- $(\eta^5\text{-C}_5\text{H}_5)\text{CoC}_2\text{B}_5\text{H}_7$,¹⁷ and 3,1,7- $(\text{CO})_3\text{FeC}_2\text{B}_5\text{H}_7$,¹⁸ in both of which the cage carbon atoms have been assigned nonadjacent positions in the cage. From electron-counting considerations,⁶ $(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5^{2-}$ is expected to have a nido 7-vertex structure so that the metal atom completes a closo 8-vertex polyhedron, as shown in Figure 3.

Reaction of B_3H_8^- , 2,3- $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$, and FeCl_2 . When the preceding experiment was repeated with $(\text{CH}_3)_4\text{N}^+\text{B}_3\text{H}_8^-$ in place of $\text{Na}^+\text{B}_5\text{H}_8^-$, the isolated products were again II, III, and $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$, and in nearly identical yields. These findings indicate that both B_3H_8^- and B_5H_8^- are capable of attacking the $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ group in the presence of FeCl_2 ¹⁹ with net addition of three BH units to generate the unstable $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{FeH}_2[(\text{CH}_3)_2\text{C}_2\text{B}_7\text{H}_7]$ complex (IV), which in turn degrades to form the stable, isolable species $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{FeH}_2[(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5]$ (II). In principle, the replacement of B_5H_8^- by B_3H_8^- as a borane reagent is highly advantageous (if the ultimate products are the same) because of the accessibility and ease of handling of B_3H_8^- salts, which are air-stable and can be easily prepared²⁰ from NaBH_4 or obtained commercially. The present work is

evidently the first instance in which $B_3H_8^-$ has been employed as a source of boron in polyhedral metallaborane or metallocarborane syntheses, although we have recently reported the multi-step, one-pot preparation of the nido-carborane 2,3- $(CH_3)_2C_2B_4H_6$ from $(C_2H_5)_3N^+B_3H_8^-$ via B_5H_9 generated in situ.²¹ Many metal chelate (non-polyhedral) complexes of $B_3H_8^-$ are, of course, known.²²

Reaction of BH_4^- , 2,3- $(CH_3)_2C_2B_4H_5^-$, and $FeCl_2$. As an extension of the work involving $B_3H_8^-$ and $B_5H_8^-$, $FeCl_2$ was treated with a THF solution containing a mixture of BH_4^- and $(CH_3)_2C_2B_4H_5^-$ ions. The sole carborane product was the known dark red complex³ [2,3- $(CH_3)_2C_2B_4H_4$] $_2FeH_2$, obtained in 97% yield. Clearly no incorporation of boron from BH_4^- took place, but the nearly quantitative yield of ferracarborane was significantly higher than we have previously obtained³ in the absence of BH_4^- (~80 to 90%). The probable explanation is that trace oxidants present in the THF, which survive drying and distillation and reduce the yield of ferracarborane complex, are removed by the BH_4^- (which does not attack the ferracarborane itself). However, the role of BH_4^- in this reaction was not further explored.

Reaction of $B_5H_8^-$ and $FeCl_2$ Followed by $(CH_3)_2C_2B_4H_5^-$. The preceding experiments did not establish whether the observed metallocarborane products formed via attack of the carborane ligand on a ferraborane species produced initially, attack of $B_5H_8^-$ or $B_3H_8^-$ on a metallocarborane intermediate, or both. In order to investigate the actual sequence, $FeCl_2$ and a 2:1 excess of $Na^+B_5H_8^-$ in THF were

allowed to react, forming a brown nonvolatile complex which was not isolated. After treatment of this dark solution with excess $\text{Na}^+(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ followed by workup with CCl_4 as described above, the sublimable products were found on mass spectroscopic analysis to consist of a complex mixture of species including II, $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$, and $[(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5]_2\text{FeH}_2$ (V), a product which was not observed to form in the "simultaneous" $\text{B}_5\text{H}_8^-/(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-/\text{FeCl}_2$ reaction described above. Complex V unfortunately did not survive column chromatography on silica, even when conducted in an oxygen- and water-free chamber, and was not isolated. The observation that V forms in this reaction, but is absent when both B_5H_8^- and $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ are present initially, suggests that FeCl_2 reacts more rapidly with the carborane anion than with B_5H_8^- ; this question will be discussed further in the concluding section.

Mass spectra of the product mixture following exposure to air for 16 hr revealed the presence of a new carborane species, $(\text{CH}_3)_4\text{C}_4\text{B}_{10}\text{H}_{10}$, whose formation can be attributed to oxidative fusion of the $(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5^{2-}$ ligands in V. Also present were $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ and traces of a compound assigned as $[(\text{CH}_3)_2\text{C}_2\text{B}_8\text{H}_8]_2$.

Reaction of $[2,3-(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5]_2\text{FeH}_2$ with B_5H_9 . From earlier work³ it was known that the $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ ion and FeCl_2 rapidly form the red complex $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5]_2\text{FeH}_2$ in nearly quantitative yield. This compound was detected only in traces in the present study, implying that it undergoes further attack by borane reagents during the course of the reaction. As a direct test, a sample of $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5]_2\text{FeH}_2$ was

treated with an equimolar quantity of B_5H_9 in THF at room temperature, giving as sublimable products only the mixed-ligand complex II and the carboranes $(CH_3)_4C_4B_{11}H_{11}$ (III) and $(CH_3)_4C_4B_8H_8$; no starting metallacarborane was recovered.

Formation of C_4B_n Carboranes by Oxidative Fusion. The reactions described above produced several species lacking metal atoms and having the formulae $(CH_3)_4C_4B_nH_m$, where $n-2 \leq m \leq n+2$ and n has values from 8 to 16. Of these, the known compound $(CH_3)_4C_4B_8H_8$ was obtained in largest yield (up to 64%); since this carborane is known to form by fusion of two $[(CH_3)_2C_2B_4H_4]^{2-}$ ligands in transition metal complexes,²⁻⁴ it can be assumed that the new C_4B_n species originate in similar fashion, either by fusion to produce a single large cage or by coupling to generate biscarboranes in which two discrete polyhedra are linked by single B-B bonds. Where $m < n$ (as in $[(CH_3)_4C_4B_8H_7]_2$, mentioned above), a linked-cage structure is almost certain, but when $m \geq n$ a single polyhedral system is more likely; this conclusion is strengthened by the crystallographic characterizations of $(CH_3)_4C_4B_8H_8$ ⁵ and the $Co_2C_4B_6$ fusion products^{8,9} mentioned earlier. The parent-daughter relationship between the bis(carboranyl) metal complexes and the observed tetracarbon carborane products is further supported by experimental evidence in the case of the C_4B_{11} and C_4B_9 systems, as follows.

$(CH_3)_4C_4B_{11}H_{11}$: The white, air-stable, surprisingly volatile crystalline solid (III) was characterized from its unit- and high-resolution mass spectra, ^{11}B NMR, and 1H FT spectra (Tables I and II).

The mass spectrum exhibits an intense parent grouping (cutoff at m/e 240) with no other major peaks, indicating little fragmentation or hydrogen abstraction; this spectrum is compatible with a compact polyhedral species lacking bridge hydrogens. Thus, although III could be formulated as a linked-cage molecule, e.g., $(CH_3)_2C_2B_4H_4^-H_7B_7C_2(CH_3)_2$ with a B-H-B bridge on each cage (terminal hydrogens being absent on the boron atoms involved in the B-B linkage) such a structure appears inconsistent with the mass spectrum. The NMR spectra are complex, despite the mass spectroscopic purity of the sublimed product, and suggest that more than one isomer is present in solution. However, at $110^\circ C$ in C_6D_6 solution the original species, III, rearranged to a new isomer (IIIA) whose ^{11}B NMR spectrum was considerably simpler than that of III (Table I); the mass spectrum of the new isomer was essentially unchanged from that of III.

All efforts at obtaining crystals of III or IIIA suitable for X-ray analysis have thus far been unsuccessful, but if we are in fact dealing with a C_4B_{11} cage as suggested above, electron-counting arguments⁶ predict that it should be of the nido class, and specifically should resemble a 16-vertex closo polyhedron with one missing vertex. No known examples of either 16-vertex closo or 15-vertex nido cages have been structurally established (or even postulated prior to the present study), but Brown and Lipscomb²³ have predicted a unique stable geometry for closo- $B_{16}H_{16}^{2-}$, consisting of a tetracapped tetratruncated tetrahedron of T_d symmetry, i.e., a tetrahedron whose four corners have been truncated, each of

whose four hexagonal faces is capped by an additional vertex. Removal of one of the high-coordinate vertices yields an open basket (Figure 4) which may represent the gross geometry of the C_4B_{11} species; there are, of course, many possible ways of arranging the four carbons on this framework consistent with the NMR spectra (i.e., unsymmetrical) and a unique assignment is not possible at present.

Evidence that III is generated from an unstable precursor $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_7H_7]$ (IV) is given by the mass spectrum of the product mixture prior to workup in air, which revealed a strong peak grouping with a cutoff at m/e 298 corresponding to IV; on exposure to air this material disappeared, and III, not previously present, was observed.

$(CH_3)_4C_4B_9H_{11}$: The red mixed-ligand $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$ (II), obtained as described earlier, is moderately air-stable and does not appreciably degrade as a solid on exposure to air for several hours. In THF solution, however, treatment with a stream of O_2 gas did effect oxidative ligand fusion over an 18-hr period, forming a new tetracarbon carborane, $(CH_3)_4C_4B_9H_{11}$ (V). This compound was purified by sublimation and obtained in 42% yield as a colorless crystalline air-stable solid, m.p. $65^\circ C$. The mass spectrum of V exhibits a strong parent envelope and a more intense grouping corresponding to $(CH_3)_4C_4B_8H_8^+$ (m/e 204); the latter species must form by loss of BH_3 from V in the mass spectrometer, since contamination of the product by $(CH_3)_4C_4B_8H_8$ is excluded by the ^{11}B and 1H NMR spectra. The 1H FT NMR spectrum reveals only two methyl environments, consistent


with the presence of either a C_2 axis or, more probably, a mirror plane. The carborane skeleton is a formal 32-electron ($2n + 6$) system (taking into account the two "extra" hydrogens), on the basis of which we would expect a C_4B_9 arachno cage corresponding to a 15-vertex closo polyhedron with two vacant vertices. Elucidation of the structure clearly will require an X-ray crystallographic study.

Conclusions


Scheme I outlines the observed reactions, proposed intermediates, and probable sequences of product formation in the iron-borane-carborane system. While the tetracarbon carborane products other than $(CH_3)_4C_4B_8H_8$ have not yet been structurally characterized, the origin of these compounds from bis(carborane) iron complexes is clearly indicated. These results, combined with studies previously reported, further demonstrate that metal-promoted fusion and coupling²⁴ of borane and carborane substrates is a general phenomenon that can be exploited as a useful synthetic tool.

Observations regarding mechanism(s) have been presented elsewhere^{9,25} and it has been noted that fusion of carborane ligands seems to be associated with the presence of hydrogen atoms bound to the metal; significantly, this empirical correlation still holds true on the basis of the work reported here. However, we have established that the presence of an external oxidant (such as O_2) is not an absolute requirement for fusion to take place. Thus, while $[(CH_3)_2C_2B_4H_4]_2FeH_2$ converts rapidly to $(CH_3)_4C_4B_8H_8$ on exposure to

Scheme I



air with concomitant formation of an iron oxide/hydroxide mixture, the carborane forms slowly even when oxygen is rigorously excluded; in this case metallic iron is produced (Experimental Section). The net process involves a two-electron transfer from two carborane ligands to Fe^{2+} :



90%

The reaction is accompanied by color changes corresponding to intermediate iron-carborane complexes (including the known red compound $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]_2\text{FeH}_2$). It should be emphasized that this reaction is extremely slow in comparison to the very rapid (minutes) conversion of the red complex to $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ in the presence of oxidants such as O_2 or PbBr_2 .^{3,14}

Current investigations in our laboratory are concerned with mixed-ligand complexation and fusion involving large anions including $\text{C}_2\text{B}_9\text{H}_{12}^-$ and will be reported at a later time.

Experimental Section

Materials. 2,3-Dimethyl-2,3-nido-dicarbaheptaborane(8), 2,3- $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_6$, was prepared by the reaction of 2-butyne with pentaborane(9) in the presence of triethylamine as described elsewhere.²¹ Pentaborane(9) was obtained from the Callery Chemical Co., Callery, Pa. All other reagents were commercially obtained and used as re-

ceived. Tetrahydrofuran (THF) was dried over lithium aluminum hydride before use.

Spectra. Boron-11 and proton pulse Fourier transform NMR spectra at 32.1 and 100 MHz, respectively, were recorded on a JEOL PS-100P spectrometer interfaced to a JEOL-Texas Instruments EC-100 computer system. Unit-resolution mass spectra were obtained on a Hitachi Perkin-Elmer RMU-6E mass spectrometer, while high-resolution mass measurements were conducted on an AEL MS-902 double-focusing instrument equipped with an SRI chemical ionization source. All high-resolution spectra were recorded under chemical ionizing conditions in methane or argon-water. Infrared spectra were obtained on a Beckman IR-8 instrument.

Reaction of $B_5H_8^-$, $[(CH_3)_2C_2B_4H_5^-]$, and $CoCl_2$. A solution of $Na^+B_5H_8^-$ and $Na^+[(CH_3)_2C_2B_4H_5]^-$ was prepared by distillation of B_5H_9 (0.25 g, 3.87 mmol) and $(CH_3)_2C_2B_4H_6$ (0.40 g, 3.85 mmol) together onto NaH (0.553 g, 24.0 mmol) in 50 ml of THF. This solution was filtered in vacuo onto anhydrous $CoCl_2$ (0.63 g, 4.85 mmol) in a 250 ml round-bottom flask cooled in dry ice. The solution was allowed to warm to $-30^\circ C$ and stirred for 4 hr at that temperature, during which it turned dark brown and gas was evolved. After cooling the reaction flask in liquid nitrogen, the noncondensable gas (5.0 mmol) was pumped off. The solvent (THF) was distilled out at $-30^\circ C$ over a period of 2 h, leaving a dark brown residue. Dry N_2 was introduced into the flask and a vacuum sublimator was attached to the vessel. On heating

the flask at 70°C no material was collected on the cold finger, which was maintained at -78° by dry ice.

To the brown residue was added 100 ml of an equal-volume mixture of CH₂Cl₂ and hexane and the solution was stirred in air for 1 h. The dark brown solution was filtered and the solvent was distilled away from the filtrate. This residue was placed in a 200 ml flask and a vacuum sublimator was attached. On heating at 30°C in vacuo, a yellow sublimate condensed on the dry ice-filled cold finger. This material was characterized as [(CH₃)₂C₂B₄H₄]CoH[(CH₃)₂C₂B₇H₇], I, m.p. 79°C, 0.12 g (0.43 mmol), 22% yield based on carborane consumed. Mass measurement: calc. for ⁵⁹Co¹²C₈¹¹B₁₁¹H₂₅⁺ (P + 1 peak), 301.231; found, 301.235. Ir (CCl₄ vs. CCl₄): C-H bands at 3050(m), 3018(vs), 2845(s); B-H, 2570(vs); other bands at 1450(m) 1370(m), 1100(w,br).

When the reaction was conducted at room temperature (all other conditions being the same as described above), the results were identical except that the yield of I was slightly lower (15%).

Reaction of B₅H₈⁻, (CH₃)₂C₂B₄H₅⁻ and FeCl₂ (Simultaneous Addition of Reagents). A solution of Na⁺B₅H₈⁻ and Na⁺[(CH₃)₂C₂B₄H₅]⁻ in THF was prepared as described in the preceding experiment, from 1.48 g (23.1 mmol) of B₅H₉, 2.39 g (23.0 mmol) of (CH₃)₂C₂B₄H₆, and 1.50 g (62.5 mmol) of NaH in 50 ml of THF. This solution was filtered in vacuo onto anhydrous FeCl₂ (4.44 g, 35.0 mmol) in a 250 ml round-bottom flask cooled in dry ice. The solution was allowed to warm to room temperature and stirred for 24 hr, during which it became purple and gas was evolved. The vessel was cooled in liquid nitrogen

and the noncondensable gas (7.0 mmol) was expelled. The solvent (THF) was removed over a period of 2.0 hr, leaving a dark red-brown residue. Fractionation of the volatile products gave B_2H_6 (10.0 mmol) and $BH_3 \cdot THF$, identified from their ir spectra. Dry N_2 was introduced and a vacuum sublimator was attached to the vessel; however, heating the reaction residue at $70^\circ C$ produced no sublimate on the dry ice-cooled finger.

A few mL of liquid CCl_4 was condensed into the flask at $-196^\circ C$ and the mixture was allowed to warm to room temperature and stirred for 30 min, during which the solution was at first green and then turned to reddish brown; there was considerable heat and gas evolution. After cooling the flask in liquid nitrogen the noncondensable gas (2.50 mmol) was expelled, the flask was warmed once again to room temperature, and the solvent was removed in vacuo. Heating the reaction residue at $80^\circ C$ for 6 h caused a mixture of red and white crystalline solids to collect on the -78° cold finger. A mass spectrum of this mixture disclosed the presence of perchloroethane (C_2Cl_6) together with $[(CH_3)_2C_2B_7H_7]FeH_2[(CH_3)_2C_2B_4H_4]$, $(CH_3)_4C_4B_{11}H_{11}$, $(CH_3)_4C_4B_8H_8$, $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$, and $[(CH_3)_2C_2B_4H_4]_2FeH_2$.

The sublimed mixture was fractionated in vacuo through a trap at $0^\circ C$, which allowed the C_2Cl_6 (0.21 g) to pass through but retained the other materials. These substances were separated in air on a Waters Prep/500 High Pressure Liquid Chromatograph (HPLC) with two 500-ml silica gel columns employed in sequence and pure n-hexane as eluent. The pure products obtained included red $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$, II, 0.351 g (1.28 mmol), 11% yield, retention time (R_T) = 4.0 min, m.p.

71°C; mass measurement: calc. for $^{56}\text{Fe}^{12}\text{C}_8^{11}\text{B}_9^1\text{H}_{23}^+$, 274.1987, found 274.1981. $\text{Ir}(\text{CDCl}_3 \text{ vs. } \text{CDCl}_3)$: C-H, 2950(m), 2920(vs), 2850(m); B-H, 2570(vs). The previously reported compound $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8^{3,5a}$ appeared with $R_T = 7.6$ min, 1.51 g (7.40 mmol), 64 % yield. The new carborane $(\text{CH}_3)_4\text{C}_4\text{B}_{11}\text{H}_{11}$ eluted with $R_T = 10.0$ min as colorless crystals, m.p. 68°C, 63.8 mg (0.27 mmol), 2.3 % yield. Mass measurement: calc. for $^{12}\text{C}_8^{11}\text{B}_{11}^1\text{H}_{23}^+$, 240.2824, found 240.2820.

Small quantities of colorless products with $R_T > 10$ min were eluted and tentatively identified from their mass spectra as $[(\text{CH}_3)_2\text{C}_2\text{B}_8\text{H}_8]_2$ (mass 300) and $[(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_7]_2$ (mass 406).

Reaction of B_3H_8^- , $[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5]^-$, and FeCl_2 . Anhydrous ferrous chloride (0.700 g, 5.52 mmol) and $(\text{CH}_3)_4\text{N}^+\text{B}_3\text{H}_8^-$ (0.516 g, 4.49 mmol, prepared as described elsewhere)²⁰ were placed in a greaseless Pyrex reactor which was evacuated on the vacuum line, and 10 mL of dry THF was distilled into the reactor at -196°C. Separately, $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_6$ (4.4 mmol) was condensed onto 0.305 g (12.7 mmol) of NaH in 25 ml dry THF in vacuo. After reaction, the solution of $\text{Na}^+(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ was filtered in vacuo through sintered glass into the $\text{FeCl}_2/(\text{CH}_3)_4\text{N}^+\text{B}_3\text{H}_8^-$ solution, and the mixture was allowed to stand at room temperature for 48 hr. After cooling to -196°C, the noncondensable gas (4.2 mmol) was pumped off, the THF was removed under vacuum at room temperature over a 2-hr period, and 30 mL of dry CCl_4 was added. The mixture was stirred at room temperature for 2 hr, during which the color changed from greenish-brown to red-brown. Following removal of the CCl_4 by vacuum distillation over a 2-hr period, a sublimator containing a

U-tube was attached to the reactor under an N_2 atmosphere and the reactor was heated to $70-80^\circ C$ while the U-tube was immersed in a -78° bath. After materials had ceased collecting in the U-trap, the trap was warmed to $0^\circ C$ to permit the C_2Cl_6 (see above) to be pumped off. The remaining sublimate was separated via HPLC as described above, and the major products were the same as in the preceding experiment. Yields: $(CH_3)_4C_4B_8H_8$, 0.301 g (1.48 mmol, 67 %); $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$ (II), 90.2 mg (0.329 mmol, 15%); $(CH_3)_4C_4B_{11}H_{11}$, 13.2 mg (0.055 mmol, 2.5%).

Reaction of BH_4^- , $(CH_3)_2C_2B_4H_5^-$, and $FeCl_2$. A mixture of $NaBH_4$ (0.21 g, 5.6 mmol) and anhydrous $FeCl_2$ (0.55 g, 4.4 mmol) in a Pyrex bulb was evacuated on the vacuum line for 4 hr, and a solution of 3.0 mmol of $Na^+(CH_3)_2C_2B_4H_5^-$ in THF was added through a glass filter in vacuo. After stirring for 18 hr at room temperature in THF, during which a black solid formed and noncondensable gas was evolved, the THF was removed by distillation. The reaction residue was heated to $78^\circ C$, causing a bright red solid, identified mass spectroscopically as the known³ complex $[(CH_3)_2C_2B_4H_4]_2FeH_2$, to sublime into an adjoining U-trap cooled to $-78^\circ C$. The yield was 0.38 g (1.45 mmol, 97% yield). No other volatile products were detected.

Reaction of $B_5H_8^-$ and $FeCl_2$ Followed by $(CH_3)_2C_2B_4H_5^-$. Pentaborane(9) (13.9 mmol) was treated with 0.803 g (33.5 mmol) of NaH in 20 mL of THF, and the resulting solution of $Na^+B_5H_8^-$ was filtered in vacuo into a Pyrex bulb containing 0.940 g (7.40 mmol) of anhydrous $FeCl_2$. The

mixture was stirred for 1 hr at room temperature, during which the solution acquired a dark brown color and gas evolution occurred. At this point a solution of 13.7 mmol of $\text{Na}^+(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ in 20 ml THF (prepared as described above) was added by filtration, and the mixture was stirred at room temperature for 24 hr. No further color change was detected, but some additional gas evolution occurred. Following removal of THF by distillation in vacuo for 2 hr, a U-trap with an attached side-arm and bulb was attached to the reactor under a flow of N_2 , the apparatus was evacuated, and the reactor was heated to $70-80^\circ\text{C}$ while cooling the U-trap to -78°C . The sublimed material was a mixture of red and white solids which on mass spectroscopic analysis was found to contain II, $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$, and a new compound, $[(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5]\text{FeH}_2$ (V), identified from its parent grouping (m/e 286) and pattern of peak intensities, which conform to the postulated composition. Attempts to separate and purify this new material by HPLC in air and by column chromatography on silica in an atmosphere of dry N_2 did not succeed due to the instability and small quantity of V. The residue remaining after sublimation was treated with CCl_4 as described above, which gave additional quantities of the same sublimable red and white products.

When the product mixture containing V was exposed to air for 16 hr, the mass spectrum of the material contained a new, intense peak grouping at m/e 228 corresponding to $(\text{CH}_3)_4\text{C}_4\text{B}_{10}\text{H}_{10}$, together with $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ (m/e 204) and a small grouping at m/e 300 assigned to $[(\text{CH}_3)_2\text{C}_2\text{B}_8\text{H}_8]_2$.

Reaction of $[(CH_3)_2C_2B_4H_5]_2FeH_2$ with B_5H_9 . A 10 mg sample of the iron complex^{and} an equimolar quantity of B_5H_9 were stirred in THF at room temperature for 24 hr, after which the solvent was removed by distillation at room temperature and the products were analyzed by mass spectroscopy. The mass spectrum revealed three major components: $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$ (II), $(CH_3)_4C_4B_{11}H_{11}$ (III), and $(CH_3)_4C_4B_8H_8$. The original complex was not present in significant quantity.

Synthesis of $(CH_3)_4C_4B_9H_{11}$ from $[(CH_3)_2C_2B_4H_4]FeH_2[(CH_3)_2C_2B_5H_5]$. A solution of 30 mg (0.11 mmol) of the iron complex in 50 ml of THF was placed in a 3-necked 200 ml flask, and O_2 gas was bubbled through the solution over an 18-hr period with 30 ml of THF added every 2 hr. During this time the originally violet-red solution changed to pale yellow-orange with some insoluble residue at the bottom of the flask. The solution was filtered and the solvent removed under vacuum at $0^\circ C$ over a 40 min period.

A vacuum sublimator was attached to the flask and the apparatus was evacuated; a white sublimate condensed on the dry ice-filled cold finger while the reactor was maintained at room temperature. The white compound, m.p. $65^\circ C$ (10 mg, 0.046 mmol, 42% yield) was characterized as $(CH_3)_4C_4B_9H_{11}$ from its mass spectrum, ^{11}B NMR and 1H NMR spectra.

Conversion of $[(CH_3)_2C_2B_4H_4]_2FeH_2$ to $(CH_3)_4C_4B_8H_8$ in the Absence of Oxygen. A THF solution of $Na^+[(CH_3)_2C_2B_4H_5]^-$, prepared as described above from 2.38 mmol of $(CH_3)_2C_2B_4H_6$ and excess NaH, was filtered

through sintered glass into an evacuated flask containing 0.816 g FeCl_2 (1.48 mmol) and the mixture was stirred at room temperature over a 24-hr period. At the end of this time the solution was dark purple. Following removal of solvent by vacuum distillation, the purple residue was heated at 100° , causing white crystalline $(\text{CH}_3)_4\text{C}_4\text{B}_8\text{H}_8$ (0.22 g, 1.08 mmol, 90.6% yield) to condense in an adjacent U-trap cooled to 0°C . The carborane was identified from its mass spectrum by comparison with the spectrum of an authentic sample. The dark solid remaining in the reactor was recognized as metallic iron from its attraction to a bar magnet.

Acknowledgments. This work was supported in part by the Office of Naval Research. We thank Mr. David Finster for recording the ^{11}B FT NMR spectra.

References and Notes

- (1) Recent reviews: (a) Nicholas, K.M.; Nestle, M.O.; Seyferth, D., in "Transition Metal Organometallics in Organic Synthesis", Vol. II, H. Alper, Ed.; Academic Press:New York, 1978, Chapter 1; (b) Advances in Organometallic Chemistry, Vol. 17, Stone, F.G.A. and West, R., Eds.: Academic Press, New York, 1979; (c) Maitlis, P. M., Accounts Chem. Res., 1976, 9, 93.
- (2) Grimes, R. N., Accounts Chem. Res., 1978, 11, 420.
- (3) Maxwell, W. M.; Miller, V. R.; Grimes, R. N. Inorg. Chem. 1976, 15, 1343.
- (4) Maxwell, W. M.; Miller, V. R.; Grimes, R. N. J. Am. Chem. Soc., 1976, 98, 4818.
- (5) (a) Freyberg, D. P.; Weiss, R.; Sinn, E.; Grimes, R. N. J. Am. Chem. Soc. 1977, 16, 1847; (b) Grimes, R. N.; Maxwell, W. M.; Maynard, R. B.; Sinn, E., submitted for publication.
- (6) (a) Wade, K. Adv. Inorg. Chem. Radiochem. 1976, 18, 1; (b) Rudolph, R. W. Accounts Chem. Res., 1976, 9, 446; (c) Mingos, D.M.P. Nature (London), Phys. Sci. 1972, 236, 99.
- (7) Hosmane, N. S.; Furia, J.; Finster, D. C.; Grimes, R. N., to be published.
- (8) Wong, K-S.; Bowser, J. R.; Pipal, J. R.; Grimes, R. N., J. Am. Chem. Soc. 1978, 100, 5045.
- (9) Pipal, J. R.; Grimes, R. N. Inorg. Chem. 1979, 18, 1936.
- (10) Grimes, R. N.; Beer, D. C.; Sneddon, L. G.; Miller, V. R.; Weiss, R. Inorg. Chem. 1974, 13, 1138.

- (11) Miller, V. R.; Weiss, R.; Grimes, R. N. J. Am. Chem. Soc. 1977, 99, 5646.
- (12) George, T. A.; Hawthorne, M. F. J. Am. Chem. Soc. 1969, 91, 5475.
- (13) This is evidently the case in the closely related species $[2,3-(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{FeH}_2$, in which the ligands are mutually rotated²⁵ by 90° , yet the four methyl groups are equivalent in the proton NMR spectrum.³
- (14) Maxwell, W. M.; Wong, K-S.; Grimes, R. N. Inorg. Chem. 1977, 16, 3094.
- (15) Observed ^{11}B shifts for $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4^{2-}$ complexes range from +10.8 to -18.3,^{3,4,10} and for complexes of $\text{C}_2\text{B}_7\text{H}_9^{2-}$ range from +21.0 to -25.2.^{12,16}
- (16) Jones, C. J.; Francis, J. N.; Hawthorne, M. F. J. Am. Chem. Soc. 1972, 94, 8391.
- (17) Miller, V. R.; Grimes, R. N. J. Am. Chem. Soc. 1973, 95, 2830; 1975, 97, 4213.
- (18) Sneddon, L. G.; Beer, D. C.; Grimes, R. N. J. Am. Chem. Soc. 1973, 95, 6623.
- (19) Control experiments demonstrate that the $(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5^-$ and B_5H_8^- ions do not interact significantly with each other in THF at room temperature, in the absence of a transition metal reagent.
- (20) Nainan, K. C.; Ryschkewitsch, G. E. Inorg. Nucl. Chem. Lett. 1970, 6, 765.
- (21) Hosmane, N.S.; Grimes, R. N. Inorg. Chem. 1979, 18, 3294.

- (22) Wegner, P. A. in "Boron Hydride Chemistry", E. L. Muetterties, Ed., Academic Press: New York, 1975, Chapter 12, and references therein.
- (23) Brown, L. D.; Lipscomb, W. N. Inorg. Chem., 1977, 16, 2989.
- (24) An example of oxidative coupling (single-bond linkage of two cages) in a bis(carboranyl) metal complex is the thermal conversion of $\mu, \mu' - [(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5]_2\text{Hg}$ to $5,5' - [(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_5]_2$ and elemental mercury: Hosmane, N. S.; Grimes, R. N. Inorg. Chem. 1979, 18, 2886.
- (25) Pipal, J. R.; Grimes, R. N. Inorg. Chem. 1979, 18, 263.
- (26) The order of addition of reagents is also important in the cobalt system, which under certain conditions (reaction of CoCl_2 with B_5H_8^- prior to addition of carborane anion) generates mixed borane-carborane complexes of Co(III): L. Borodinsky and R.N. Grimes, to be submitted.

Table I. ^{11}B FT NMR Data (32.1 MHz)^a

Compound	δ^b (J, Hz)	rel. areas
$[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{CoH}[(\text{CH}_3)_2\text{C}_2\text{B}_7\text{H}_7]$, I	12.2(141), 0.0(142), -7.7(150), ^c -19.3(161), -26.4(171)	3:3:1:2:2
$[(\text{CH}_3)_2\text{C}_2\text{B}_4\text{H}_4]\text{FeH}_2[(\text{CH}_3)_2\text{C}_2\text{B}_5\text{H}_5]$, II	26.3(176), 2.9(-170), ^c -2.7(157), -14.5(156) ^d	1:1:4:3
$(\text{CH}_3)_4\text{C}_4\text{B}_{11}\text{H}_{11}$, III ^{e,f} (C_6D_6)	16.9, 10.8, 2.9, -2.6, -8.7, -20.9, -23.6, -26.6, -30.0	1:3:3:5:3 1:1:1:1 ^g
$(\text{CH}_3)_4\text{C}_4\text{B}_{11}\text{H}_{11}$, IIIA (rearranged isomer, in C_6D_6 .	8.0(170), 1.65(160), -3.6(174) -32.2(155) ^h	1:1:8:1
$(\text{CH}_3)_4\text{C}_4\text{B}_9\text{H}_{11}$, V ⁱ	1.3, -3.6, -5.4, -15.2	2:3:3:1

^aAll spectra were obtained in CDCl_3 solution except where otherwise indicated. In each case, both uncoupled and ^1H -decoupled spectra were obtained; in the decoupled spectra all H-B doublets collapse to singlets. ^bChemical shifts are relative to $\text{BF}_3 \cdot \text{O}(\text{C}_2\text{H}_5)_2$ with a positive sign indicating deshielding. ^cEstimated from overlapped peaks. ^dAsymmetric peak with shoulder on upfield side. ^eChemical shifts are given for the proton-decoupled spectrum. ^fProbable mixture of isomers. ^gApproximate areas. ^hSmall peaks at $\delta + 36.6$, -9.0 , and -13.6 , also observed, were assigned to minor side products of the isomerization. ⁱProton-decoupled spectrum; resonances in the uncoupled spectrum were heavily overlapped and couplings were not measurable.

Table II. ^1H FT NMR Data (100 MHz)

Compound	δ^a (rel area)	assignment
I (CDCl_3)	2.35(3), 2.23(3)	CH_3
	-12.3^b ($w_{1/2} = 45 \text{ Hz}$)	Co-H
II (CDCl_3)	2.43(3), 2.26(3)	CH_3
	-12.4^b ($w_{1/2} = 75 \text{ Hz}$)	Fe-H
III (C_6D_6) ^c	1.60(8), 1.43(5), 1.17(1), -0.05(1)	CH_3
IIIA (rearranged isomer in C_6D_6)	2.14(1), 1.73(1), 1.38(1), 1.27(1)	CH_3
V (CDCl_3)	2.30(1), 1.97(1)	CH_3

^aChemical shifts relative to $(\text{CH}_3)_4\text{Si}$ with positive sign indicating deshielding. ^bBroad peak, area not measurable. ^cProbable mixture of isomers; areas given are approximate.

Figure Captions

- Figure 1. Schematic diagram of the oxidative fusion process, where hemispheres represent anionic carborane ligands and M is a transition metal cation.
- Figure 2. Formation of $[(CH_3)_2C_2B_4H_4]CoH[(CH_3)_2C_2B_7H_7]$, I, with proposed structure of I indicated.
- Figure 3. Formation of mixed-ligand ferracarboranes and conversion to tetracarbon carboranes.
- Figure 4. Possible geometry of a 15-vertex nido cage corresponding to $(CH_3)_4C_4B_{11}H_{11}$, derived from a T_d closo 16-vertex polyhedron²³ by removal of a high-coordinate vertex.




Figure 1





Figure 2

Figure 3






Figure 4

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 33	2. GOVT ACCESSION NO. AD-A087152	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Transition Metal-Promoted Oxidative Fusion With Carborane and Borane Anions. Synthesis of C_4B_n (n = 8-11) Carborane Systems in the Presence of $B_5H_8^-$ or $B_3H_8^-$ Ions	5. TYPE OF REPORT & PERIOD COVERED Interim	
7. AUTHOR(s) Narayan S. Hosmane and Russell N. Grimes	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Virginia Department of Chemistry Charlottesville, Va. 22901	8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0305	
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Branch, Office of Naval Research Arlington, Va. 22217	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-569	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE July, 1980	
	13. NUMBER OF PAGES 33	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Metallaborane Carborane Metallacarborane Supra-icosahedral Carborane Ferracarborane Oxidative Fusion Cobaltacarborane Tetracarbon Carborane		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See title page		

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
		Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024	1	Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Dr. D. Seyferth Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Professor H. Abrahamson University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Branch Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Branch Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

P4-2/A25

472:GAN:716:tan
78u472-608

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1