

AD-A087 628

VIRGINIA UNIV CHARLOTTESVILLE SCHOOL OF MEDICINE F/G 6/5  
THE EFFECTS OF ACUTE ALTERATIONS IN HEMODYNAMICS, OXYGEN AVAILA--ETC(U)  
MAR 80 W P RITCHIE DAMD17-74-C-4014

NL

UNCLASSIFIED

1 of 1  
2/2/80




END

DATE

FILED

9-80

DTIC



LEVEL

AD \_\_\_\_\_

THE EFFECTS OF ACUTE ALTERATIONS IN HEMODYNAMICS,  
OXYGEN AVAILABILITY AND ACID-BASE BALANCE ON  
THE PERMEABILITY OF THE GASTRIC MUCOSA

12

ADA 087628

ANNUAL PROGRESS REPORT  
(FOR THE PERIOD 1 OCT. 78 to 30 SEPT. 79)

DATE OF REPORT MARCH 1980

BY

WALLACE P. RITCHIE, JR., M.D.

SUPPORTED BY  
U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND  
FORT DETRICK, FREDERICK, MARYLAND 21701

CONTRACT NO. DAMD 17-74-C-4014

UNIVERSITY OF VIRGINIA SCHOOL OF MEDICINE  
CHARLOTTESVILLE, VIRGINIA 22908

DTIC  
SELECTED  
AUG 7 1980  
C

APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED

THE FINDINGS IN THIS REPORT ARE NOT TO BE CONSTRUED AS AN  
OFFICIAL DEPARTMENT OF THE ARMY POSITION UNLESS SO DESIGNATED  
BY OTHER AUTHORIZED DOCUMENTS.

DDC FILE COPY

80 7 31 066

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A087628	(9)
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
The Effects of Acute Alterations in Hemodynamics, Oxygen Availability, and Acid-Base Balance on the Permeability of the Gastric Mucosa.		Annual Progress Report. 1 Oct 1978-30 Sep 1979
6. PERFORMING ORG. REPORT NUMBER		
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Wallace P. Ritchie, Jr., M.D., Ph.D.		(15) DAMD 17-74-C-4014
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
University of Virginia School of Medicine Charlottesville, Virginia 22901		61102A (16) 3A161102B71R 01.073 (17) 01
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
U.S. Army Medical Research & Development Command Frederick, Maryland 21701		(11) March 1980
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
(12) 8		8
15. SECURITY CLASS. (of this report)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
Unclassified		
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release, distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Blood flow, ischemia, permeability, hydrochloric acid, histamine receptors, primary bile acids, secondary bile acids, ulcerogenesis.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Using a previously described model for acute gastric mucosal ulcerogenesis, developed under the auspices of the current contract ( <del>Gastroent. 68-699, 1975</del> ), studies from this laboratory during the period covered by this progress report indicate (1) that, in the face of an imposed topical acid load, H <sub>1</sub> and H <sub>2</sub> receptor blockade, either alone or in combination, effects no change in gastric mucosal permeability to cations or in mucosal blood flow and affords no protection against bile acid-ischemia induced ulcerogenesis; (2) that the addition of exogenous histamine neither protects nor augments lesion formation under		

#20 Continued:

these circumstances; (3) that the dihydroxy secondary bile acids are more damaging to gastric mucosa than are the trihydroxy primary bile acids; and (4) that in non-ischemic bile acid-treated gastric mucosa, mucosal blood flow increases in proportion to the magnitude of H<sup>+</sup> back-diffusion induced, a response not mediated by histamine receptors.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution / _____	
Availability Codes	
Dist	Availability Codes
<b>A</b>	Special

PROGRESS REPORT: CONTRACT DAMD 17-74-C-4014

I. TITLE OF RESEARCH CONTRACT:

The Effects of Acute Alterations in Hemodynamics, Oxygen Availability, and Acid-Base Balance on the Permeability of the Gastric Mucosa.

II. PRINCIPAL INVESTIGATOR:

Wallace P. Ritchie, Jr., M.D., Ph.D.  
Department of Surgery  
University of Virginia School of Medicine  
Charlottesville, Virginia 22908  
472-36-6392

III. PERIOD COVERED: 1 October 1978 to 30 September 1979

IV. PROGRESS REPORT:

(1) Effect of H<sub>1</sub> and H<sub>2</sub> receptor blockade on gastric mucosal permeability to cations and bile acid-induced ulcerogenesis. Recent reports suggest that, in the presence of topical acid, combined H<sub>1</sub> and H<sub>2</sub> receptor blockade decreases gastric mucosal permeability and H<sub>2</sub> blockade alone ameliorates bile-acid-induced "stress ulcer". The present study was undertaken to re-examine these assertions using the sensitive model outlined. Ex-vivo chambered wedges of mucosa were prepared in four groups of animals, each of which was studied during nine sequential 15 minute periods. Group A mucosa were exposed, during 1-3, to topical acid test solution alone (ATS), during 4-6, to ATS+2mM Na taurocholate (TC), and during 7-9 to ATS+TC+wedge-specific vasopressin (VP),  $5 \times 10^{-3}$ U/Kg-min. Groups B, C, and D were treated analogously except that group B was pretreated with cimetidine, 300mgm IV, group C with pyribenzamine, 10mgm/Kg IM, and group D, with both. The dose of cimetidine employed has been shown to produce significant inhibition of histamine stimulated gastric acid production in this model for a time period comparable to that of the current experiment. The dose of the H<sub>1</sub> receptor blocking agent was precisely the same as that reported to reduce, in combination with cimetidine, mucosal permeability to cations.

The parameters evaluated included (1) net flux of H<sup>+</sup> and Na<sup>+</sup> ( $\mu$ Eq); (2) the potential difference (mV); (3) aminopyrine clearance (ml/min); and (4) the lesion index, graded 0-5. The results per 15 minutes ( $\pm$ SEM) indicate that, in mucosa exposed to ATS,  $\Delta$ H<sup>+</sup> = -43 $\pm$ 12,  $\Delta$ Na<sup>+</sup> = +133 $\pm$ 13, PD = -63 $\pm$ 1, AC: 2.3 $\pm$ 0.4, and LI=0. During ATS+TC,  $\Delta$ H<sup>+</sup> = -219 $\pm$ 21,  $\Delta$ Na<sup>+</sup> = +206 $\pm$ 17, PD = -45 $\pm$ 1, AC = 2.8 $\pm$ 0.5, and LI=0. H<sub>1</sub>, H<sub>2</sub> and H<sub>1</sub> + H<sub>2</sub> blockade effected no significant change under either circumstance. In bile acid treated ischemia mucosa;

GROUP	$\Delta H^+$	$\Delta Na^+$	$\overline{PD}$	AC	LI
Control (n=7)	-233±20	+323±20	-24±1	1.1±0.1	3.3±0.4
H <sub>1</sub> (n=6)	-233±27	+333±24	-20±2	0.9±0.2	3.7±0.2
H <sub>2</sub> (n=5)	-266±32	+361±22	-24±1	1.2±0.1	3.2±0.4
H <sub>1</sub> +H <sub>2</sub> (n=6)	-277±24	+324±17	-21±1	1.0±0.1	3.5±0.2

Thus, in the face of an imposed topical acid load, H<sub>1</sub> and H<sub>2</sub> blockade, both alone and in combination, effects no change in gastric mucosal permeability to cations or blood flow and affords no protection against bile acid-induced lesion formation.

(2) Effect of histamine stimulation on the magnitude of acute gastric mucosal damage induced by bile acids and ischemia.

It has been demonstrated in-vitro that burimamide-inhibited amphibian gastric mucosa is less resistant to bile acid induced damage, as judged by electrical indices, than is spontaneously secreting mucosa. The present study used ex-vivo vascularized chambered wedges of proximal canine mucosa to examine the converse: the potential protective effect of active H<sup>+</sup> secretion induced by histamine on acute mucosal ulcerogenesis caused by the topical application of Na taurocholate in acid solution and concomitant pharmacologic ischemia. Methods: Groups of dogs (n=5-6ea) were studied during 3 consecutive 30 minute periods. Group A= topical acid test solution (ATS) during periods (1), (2), (3). Group B= (1) ATS, (2) ATS, (3) ATS+vasopressin (VP), 0.1x10<sup>-2</sup>U/Kg-min delivered via the splenic artery. Group C= (1) ATS, (2) ATS+H<sup>+</sup>TC, (3) ATS+H<sup>+</sup>TC+VP. Parameters evaluated during each period=net H<sup>+</sup> flux, aminopyrine clearance, and lesion index, graded 0-5. Results: In the absence of VP, TC significantly increased net H<sup>+</sup> loss and AC relative to ATS; H resulted in significant H<sup>+</sup> gain and further increased AC relative to ATS+TC; no lesions were noted under any circumstance. The results observed during period 3 (±SEM/30min):

	ATS	ATS+VP	ATS+TC+VP	ATS+H <sup>+</sup> TC+VP
$\Delta H^+$ ( $\mu$ Eq)	-109±43*	-108±60*	-242±53	-55±57*
AC (ml/min)	1.57±0.27	0.70±0.16*	1.13±0.27	1.57±0.49
LI (0-5)	0*	0.3±0.2*	2.4±0.3	1.6±0.5

\*Significant difference vs. ATS+TC+VP.

Thus, the combination of topical bile acid in acid solution and relative mucosal ischemia (compared to ATS+TC, where AC=3.36±0.69ml/min) is acutely ulcerogenic. Concomitant histamine infusion neither protects nor augments lesion formation under these experimental conditions.

(3) Mediators of Bile Acid-Induced Enhancement of Gastric Mucosal Blood Flow. The topical application of bile acid in acid solution to non-ischemic gastric mucosa (GM) increases mucosal blood flow (MBF), an important cytoprotective phenomenon since, when it alone is blunted, mucosal ulcerogenesis results. The present study addressed the hypothesis that the magnitude of this response is related to the absolute amount of H<sup>+</sup> "back diffusion" induced and that endogenous histamine release is the intramucosal mediator. Using vascularized chambered ex-vivo wedges of GM, 3 groups of dogs were studied during 3 sequential 30 minute periods. Group A (n=6): topical neutral test

solution (NTS; pH=7.0) during 1, NTS+2.5 mM sodium taurocholate (TC) during 2, and during 3, NTS+5 mM TC. Group B (n=11): topical acid test solution (ATS=100 mM HCl) during 1, ATS=2.5 TC during 2, and ATS=5 TC during 3. Group C(n=7) was treated in the same manner as group B except that each dog was pretreated with the H<sub>1</sub> and H<sub>2</sub> blockers, pyribenzamine (10 mg/Kg IM), and cimetidine (300 mg IV). Parameters evaluated: (1) net flux H<sup>+</sup>, Na<sup>+</sup> ( $\Delta$ H<sup>+</sup>,  $\Delta$ Na<sup>+</sup>), (2) MBF, determine by radiolabelled microsphere embolization, (3) potential difference (PD). Results ( $\pm$ SEM)/30 minutes:

	NTS		ATS		NTS		ATS	
	0	-96 $\pm$ 22*	0	-405 $\pm$ 40* <sup>+</sup>	0	-683 $\pm$ 37* <sup>+</sup>	0	-38 $\pm$ 1* <sup>+</sup>
$\Delta$ H <sup>+</sup> ( $\mu$ Eq)								
PD (mV)	-59 $\pm$ 2	-62 $\pm$ 1	-53 $\pm$ 2*	-47 $\pm$ 1* <sup>+</sup>	-45 $\pm$ 2*	-38 $\pm$ 1* <sup>+</sup>		
MBF (ml/min)	4.7 $\pm$ 0.8	6.0 $\pm$ 0.7	4.4 $\pm$ 0.8	8.4 $\pm$ 1.0* <sup>+</sup>	4.2 $\pm$ 1.0	10.4 $\pm$ 1.0* <sup>+</sup>		

\*Significant difference vs NTS; <sup>+</sup> Significant difference vs 0 nM TC, same group.

The addition of blockers effected no change from ATS alone in any parameter at any {TC} except at 5 TC, where MBF increased (to 15.8 $\pm$ 2.6 ml/min.). CONCLUSIONS: (1) In non-ischemic GM, MBF increases in the presence of greater than normal  $\Delta$ H<sup>+</sup>. (2) The amplitude of this response is dependent on the absolute magnitude of the increase in H<sup>+</sup> "back diffusion" observed (p<0.01). (3) Enhanced MBF is not mediated by H<sub>1</sub> and H<sub>2</sub> receptor related histamine release. In fact, receptor blockage augments the response at 5mM TC.

(4) Ulcerogenic Potential of Primary vs. Secondary Bile Acids in Gastric Mucosa. Although the secondary (2<sup>o</sup>) bile acids are more damaging to colonic mucosa than are the primary (1<sup>o</sup>) bile acids, the existence of a similar circumstance in gastric mucosa has been inadequately documented. The present study addressed this problem by comparing the effect of an identical (and physiologic) concentration of the 2<sup>o</sup> BA, taurodeoxycholic a. (TDC) to its parent 1<sup>o</sup> BA taurocholic a. (TC) in both ischemic and non-ischemic GM. METHODS: Using chamber ex-vivo wedges of proximal GM, 2 groups of dogs were studied during 9 sequential 15 minute periods. Group A (4 dogs)=topical acid test solution alone (ATS=100mM HCl) during periods 1-3; ATS+1mM TC during 4-6; and, during 7-9, ATS+1mM TC+vasopressin (VP=5x10<sup>-3</sup>U/Kg-min) infused into the splenic artery. Group B (4 dogs) were similarly treated except that the BA employed was TDC. Parameters evaluated=(1) net flux H<sup>+</sup>, Na<sup>+</sup> ( $\Delta$ H<sup>+</sup>,  $\Delta$ Na<sup>+</sup>), (2) potential difference (PD), (3) mucosal blood flow (MBF), determined by radiolabeled microsphere embolization, and (4) the lesion index (LI), graded 0-5 by an independent observer using photographs. RESULTS ( $\pm$ SEM)/15 minutes:

GROUP	ATS (1-3)		ATS+1mM BA(4-6)		ATS+1mM BA+VP (7-9)	
	A	B	A	B	A	B
PD (mV)	-64 $\pm$ 1	-63 $\pm$ 1	-60 $\pm$ 1 <sup>+</sup>	-48 $\pm$ 1* <sup>+</sup>	-41 $\pm$ 1 <sup>+</sup>	-28 $\pm$ 1* <sup>+</sup>
$\Delta$ H <sup>+</sup> ( $\mu$ Eq)	-8 $\pm$ 15	-1 $\pm$ 18	-97 $\pm$ 13 <sup>+</sup>	-157 $\pm$ 20* <sup>+</sup>	-105 $\pm$ 21 <sup>+</sup>	-250 $\pm$ 30* <sup>+</sup>
MBF (ml/min)	11.2 $\pm$ 2.0	12.0 $\pm$ 2.5	9.0 $\pm$ 1.4	13.6 $\pm$ 4.8	2.5 $\pm$ 0.7 <sup>+</sup>	2.9 $\pm$ 0.5 <sup>+</sup>
LI (0-5)	0	0	0	0	0.9 $\pm$ 0.4 <sup>+</sup>	2.6 $\pm$ 0.4 <sup>+</sup>

+P 0.05 vs. ATS, same group; \*p 0.025 vs. A(TC)

Thus, at an identical (and physiologic) concentration, 2° BA induce a greater depression on PD, a more profound increase in mucosal permeability to H<sup>+</sup>, and in ischemic GM, a more severe degree of gross mucosal damage than do 1° BA. CONCLUSION: 2° BA are more damaging to GM than are 1° BA, an effect undoubtedly related to their greater lipid solubility and consequent capacity to disrupt cell membranes.

V. PUBLICATIONS DURING THE CONTRACT YEAR:

1. Ritchie, W.P., Jr. and Cherry, K.J., Jr.: Influence of [H<sup>+</sup>] on Bile Acid Induced Acute Gastric Mucosal Ulcerogenesis. Ann. Surg. 189:637-642, 1979.
2. Ritchie, W.P., Jr. and McRae, D.B., Jr.: Effect of H<sub>1</sub> and H<sub>2</sub> Receptor Blockade on Gastric Mucosal Permeability to Cations and Bile Acid Induced Ulcerogenesis. Gastroent. 76:1227, 1979. Abstract.
3. Ritchie, W.P., Jr., McRae, D.B., Jr. and Felger, T.S.: A-V Acid-Base Balance During Acute Gastric Mucosal Ulcerogenesis. Am. J. Surg. 139:22-26, 1980.
4. Ritchie, W.P., Jr.: Invited Commentary on "Restraint Induced Stress Ulcer. Parts I and II" World Journal of Surgery 4:101-103, 1980.
5. Felger, T.S., and Ritchie, W.P., Jr.: Ulcerogenic Potential of 1° vs. 2° Bile Acids in Gastric Mucosa. Surgical Forum 30:324-326, 1979.

## DISTRIBUTION LIST

4	copies	HQDA (SGRD-SI) Fort Detrick Frederick, MD. 21701
12	copies	Defense Technical Information Center (DTIC) ATTN: DTIC-DDA Cameron Station Alexandria, Virginia 22314
1	copy	Dean School of Medicine Uniformed Services University of the Health Sciences 4301 Jones Bridge Road Bethesda, Maryland 20014
1	copy	Superintendent Academy of Health Sciences, US Army ATTN: AHS-COM Fort Sam Houston, Texas 78234
4	copies	Commander Letterman Army Institute of Research (LAIR) Bldg. 1110 ATTN: Dr. J. Ryan Neville Presidio of San Francisco, CA 94129