

AD-A088 334

FLORIDA STATE UNIV TALLAHASSEE DEPT OF PSYCHOLOGY F/G 6/16
EVOKED RESPONSE MEASURES OF RESOURCE ALLOCATION: EFFECTS OF PRI--ETC(U)
MAY 80 L F ELFNER, R R STANNY, W R HOWSE AFOSR-79-0097

UNCLASSIFIED

AFOSR-TR-80-0597

NL

1 - 1
2 - 100-100


END

DATE

FILED

9 80

DTIC



Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

LEVEL II

1. REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
2. REPORT NUMBER AFOSR/TR-80-0597		3. RECIPIENT'S CATALOG NUMBER AD-A088334	
4. TITLE (and Subtitle) Evoked Response Measures of Resource Allocation: Effects of Primary Task Load		5. TYPE OF REPORT & PERIOD COVERED Final Scientific Report	
6. PERFORMING ORG. REPORT NUMBER		7. CONTRACT OR GRANT NUMBER(s) AFOSR-79-0097	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Psychology Florida State University Tallahassee, Florida 32306		9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2313 D9	
10. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research (NL) Bolling Air Force Base, D.C. 20332		11. REPORT DATE 31 May 1980	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 12 11		13. NUMBER OF PAGES 11	
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. SUPPLEMENTARY NOTES			
18. KEY WORDS (Continue on reverse side if necessary and identify by block number) Evoked Potential Wave V Brainstem Evoked Potential Resource Allocation Perpetual-motor task			
19. ABSTRACT (Continue on reverse side if necessary and identify by block number) The major emphasis of the research was to assess the utility of employing event-related potentials in monitor type tasks. The work manipulation paradigm was employed. Auditory probes were presented along a secondary channel unrelated to the perceptual-motor task. In the initial study amplitude and latency measures of Wave V of the brain stem evoked potential were measured to auditory probes under conditions of tracking a visual target with a joystick, and just observing the visual display. It was noted that a reduction in amplitude and an increase in latency of Wave V occurred during the			

AD A088334

DDC FILE COPY

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 63 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

401412 Lu

80 8 14 01

20. CONTINUED

tracking task. Although not statistically significant all subjects demonstrated the same effect.

Accession For	
NTIS GRA&I	☒
DDC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

INTRODUCTION

That ERP is a useful measurement of higher-order functioning is "fait accompli." The level of the central nervous system that reflects higher-order functioning is in dispute. Oatman (1976) has demonstrated attentional effects at several levels in the auditory system of cat.

Since attentional diversion from auditory stimuli is a measure of import in assessing the effect of a visual primary task, we decided to observe the early potentials of the auditory evoked response to determine if a visual motor task would affect the neuronal processing of auditory stimuli at the level of the brainstem. An odd-ball uncertainty task (Sutton, 1965) was employed since the P_3 wave is very evident under this procedure. The visual task consisted of playing one of the commercial pong games on a TV screen.

It is hypothesized that the Wave V of the brainstem ERP will show a reduction in amplitude and a longer latency during a visual motor task than during a simple odd-ball procedure.

METHOD

SUBJECTS: Five undergraduate students volunteered to perform in the study. All exhibited normal hearing with no demonstrable central nervous system problems.

APPARATUS: A TV game (Radio Shack Model 603056) provided the perceptual-motor task. The game was projected on a Sony 19" video screen. The auditory signal (2000 Hz) was generated by a Hewlett-Packard (Model 200) audio oscillator and delivered to a set of TDH-39 earphones. A Grason-Stadler probability generator (Model 1284) randomized the presentation of the target stimuli. Intensity of the Auditory stimuli was controlled by a pair of (Tech Lab,

Approved for public release;
distribution unlimited.

Model 850) attenuators. Shaping (onset, off-set 2.5 msec) and duration (7.5 msec) of the signals and the ISI were controlled by Grason-Stadler (Model 829E) switches and (Model 471) timers. The ERP were averaged and hard copied on a Nicolet (Model 1074) averager and a Hewlett-Packard XY plotter. The recording electrode was at vertex, the reference electrode at right mastoid, a ground electrode at left mastoid. The EEG was filtered below 0.1 kHz and above 3.0 kHz and amplified by 10^6 through the use of cascaded preamplifiers (Grass P15, and Tektronix type 122).

PROCEDURE: Initially S was wired with the three electrodes. Impedance was kept below 5 k ohms. Then a short session of training to detect the 2dB softer target stimuli was employed. S then was given practice in playing the pong game until reasonably stable in terms of performance level.

Auditory stimuli were delivered at a repetition rate of 30 per sec. Detection of the softer rare-occurring stimulus (33dB SL) was reported with a button press. Each ERP represented a minimum of 4096 evoked presentations of the target stimulus.

RESULTS

The individual data are shown in figures 1, 2, 3. The upper trace shows the wave V ER elicited while playing the pong game. The second trace shows the wave V ER elicited while the subject detected auditory signals. Figure 3 shows

Insert Figs. 1, 2 & 3 about here

the above plus an evoked response when the subject was resting. (not instructed to listen to the low intensity signals with the video tube off.) A t-test performed on the mean latencies and amplitudes of Wave V failed to demonstrate statistical significance. However, all amplitudes were somewhat greater in the auditory alone condition when compared to the potentials evoked while playing the pong game.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

It is interesting to note in Figures 1 and 2, subjects 1 and 2, that during the auditory attention task a noticeable post-auricular response occurred following Wave V.

DISCUSSION

Since Hernandez-Peon (1955) demonstrated the inhibition of auditory neural responses by visual distraction in cat, a great deal of effort has been expended in the area of intersensory attentional effects. As mentioned in the introduction the late components of ERP reflect cerebral activity and do exhibit selective attentional effects (Hillyard et al, 1973). However, little is known about the lower levels of the central nervous system, in terms of attentional effects. Lukas (1979) recently reported a reduction in amplitude of brain stem evoked response potentials both in the auditory nerve and the inferior colliculus response to auditory tonal bursts. This reduction was effected by introducing a visual letter display which the subject attended to. The procedure controlled for middle ear effects by using short duration high frequency stimuli. Lukas's results suggest that during concentrated attention to a visual stimuli, irrelevant auditory may be suppressed at a peripheral level possibly through the action of the olivocochlear bundle (Rasmussen, 1939).

The present data lend support to the concept that paying attention to a relevant visual motor task effects a reduction in amplitude (and a slight lengthening of the latency) of the evoked response to an irrelevant auditory stimulus. In the present study only the Wave V of the brainstem evoked response was studied. No attempt was made to evaluate the auditory nerve component. Nevertheless, it seems apparent that the inhibitory effect is present at least by the level of the inferior colliculus.

The results of the present study are very encouraging in terms of providing information concerning the "attentional" process in humans. The study of higher-order processing of information in humans has been hindered by the lack of a broadly applicable model. Understanding the underlying neural processes should provide the necessary information to formulate such a model.

REFERENCES

- Hernandez-Peon, R., and Scherer, H. Federal Proceedings, 1955, 14-71.
- Hillyard, S. A., Hink, R. F. and Picton, T. W. Electrical signs of selective attention in the human brain, Science, 1973, 182, 177-180.
- Lukas, J. H. The effects of attention on the human auditory brainstem potentials. Paper presented at the 19th meeting of the Society for Psychophysiological Research, Cincinnati, OH. 18-21 October, 1979.
- Oatman, Effects of Visual Attention on the Intensity of Auditory Evoked Potentials. Exp. Neurology, 1976, 51, 45-53.
- Rasmussen, G. L. An efferent cochlear bundle. Anatomical Record, 1942, 82, 441.
- Sutton, S., Braren M., Zubin, J. and John, E. R. "Evoked P₃₀₀ Potentials and Variations in Stimulus Probability. Psychophysiology, 1975, 12, 591-595.

FIGURE CAPTIONS

- Figure 1. Brainstem evoked response to an auditory stimulus average of 4096 target stimuli in an odd-ball task. For Subject 1.
- Figure 2. Brainstem evoked response to an auditory stimulus average of 4096 target stimuli in an odd-ball task. For Subjects 2 and 3.
- Figure 3. Brainstem evoked response to an auditory stimulus average of 4096 target stimuli in an odd-ball task. For Subjects 4 and 5. The lower evoked potential is during a rest only control condition.

VISUAL



2 msec

S₁

AUDITORY



2 msec

VISUAL



S₂

AUDITORY



2 msec

VISUAL




S₃

AUDITORY




2 msec

S₄



2 msec

S₅



2 msec