

1-A089 426

LETTERMAN ARMY INST OF RESEARCH SAN FRANCISCO CA F/G 6/20
HEMATOLOGIC AND SELECTED HEPATIC CHANGES PRODUCED BY SUBSTITUTE--ETC(U)
JUL 80 S T OMAE, R A WIRTZ, L S GUTHERTZ
LAIR-TN-14 NL

CLASSIFIED

1 of 1

20-24-70



END
DATE
FILMED
-10-80
DTIC

AD A089426



10

9) TECHNICAL NOTE NO. 14

6) HEMATOLOGIC AND SELECTED HEPATIC CHANGES
PRODUCED BY SUBSTITUTED p-BENSOQUINONES IN THE RAT,

10) S.T./OMAYE, PhD
R.A./WIRTZ, PhD, CPT MS*
L.S./GUTHERTZ, MA
P.C./TAYLOR, BS
J.T./FRUIN, DVM, PhD, LTC VC

11) Jul 11

13) 126

DIVISION OF NUTRITION TECHNOLOGY
and
*DIVISION OF CUTANEOUS HAZARDS


17) LAIR-TN-14

DTIC
ELECTE
SEP 24 1980
S E D

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution unlimited

DDC FILE COPY

JULY 1980



TERMAN ARMY INSTITUTE OF RESEARCH PRESIDIO OF SAN FRANCISCO CALIFORNIA 94129

77112 80 8 11 049

1

Reproduction of this document in whole or in part is prohibited except with the permission of Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, DDC is authorized to reproduce the document for United States Government purposes.

Destroy this report when no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

In conducting the research described in this report, the investigators adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Revision of the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Research Council.

Jm 20 July 80
.....
(Date)

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

HEMATOLOGIC AND SELECTED HEPATIC CHANGES
PRODUCED BY SUBSTITUTED p-BENZOQUINONES IN THE RAT

Accession For	
NTIS GRA&I	
DDC TAB	
Unannounced	
Justification <i>Letter on File</i>	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
<i>A</i>	

Benzoquinones are secreted by a number of tenebrionid flour beetles of which the Tribolium spp. are the most prominent food pests (1). The major components of the odoriferous secretion have been identified as 2-methyl-1,4-benzoquinone (MBQ) and 2-ethyl-1,4-benzoquinone (EBQ) (1-5). Benzoquinones are highly reactive compounds and evidence indicates that they are acutely toxic and perhaps carcinogenic to laboratory animals (6,7). Poisoning by hydroquinone-quinone-systems in man is characterized by jaundice, anemia, hemoglobinuria, and cachexia (8). In previous studies, we found that benzoquinone toxicity in rats resulted in respiratory depression, skin blanching, and cyanosis before death (9,10).

Since respiratory impairment may be due in part to inadequate blood oxygen, we conducted preliminary studies to examine the effects of substituted benzoquinones on the heme in red blood cells and in hepatic microsomal enzymes.

METHODS

All chemicals were of the highest purity available. Agar (No. 0140) was obtained from Difco Laboratories, Detroit, MI. MBQ was obtained from J.T. Baker Chemical Co., Phillipsburg, NJ. EBQ was synthesized following the procedures of Ladisch and Suter (6). The adult male Sprague-Dawley rats (50-days old, weighing 200 ± 50 g) were obtained from Charles River, Wilmington, MA. Upon arrival, each animal was examined for abnormalities and normal rats were randomly housed in stainless steel wire-bottomed cages, one to a cage, and fed a stock diet (Rodent Laboratory Chow #5001, Ralston Purina Co., St. Louis, MO) ad libitum. The rats were divided into three groups of five animals each. The animals in each group were dosed daily with either the aqueous vehicle (0.25% agar), MBQ, or EBQ for four days before killing. The benzoquinones, suspended by trituration in 0.25% agar at concentrations of 90 mg MBQ/5 ml and 110 mg EBQ/5 ml, were administered at the

LD₀₅ level (9,10). Dosage was by oral intubation at a constant volume of 5 ml/kg body weight.

Twenty-four hours after the last dosing the rats were weighed and killed by decapitation. Blood from decapitation was collected into tubes containing EDTA and kept at 4 C for subsequent hematologic measurements. The livers were excised, weighed, perfused with ice-cold 1.15% KCl, and minced. Liver preparation, subcellular fractionation, and hepatic cytochrome-drug assays were done as previously described (11,12). Nonprotein sulfhydryls were measured in whole liver homogenates (13) and glutathione. GSH-s-transferase activity was measured in the post-100,000 xG supernatant (cytosol) (14).

Data were analyzed statistically by the Student's t test or by analysis of variance.

RESULTS

After four days of dosing with substituted p-benzoquinone the rats appeared passive and listless compared to agar-treated controls. Table 1 shows that blood leukocytes were significantly increased (67%) after MBQ or EBQ treatment and that there was also a tendency for the differential lymphocyte to neutrophil ratio to increase compared to controls. In general, Table 1 shows that MBQ and EBQ treatment resulted in fewer erythrocytes (8.4% and 10.7%, respectively) and lower hemoglobin and hematocrits than agar-treated controls (5.6 to 14.8%).

Table 1. Hematology on rats treated with substituted p-benzoquinones*

Compound	WBC x10 ³	RBC 10 ⁶	HGB gm	HCT %	Differential WBC Counts, % Lymphocytes	Neutrophils
Control	10.3 ± 0.8	6.45 ± 0.18	14.3 ± 0.4	39.1 ± 0.8	24 ± 3	70 ± 4
MBQ	17.2 ± 2.1†	5.91 ± 0.26	13.5 ± 0.7	34.7 ± 1.8	36 ± 4†	61 ± 4
EBQ	17.2 ± 1.2†	5.76 ± 0.20†	13.0 ± 0.3†	33.3 ± 1.0†	34 ± 4	63 ± 4

* Mean S.E.M.; N = 5; MBQ = 2-methyl-1,4-benzoquinone; EBQ = 2-ethyl-1,4-benzoquinone; WBC = white blood cells; RBC = red blood cells; HGB = hemoglobin; HCT = hematocrit.

† P < 0.05 compared to control groups.

Table 2 demonstrates that the treatment of rats with substituted p-benzoquinones resulted in no statistical change of body weight, liver weight, or liver microsomal protein. Table 3 shows that MBQ and EBQ

Table 2. Body and liver weight and protein content of liver microsomal protein in rats treated with substituted p-benzoquinones*

Compound	Body Weight g	Liver Weight g	Microsomal Protein mg/g
Control	285 ± 10.0	9.5 ± 0.59	13.78 ± 1.09
MBQ	288 ± 05.0	10.5 ± 0.69	14.87 ± 0.87
EBQ	274 ± 06.6	10.3 ± 0.43	16.68 ± 1.07

* Mean ± S.E.M.; N = 5; MBQ = 2-methyl-1,4-benzoquinone; EBQ = 2-ethyl-1,4-benzoquinone.

Table 3. Liver microsomal cytochrome levels in rats treated with substituted p-benzoquinones*

Compound	Cytochromes P-450	(nmoles/mg protein) b ₅
Control	1.221 ± 0.079	0.351 ± 0.019
MBQ	1.076 ± 0.065	0.270 ± 0.010†
EBQ	0.926 ± 0.106†	0.258 ± 0.024†

* Mean ± S.E.M.; N = 5; MBQ = 2-methyl-1,4-benzoquinone; EBQ = 2-ethyl-1,4-benzoquinone.

† P ± 0.05 compared to control groups.

treatment of rats resulted in decreases of both cytochrome P-450 and cytochrome b₅ hepatic microsomal heme proteins (23.1 to 26.5%). Table 4 documents that the cytochrome P-450 dependent mixed-function oxidase activity of aniline hydroxylase decreased significantly after treatment with substituted p-benzoquinones. Similar trends were also found for aminopyrine demethylase activity after treatment with MBQ and EBQ.

Table 4. Liver microsomal mixed-function oxidase activity in rats treated with substituted p-benzoquinones*

Compound	Aminopurine N-demethylase (nmole/mg protein)	Aniline hydroxylase (nmole/mg protein)
Control	5.164 ± 0.458	0.701 ± 0.041
MBQ	4.718 ± 0.333	0.521 ± 0.031†
EBQ	4.970 ± 0.373	0.402 ± 0.050†

* Mean ± S.E.M.; N = 5; MBQ = 2-methyl-1,4-benzoquinone; EBQ = 2-ethyl-1,4-benzoquinone.

† P < 0.05 compared to control group

In contrast to the suggested mechanism of action (15), Table 5 shows that p-benzoquinones do not reduce liver sulfhydryls in vivo. In addition, MBQ and EBQ had no effect on cytosolic GSH-s-transferase activity.

Table 5. Liver nonprotein sulfhydryls and glutathione-s-transferase activity in rats treated with substituted p-benzoquinones*

Compound	Nonprotein sulfhydryls (nmole/mg protein)	GSH-s-transferase (μmole/mg protein)
Control	3.53 ± 0.152	0.845 ± 0.051
MBQ	4.25 ± 0.514	0.764 ± 0.035
EBQ	3.92 ± 0.186	0.890 ± 0.076

* Mean ± S.E.M.; N = 5; MBQ = 2-methyl-1,4-benzoquinone; EBQ = 2-ethyl-1,4 benzoquinone.

DISCUSSION

Several in vitro reactions have been described as explanation for the p-benzoquinone toxicity to animals. Synthetic EBQ and quinone mixtures isolated from Tribolium castaneum are known to form quinone conjugates with -NH and -SH groups of amino acids (15), proteins (15, 16) and similar compounds. The conjugates with amino acids are intensely colored and often fluorescent (15,17). Irreversible formation of -SH groups seems unlikely since we could find no evidence of loss of total nonprotein sulfhydryls (Table 5) in liver tissues of rats treated with MBQ or EBQ. In addition, there was no influence of substituted p-benzoquinones on GSH-s-transferase activity, a cytosolic enzyme dependent on GSH.

Microsomal heme protein and cytochrome P-450, as well as the corresponding mixed-function oxidase activity and blood heme parameters, seem to be markedly influenced by substituted p-benzoquinone treatment. This is consistent with the hydroquinone-quinone poisoning, characterized in man by anemia and hemoglobinuria (8). Our observations of decreased heme-containing proteins and blood heme after MBQ and EBQ treatments are also consistent with our previous observations of cyanosis, extreme skin blanching, and depressed respiration in rats treated with p-benzoquinone and substituted p-benzoquinones (9,10).

Affinity of benzoquinones for amino acids and proteins may result in decreased synthesis or increased degradation of heme or apoprotein. Ladisch and Suter (16) have also suggested that the affinity of the benzoquinones for protein is responsible for their carcinogenic properties; however, further investigation is needed to determine any resultant alterations of biochemical function which may be responsible for the observed toxicity of these quinones.

SUMMARY

Rats were given daily oral doses of 2-methyl-1,4-benzoquinone (MBQ) or 2-ethyl-1,4-benzoquinone (EBQ) at the LD₀₅ level for four days. There were 5.6 to 14.8% decreases of blood heme parameters in rats treated with substituted p-benzoquinone compared to agar-treated controls. Benzoquinone treatment resulted in 67% increases of blood leukocyte counts. Hepatic microsomal cytochrome P-450 and cytochrome b₅ contents were decreased 11.9 to 26.5% in benzoquinone-treated rats. Aniline hydroxylase activity decreased 25.7 to 42.7%. There was no change in liver weight, microsomal protein, nonprotein sulfhydryls, or GSH-s-transferase activity. The results of this study suggest that substituted p-benzoquinones may have exerted their toxic action in rats by altering heme metabolism.

ACKNOWLEDGEMENTS

The authors thank Mr. D. Darr and Ms. F. Pulliam for their technical assistance and the Division of Research Support for hematologic measurements.

REFERENCES

1. LADISCH, R.K. Quinone toxins and allied synthetics in carcinogenesis. Proc Penn Acad Sci 38:144-149, 1965
2. ALEXANDER, P., and D.H.R. BARTON. The excretion of ethylquinone by the flour beetle. Biochem J 37:463-465, 1943
3. ENGLEHARDT, M., H. RAPOPORT, and A. SOKOLOFF. Odor secretions of normal and mutant Tribolium confusum. Science 150:632-633, 1965
4. WIRTZ, R.A., S.L. TAYLOR, and H.G. SEMEY. Concentrations of substituted p-benzoquinones and 1-pentadecene in the flour beetles Tribolium confusum J. Du val and Tribolium castaneum (Herbst). Comp Biochem Physiol 61:25-28, 1978
5. WIRTZ, R.A., S.L. TAYLOR, and H.G. SEMEY. Concentrations of substituted p-benzoquinones and 1-pentadecene in the flour beetles Tribolium madens (Charp.) and Tribolium brevicornis (LEC.) (Coloptera:Tenebrionidae). Comp Biochem Physiol 61:287-290, 1978
6. LADISCH, R.K. and M.A. SUTER. Teritoxin: synthesis and toxicity rating. Proc Penn Acad Sci 39:42-47, 1966
7. LADISCH, R.K., M.A. SUTER, and G.F. FROIO. Sweat gland carcinoma produced in mice by insect quinones. Proc Penn Acad Sci 42:87-91, 1968
8. SEIFTER, J. General pharmacology and toxicology of quinones and hydroquinones. Biological Antioxidants. Trans. 3rd Conference, Josiah Macy, Jr. Foundation, New York, N.Y., 1948. pp 50-57
9. OMAJE, S.T., R.A. WIRTZ, and J.T. FRUIN. Acute oral toxicity of substituted p-benzoquinones and 1-pentadecene in the male rat. LAIR Technical Note 12, San Francisco, California: Letterman Army Institute of Research, 1979
10. OMAJE, S.T., R.A. WIRTZ, and J.T. FRUIN. Toxicity of selected compounds found in the secretion of the Tenebrionid flour beetles. J Food Safety 2:97-103, 1980
11. OMAJE, S.T., and J.D. TURNBULL. Heme oxygenase activity, drug metabolism and ascorbic acid distribution in the livers of ascorbic acid deficient guinea pigs. Biochem Pharmacol 28:1415-1419, 1979

- 1
12. OMAYE, S.T., and J.D. TURNBULL. Degradation of cytochrome P-450 heme in ascorbic acid-deficient guinea pigs. *Biochem Pharmacol* 28:3651-3658, 1979
 13. SEDLACK, J., and R.H. LINDSAY. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissues with Ellman's reagent. *Anal Biochem* 25:192-205, 1968
 14. PABST, M.J., W.H. HABIG, and J.B. JAKOBY. Glutathione-s-transferase A. *J Biol Chem* 247:7140-7150, 1974
 15. LADISH, R.K. Tetritoxin: binding with amino acids and proteins, *Proc Penn Acad Sci* 39:48-56, 1966
 16. LADISH, R.K., and M.A. SUTER. Affinity of tetritoxin to animal protein. *Proc Penn Acad Sci* 41:206-212, 1967
 17. LADISH, R.K. Assay of tetritoxin in flour beetles *Tribolium* and method of detecting the toxin on cereal foods. *Proc Penn Acad Sci* 39:36-46, 1966

1

OFFICIAL DISTRIBUTION LIST

Commander
US Army Medical Research and
Development Command
Fort Detrick, Frederick, MD 21701

Defense Documentation Center
ATTN: DDC-TCA (12 cys)
Cameron Station
Alexandria, VA 22314

Director of Defense Research and Engineering
ATTN: Asst Dir (Environmental and Life Sciences)
Washington, DC 20301

The Surgeon General
ATTN: DASG-TLO
Washington, DC 20314

Superintendent
Academy of Health Sciences
US Army
ATTN: AHS-COM
Fort Sam Houston, TX 78234

Assistant Dean
Inst & Rsch Support
Uniformed Services University
of Health Sciences
6917 Arlington Road
Bethesda, MD 20014

Commander
US Army Environmental Hygiene Agency
Aberdeen Proving Ground, MD 21070

US Army Research Office
ATTN: Chem & Bio Sci Div
P.O. Box 1221
Research Triangle Park, NC 27709

Biological Sciences Division
Office of Naval Research
Arlington, VA 22217

Director of Life Sciences
USAF Office of Scientific Research (AFSC)
Bolling AFB
Washington, DC 20332

Director
Walter Reed Army Institute of Research
Washington, DC 20012

Commander
US Army Medical Research
Institute of Infectious Diseases
Fort Detrick, Frederick, MD 21701

Commander
US Army Research Institute
of Environmental Medicine
Natick, MA 01760

Commander
US Army Institute of Surgical Research
Brooke Army Medical Center
Fort Sam Houston, TX 78234

Commander
US Army Institute of Dental Research
Washington, DC 20012

Commander
US Army Medical Bioengineering
Research & Development Laboratory
Fort Detrick, Frederick, MD 21701

Commander
US Army Aeromedical Research Laboratory
Fort Rucker, AL 36362

Director
US Army Biomedical Laboratory
Aberdeen Proving Ground
Edgewood Arsenal, MD 21010

Commander
Naval Medical Research Institute
National Naval Medical Center
Bethesda, MD 20014

Commander
USAF School of Aerospace Medicine
Aerospace Medical Division
Brooks AFB, TX 78235