

AD-A091 170

NAVAL AIR DEVELOPMENT CENTER WARMINSTER PA AIRCRAFT --ETC F/G 11/9
REPETITIVE MOLECULAR EXCLUSION CHROMATOGRAPHY OF PGBX ON SEPHAD--ETC(U)
OCT 80 H # SHMUKLER, S F KWONG, W FEELY
NADC-80183-60

NL

UNCLASSIFIED

[OF]
AD-
ACQUISITION



END
DATE
FILMED
12 80
DTIC

REPORT NO. NADC-80183-60

LEVEL II

12

DTIC ELECTE
NOV 3 1980



AD A091170

**REPETITIVE MOLECULAR EXCLUSION CHROMATOGRAPHY
OF PGB_x ON SEPHADEX LH-20**

H. W. SHMUKLER, Ph.D.
S. F. KWONG, M.S.
Biochemistry Research Team
Life Sciences
Aircraft and Crew Systems Technology Directorate
NAVAL AIR DEVELOPMENT CENTER
Warminster, Pennsylvania 18974

W. FEELY, B.S.
M. G. ZAWRYT, B.A.
E. POLIS, M.S.
E. SOFFER, B.A.
HAHNEMANN MEDICAL COLLEGE AND HOSPITAL
Philadelphia, Pennsylvania 19102

PHASE REPORT
AIRTASK NO. F58527003
Work Unit No. SJ202

Approved for Public Release: Distribution Unlimited.

2 OCTOBER 1980

Prepared for
OFFICE OF NAVAL RESEARCH
Department of the Navy
Arlington, Virginia 22217

DDC FILE COPY,

8011 03 215

N O T I C E S

REPORT NUMBERING SYSTEM - The numbering of technical project reports issued by the Naval Air Development Center is arranged for specific identification purposes. Each number consists of the Center acronym, the calendar year in which the number was assigned, the sequence number of the report within the specific calendar year, and the official 2-digit correspondence code of the Command Office or the Functional Directorate responsible for the report. For example: Report No. NADC-78015-20 indicates the fifteenth Center report for the year 1978, and prepared by the Systems Directorate. The numerical codes are as follows:

CODE	OFFICE OR DIRECTORATE
00	Commander, Naval Air Development Center
01	Technical Director, Naval Air Development Center
02	Comptroller
10	Directorate Command Projects
20	Systems Directorate
30	Sensors & Avionics Technology Directorate
40	Communication & Navigation Technology Directorate
50	Software Computer Directorate
60	Aircraft & Crew Systems Technology Directorate
70	Planning Assessment Resources
80	Engineering Support Group

PRODUCT ENDORSEMENT - The discussion or instructions concerning commercial products herein do not constitute an endorsement by the Government nor do they convey or imply the license or right to use such products.

APPROVED BY: _____

DATE: _____

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 NADC-80183-60	2. GOVT ACCESSION NO. AD-A091170	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) REPETITIVE MOLECULAR EXCLUSION CHROMATOGRAPHY OF PGB _x ON SEPHADEX LH-20	5. TYPE OF REPORT & PERIOD COVERED 7 PHASE REPORT	
7. AUTHOR(s) 10 H. W. SHMUKLER, S. F. KWONG M. F. ... E. ...	8. CONTRACT OR GRANT NUMBER(s) 12 J ...	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Aircraft and Crew Systems Technology Directorate Naval Air Development Center Warminster, PA 18974	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS AIRTASK NO. F58527005 Work Unit No. SJ202	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217	12. REPORT DATE 16 OCT 1980	13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report)	
16. DISTRIBUTION STATEMENT (of this Report)		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Approved for Public Release; Distribution Unlimited.		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Repetitive Sephadex LH-20 Molecular Exclusion Chromatography PGB _x		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) PGB _x , a mixture of oligomers of 15-keto PGB ₁ , was subjected to repetitive MEC on Sephadex LH-20 in an attempt to separate and purify the active principle. The resulting fractions were analyzed and found still to be heterogeneous. Although the MEC technique did not resolve the PGB _x complex, low molecular weight fractions were obtained that may be amenable to mass spectral analysis.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6001

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

3151

315

TABLE OF CONTENTS

	Page
LIST OF FIGURES	1
LIST OF TABLES	1
LIST OF ABBREVIATIONS	2
INTRODUCTION	3
MATERIALS AND METHODS	3
RESULTS	
RESULTS	3
DISCUSSION	5
BIBLIOGRAPHY	12

LIST OF FIGURES

Figure

1	Schematic Representation of Methodology for Repetitive MEC of PGB _x on Sephadex LH-29	6
---	--	---

LIST OF TABLES

Table

I	Distribution of PGB _x in Sephadex LH-20 Fractions: 1st MEC	7
II	Distribution of PGB _x in Sephadex LH-20 Fractions: 2nd MEC	8
III	Distribution of PGB _x in Sephadex LH-20 Fractions: 3rd MEC	9
IV	The Weight Percent Distribution of PGB _x in Fractions Separated by Repetitive Sephadex LH-20 MEC.	10
V	A _{245nm} /A _{310nm} of Fractions Separated by Repetitive MEC	11

Accession For TIS GRA&I REPORTS U. S. GOVERNMENT JUSTIFICATION	By Distribution/ Availability Codes AFTR and/or Dist. Status <div style="text-align: center; font-size: 2em; font-weight: bold;">A</div>
--	---

L I S T O F A B B R E V I A T I O N S

MEC - Molecular Exclusion Chromatography

\bar{M}_n - Number Average Molecular Weight

RLM - Rat Liver Mitochondria

Type II PGB_x - Fraction 2 of 1st Sephadex LH-20 MEC

VPO - Vapor pressure osmometry

I N T R O D U C T I O N

PGB_x, a polymeric derivative of 15-keto PGB₁, was first synthesized by Polis et al (1) and shown to conserve oxidative phosphorylation during degradation of isolated rat liver mitochondria (2). More recently Ohnishi and Devlin (3) showed that PGB_x behaved as a water soluble Ca⁺⁺-ionophore in skeletal muscle sarcoplasmic reticulum. Although these properties of PGB_x are significant, the major interest in this compound is its possible role as a therapeutic agent. This is based on the animal studies of (a.) Polis and Angelakos (4) in which PGB_x treated monkeys survived cardiogenic shock, (b.) Polis and Kolata (5) in which PGB_x treated rabbits survived experimentally induced global ischemia, and (c.) Moss et al (6) in which PGB_x treated dogs survived experimentally induced "fatal" hypoxia.

PGB_x prepared according to Polis et al is a mixture of oligomers of varying molecular weight and unknown chemical composition. Before human testing may be undertaken, the active principle in the PGB_x complex must be isolated in pure form and its chemical structure determined. Studies are currently underway in this and other laboratories to attain the above goal. This report describes the use of repetitive MEC of PGB_x on Sephadex LH-20 in order to obtain the PGB_x complex as a narrower molecular weight range of oligomers and thus a purer preparation.

M A T E R I A L S A N D M E T H O D S

The PGB_x used in this study was prepared according to Polis et al (2) and was the crude extract before purification by MEC. When Sephadex LH-20 MEC was carried out, the procedure used was as described before (2). Molecular weights (Mn) were determined by VPO on the free acids dissolved in methanol, using a Wescan Molecular Weight Apparatus (Wescan, Santa Clara, CA) at 60°. Purified fractions were assayed for the in vitro PGB_x effect on the stabilization of oxidative phosphorylation during degradation of RLM by the method of Polis et al (7) as modified by Shmukler et al (8). This modification includes the definition and measurement of the PGB_x K_a and K_i effects. UV absorption spectra of PGB_x fractions were measured in a Cary Recording Spectrophotometer at a concentration of 30ug/ml dissolved in methanol.

R E S U L T S

In the original method for the preparation of PGB_x, Polis et al (2) used Sephadex LH-20 MEC in an attempt to purify the active principles of the PGB_x complex. Although the exclusion limit for Sephadex LH-20 is between 100-500 daltons, no clean chromatographic separations were obtained. Instead the PGB_x complex eluted as one diffuse chromatographic peak. In order to effect some degree of separation, Polis et al arbitrarily collected effluent fractions which were then monitored for their in vitro RLM oxidative phosphorylation effect.

The most active PGB_x preparation, fraction 2¹, gave a \bar{M}_n of 2200-2300 by VPO. Since this fraction was the PGB_x fraction sent to Office of Naval Research contractors for testing, it was of interest to try to purify this fraction even further. As a possible method to accomplish this, repetitive Sephadex LH-20 MEC was tried in order to purify the active principle and/or to obtain a PGB_x preparation with a narrower molecular weight range of components than the Type II PGB_x.

In this study 6620 mg of crude extract of PGB_x, i.e., the NaHCO₃ extract (2) were converted to the free acid, dissolved in methanol and chromatographed on Sephadex LH-20 as described by Polis et al (2). The resulting seven fractions were flash evaporated and analyzed for weight recovery, \bar{M}_n and in vitro PGB_x effects. These results are listed in table I. As seen in this table, fractions 1-2² and fraction 1-3 had approximately the same in vitro PGB_x activity, therefore these 2 fractions were combined in order to have sufficient material for further MEC and subsequent analyses. This combination contained 1932 mg of PGB_x of which 1385 mg were used in the 2nd MEC. Seven fractions were separated and analyzed as above. These results are listed in table II. As seen in this table fractions 2, 3, 4, and 5 all had approximately the same level of PGB_x activity, although the \bar{M}_n values were markedly different. For the 3rd MEC fractions, 2-2 and 2-3 were combined (679 mg) and the analyses of the resulting seven fractions are listed in table III. The separation methodology used in this study is summarized by schematic representation shown in figure 1.

The overall weight recovery of PGB_x in the 1st MEC was 88.5 percent. This suggests that 11.5 percent of the crude extract was either irreversibly adsorbed to the column, or adsorbed strongly to the column to elute outside the PGB_x range. The overall weight recovery of the 2nd and 3rd MEC was approximately quantitative, i.e., 95.7 and 101.9 percent respectively. This suggests that material not recovered in the 1st MEC is not PGB_x.

Chromatography of the crude PGB_x extract, 1st MEC, yields fractions varying in \bar{M}_n from 3049 to 372 as the retention time of the eluted material increased. Rechromatography of fractions 1-2 and 1-3 (2nd MEC) yielded fractions with \bar{M}_n varying from 2873 to 407. Even after 3rd MEC of fractions 2-2 and 2-3, the \bar{M}_n of the fractions varied from 2062-1466. Although the \bar{M}_n for fraction 3-7 could not be measured because of insufficient quantity, it would be reasonable to assume that the \bar{M}_n of fraction 3-7 would be similar to that of fraction 2-7.

The dry weights of all fractions separated in the repetitive MEC listed in tables I, II, and III were used to calculate the percent distribution of PGB_x in each fraction in each separate MEC. These results are listed in table IV. The percent distribution of PGB_x in the 1st MEC is approximately evenly distributed between fractions 1 through 6. On rechromatography of fractions 2-2 and 2-3, 59 percent of the PGB_x was found in fractions 3-2 and 3-3.

¹ Fraction 2 of the 1st Sephadex LH-20 MEC is referred to as Type II PGB_x.

² Fractions separated by repetitive MEC are described by 2 numbers: 1st number refers to MEC No.; 2nd number refers to MEC fraction. Thus 1-2 describes fraction 2 from 1st MEC which is equivalent to Type II PGB_x.

The *in vitro* PGB_x assay data listed in tables I, II and III show that the K_a was highest in fractions 2, 3 and 4 of the 1st MEC. Fraction 1-1 was about 80 percent pure while fractions 1-5, 1-6, and 1-7 were relatively impure. Rechromatography of fractions 1-2 and 1-3 (2nd MEC) showed that the K_a was equally spread throughout fractions 2-2 to 2-6 with fractions 2-1 and 2-7 exhibiting low values. When fractions 2-2 and 2-3 were rechromatographed (3rd MEC), only fractions 3-7 showed a low level of K_a . The results of the K_i distribution in the various fractions were similar to those found for the K_a distribution.

The UV absorption spectra between 200 nm to 400 nm were measured for all fractions separated in this study. In general, all fractions showed absorption maxima at 243 nm and absorption shoulders at 300- 320 nm. The major difference between the UV absorption spectra of the separated fractions was the increase in the 300- 320 nm absorption shoulder with increasing retention time of the fractions. The results are listed in table V in terms of the ratio of the absorbance at 243 nm to absorbance at 310 nm. A comparison of the retention time and the A_{243}/A_{310} shows a progressive decrease (or increase in A_{310}) with increasing retention time. A comparison of the A_{243}/A_{310} of fractions with the same retention time but successive MEC, i.e., fractions 1-2 and fraction 2-2, shows a marked increase in the ratio (or a marked decrease in A_{310}). However, the data for the 2nd and 3rd MEC showed no change in the A_{243}/A_{310} for fractions with the same retention time.

D I S C U S S I O N

The results of repetitive MEC of PGB_x on Sephadex LH-20 shows that this method does not yield homogeneous PGB_x preparations as might have been expected. Instead, the fractions that were separated still appear to be heterogeneous even after 3 MEC analyses. These results suggest that possibly even after additional MEC of fractions 3-2 and 3-3, homogeneous preparations of PGB_x would not be obtained. From these results it is obvious that Type II PGB_x, the preparation currently supplied ONR contractors for *in vitro* and *in vivo* animal studies, is a highly complex mixture of oligomers varying in \bar{M}_n from over 2800 to below 400 daltons. It is interesting to note also that the PGB_x fractions separated did not show a significant increase in the specific activity of the PGB_x, i.e., K_a .

One benefit realized with the repetitive MEC is that the PGB_x fractions separated in MEC #3 must have a narrower range of molecular weight components than found in Type II PGB_x. An additional advantage is the recovery of high K_a activity in relatively low molecular weight fraction, e.g. fraction 3-6 that had the following analytical values; \bar{M}_n , 1466; K_a 0.93, K_i 0.92. It is conceivable that such a low molecular weight preparation may be amenable to high resolution and/or field desorption mass spectral analysis.




Figure 1 - Schematic Representation of Methodology for Repetitive MEC of PGB_x on Sephadex LH-20

NADC-80183-60

Table I
Distribution of PGB_x in Sephadex LH-20
Fractions: 1st MEC

Fraction	\bar{M}_n	Wt (mg)	K_a	K_i
1	3049	767	.82	1.18
2	2554	1107	.97	1.33
3	2137	825	1.02	1.48
4	1706	706	.98	1.36
5	1257	1107	.47	.81
6	915	959	.33	.40
7	572	386	.30	.12
Total Recovery	5858 mg			
%	88.5			

NADC-80183-60

Table II

Distribution of PGB_x in Sephadex LH-20

Fractions: 2nd MEC

Fraction	\bar{M}_n	Wt (mg)	K_a	K_i
1	2873	142	.51	.67
2	2683	315	.92	1.04
3	2351	364	1.06	1.08
4	2190	267	1.11	1.08
5	1919	129	1.08	1.06
6	1541	89	.93	.83
7	407	20	.53	.06
Total Recovery	1326 mg			
%	95.7			

Table III

Distribution of PGB_x in Sephadex LH-20

Fractions: 3rd MEC

Fraction	\bar{M}_n	Wt (mg)	K_a	K_i
1	2862	98.4	.87	.89
2	2364	209.1	1.18	1.18
3	2209	200.3	1.09	1.26
4	2008	115.5	1.06	1.23
5	1886	43.3	1.02	1.26
6	1466	22.9	.93	.91
7		2.43	.17	

Total Recovery 691.9 mg

3 101.9

Table IV

The Weight Percent Distribution of PGB_x in Fractions
Separated by Repetitive Sephadex LH-20 MEC

Fraction	MEC Run		
	#1	#2	#3
1	13.1	10.7	14.2
2	19.9	23.8	30.2
3	4.1	27.5	29.0
4	12.1	20.1	16.7
5	18.9	9.7	6.5
6	16.4	6.7	3.3
7	6.6	1.5	.4

Table V

$A_{\lambda 243\text{nm}}/A_{\lambda 310\text{nm}}$
of Fractions Separated by Repetitive MEC

Fraction	1st MEC	2nd MEC	3rd MEC
1	8.14	10.42	10.22
2	6.85	9.50	9.00
3	5.58	8.08	9.89
4	4.96	7.67	8.60
5	4.12	6.20	7.33
6	3.55	5.05	6.23
7	---	4.33	4.78

B I B L I O G R A P H Y

1. Polis, B. D., Grandizio, A. M. and Polis, E.: Some In vitro and In vivo Effects of a New Prostaglandin Derivative. Neurohumoral and Metabolic Aspects of Injury. Adv. in Exper. Med. and Biol., Plenum Press, New York 33:213-220 (1973).
2. Polis, B. D., Kwong, S., Polis, E., Nelson, G., and Shmukler, H. W.: Studies on PGB_x, A Polymeric Derivative of Prostaglandin B₁: I-Synthesis and Purification of PGB_x. Physiol. Chem. and Physics 11:109 (1979).
3. Ohnishi, S. T. and Devlin, T.: Calcium Ionophore Activity of a Prostaglandin B₁ Derivative (PGB_x). Biochemical and Biophysical Research Communications 89:240 (1979).
4. Angelakos, E. T., Riley, R. L., and Polis, B. D.: Recovery of Monkeys from Cardiogenic Shock After Myocardial Infarction with Ventricular Fibrillation. Effects of PGB_x, Report No. NADC-77308-60 (1977).
5. Kolata, R. J.: The Effect of PGB_x on Neurological Recovery from Cerebral Ischemia in Rabbits. Masters Thesis, Univ. of Penna. Veterinary School (1977).
6. Moss, G.: Immediate Restoration of CNS Autonomic Cardiopulmonary Control. Survival of "Lethal" Cerebral Hypoxia by Treatment with PGB_x. Surgical Forum 29:535B (1978).
7. Polis, B. D., Polis, E. and Kwong, S.: Protection and Reactivation of Oxidative Phosphorylation in Mitochondria by a Stable Free Radical Prostaglandin Polymer (PGB_x). Proc. Nat'l Acad. Sci. 76:1598 (1979).
8. Shmukler, H. W., Soffer, E., Kwong, S. F., Zawryt, M. G., Feely, W., and Polis, E.: Studies on PGB_x: Isolation of a PGB_x with Reduced Inhibitor Content, Report No. NADC-79183-60 (1979).

D I S T R I B U T I O N L I S T

REPORT NO. NADC-80183-60

	<u>No. of Copies</u>
DTIC	12
Naval Research Laboratory, Washington, DC	6
ONR (Code 102IP) (ONRL Doc), Arlington, VA	6
ONR, Biochemistry Program (Code 442), Arlington, VA	6
Naval Medical R&D Command, Bethesda, MD.	1
Naval Medical Research Institute, Bethesda, MD	1
ONR Branch Office, Boston, MA	1
ONR Branch Office, Chicago, IL	1
ONR Branch Office, Pasadena, CA	1
US Army Science & Technology Center, APO San Francisco	1
National Library of Medicine, Bethesda, MD	1
NAMRI, Pensacola, FL	1
Army Research Office, Durham, NC	1
AF Office of Scientific Research, Bolling AFB, Washington, DC	1
Army Medical R & D Command, Washington, DC	1
ONR (Code 200), Arlington, VA	5