

AD-A091 750 TEXAS A AND M UNIV COLLEGE STATION DEPT OF CHEMISTRY

F/G 11/9

POLYMER ACTIVATED CATALYSTS. (U)

POLYMER ACTIVATED CATALYSTS 197
NOV 80 D E BERGBREITER, M S BURSTEN

N00014-77-C-0426

NL

UNCLASSIFIED TR-7


1991

END

DATE _____

FILMED

151



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A091750

12 32

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report #7	2. GOVT ACCESSION NO. AD-A091750	3. RECIPIENT'S CATALOG NUMBER (1)
4. TITLE (and Subtitle) Polymer Activated Catalysts	(12)	5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
7. AUTHOR(s) D. E. Bergbreiter, M. S. Bursten, T. J. Lynch and G. L. Parsons		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Texas A&M University Department of Chemistry College Station, TX 77843	(15)	8. CONTRACT OR GRANT NUMBER(s) N0014-77-C00426
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA	(11)	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS N00014-77-C-0426
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (14) TR-7	LEVEL II	12. REPORT DATE November 1, 1980
		13. NUMBER OF PAGES 25
		15. SECURITY CLASS. (of this Report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) C		
18. SUPPLEMENTARY NOTES Accepted for publication in proceedings of first China, Japan, U.S. symposium on organometallic chemistry.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) polymeric reagent, catalyst activation, organometallic polymer hydrogenation, homogeneous catalysis, ion exchange, chlororhodium(I) tris(triphenylphosphine), silver(I) polystyrene sulfonate, dichlororuthenium(II) tris(triphenylphosphine), hydridorhodium(I) carbonyltris(triphenylphosphine), chlororhodium(I) ethylene-bis(triphenylphosphine), polystyrene bis(cyclopentadienyl)titanium dichloride, +		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Three different ways in which functionalized polymers can activate hydrogenation catalysts will be discussed. In the first example, a sulfonated divinylbenzene-crosslinked polystyrene is converted to a silver(I) salt. This silver(I) containing polymer selectively absorbs triphenylphosphine from solution in the presence of transition metal complexes. As a result of this removal of nonvolatile phosphine ligand, the activity of conventional homogeneous hydrogenation catalysts such as $((C_6H_5)_3P)_3RhCl$ or $((C_6H_5)_3P)_3Rh(CO)H$ can be		

DTIC
NOV 14 1980

DDC FILE COPY

20. Abstract (Cont'd)

Increased in cases where excess free ligand is known to lower catalytic activity. In a second example of polymer activation of hydrogenation catalysts we have shown that the hydrogenation activity of polystyrene supported bis-(cyclopentadienyl)titanium dichloride is significantly altered when organomagnesium reagents instead of organolithium reagents are used to prepare an active catalyst. Rates of olefin isomerization of terminal alkenes to trans-2-alkenes and hydrogenation of terminal alkenes both are much faster with the organomagnesium reduced catalysts. Finally, transition metal complexes can be directly deposited on organometallic polymers to yield polymer supported heterogeneous catalysts whose activity in hydrogenation of terminal alkenes is comparable to that of catalysts supported on inorganic materials.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution	
Availability Codes	
Dist	Avail and/or Special
A	

19. Key Words (cont'd)

isomerization, heterogeneous catalysis, selectivity, polystyrene-palladium(0), polystyryl lithium.

OFFICE OF NAVAL RESEARCH

Contract No. N0014-77-C00426

Technical Report #7

POLYMER ACTIVATED CATALYSTS

by

D. E. Bergbreiter, M. S. Bursten, T. J. Lynch

and G. L. Parsons

Accepted for publication
in proceedings of first
China, Japan, U.S. Symposium on Organometallic Chemistry

Texas A&M University
Department of Chemistry
College Station, Texas 77843

November 1, 1980

Reproduction in whole or in part is permitted for any
purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

POLYMER ACTIVATED CATALYSTS

D. E. Bergbreiter, M. S. Bursten, T. J. Lynch and
G. L. Parsons
Department of Chemistry
Texas A&M University
College Station, TX 77843

Abstract

Three different ways in which functionalized polymers can activate hydrogenation catalysts will be discussed. In the first example, a sulfonated divinylbenzene-crosslinked polystyrene is converted to a silver(I) salt. This silver(I) containing polymer selectively absorbs triphenylphosphine from solution in the presence of transition metal complexes. As a result of this removal of nonvolatile phosphine ligand, the activity of conventional homogeneous hydrogenation catalysts such as $((C_6H_5)_3P)_3RhCl$ or $((C_6H_5)_3P)_3Rh(CO)H$ can be increased in cases where excess free ligand is known to lower catalytic activity. In a second example of polymer activation of hydrogenation catalysts we have shown that the hydrogenation activity of polystyrene supported bis(cyclopentadienyl)titanium dichloride is significantly altered when organomagnesium reagents instead of organolithium reagents are used to prepare an active catalyst. Rates of olefin isomerization of terminal alkenes to trans-2-alkenes and hydrogenation of terminal alkenes both are much faster with the organomagnesium reduced catalysts. Finally, transition metal complexes can be directly deposited on organometallic polymers to yield polymer supported heterogeneous catalysts whose activity in hydrogenation of terminal

alkenes is comparable to that of catalysts supported on inorganic materials.

Introduction

The preparation of polymers capable of binding transition metal compounds which are active homogeneous catalysts has been under development for several years now. Examples of polymer supported catalysts using many types of polymers and many different catalysts have been described and reviewed.^{1,2} The procedure of binding a homogeneous transition metal catalyst to a polymer has several important advantages. Succinctly, the advantages are that a polymer-supported homogeneous catalyst offers the potential of being as selective and reactive as its homogeneous counterpart while still providing the experimental simplicity commonly associated with heterogeneous catalysts. Polymer-supported catalysts are also beset by certain disadvantages. For example, polymer-bound transition metal complexes are often found to be less active than their homogeneous counterparts. Polymer-supported catalysts face disadvantages compared to typical heterogeneous catalysts on refractory supports because of the problems of mass transport and heat transport associated with typically used polymers. In addition, the experimental advantages of polymer-supported catalysts (i.e. their recoverability) are sometimes offset by slow leaching of the active (and often expensive) metal from the polymer.

In this paper, we will be discussing some rather different applications of polymers in catalysis: the alteration of a homogeneous catalyst's specificity on binding to a polymer; the activation of conventional homogeneous catalysts using metal containing polymers; and the use of organometallic polymers as precursors of highly dispersed heterogeneous catalyst species on an organic polymeric support.


Results and Discussion

Polymer-Bound Titanium Alkene Isomerization Catalysts. Typical procedures which emphasize the attachment of a homogeneous catalyst to a polymer proceed on the basis that the catalyst specificity of the polymer-supported catalyst will correspond to that of the homogeneous catalyst even if the activity is slightly altered. However, the potential of polymer-bound catalysts to catalyze reactions different from those catalyzed by their homogeneous counterparts has recently been recognized.³ Here we describe our studies of the reactions of alkenes with reduced titanium species bound to polymers in which a significant alteration in the specificity of the catalyst is observed. In addition, our studies have shown a significant effect on catalytic activity results from changing the organometallic cofactor from an organomagnesium to an organolithium reagent.

The immobilization of bis(cyclopentadienyl)titanium dichloride on a polystyrene matrix was first described by Grubbs and Brubaker⁴ who used this "immobilized" species to prepare an alkene hydrogenation catalyst whose activity was greater than its homogeneous analog. In this report we describe further examples of differences between catalysts derived from similar polystyrene-bound titanium species and their homogeneous analogs. In our studies we have shown that reduction of polystyrene-bound bis(cyclopentadienyl)titanium(IV) dichloride (**1**) with alkylmagnesium halides in ethereal solutions produces an alkene isomerization catalyst. We have also studied alkene hydrogenations using organomagnesium reduced **1** and its soluble analog as a further comparison to the organolithium-titanium(IV) systems. Although the exact nature of the catalytically active species could not be determined, ESR studies did establish that Ti(III) was a component, although not an active catalytic species, of these catalyst mixtures.


Originally we had hoped to prepare a polymeric version of the

titanium hydrometalation catalyst first described by Finkbeiner and Cooper (eq 1).^{5,6} Indeed, while our studies were in progress cata-



lytic hydroalumination of alkenes using the same polymer-bound catalyst we employed was reported.⁷ However, when we attempted reaction 1 with polystyrene-bound bis(cyclopentadienyl)titanium dichloride we saw mainly alkene isomerization and little or no hydrometalation. These alkene isomerization catalysts were prepared either using a 2% divinylbenzene-crosslinked polystyrene support or a 20% divinylbenzene-crosslinked polystyrene support. Addition of excess alkylmagnesium bromide or chloride to either polymeric titanium complex suspended in ether resulted in a change in the color of either polymeric titanium complex from an initial red to black. The rate of the color change varied for different alkylmagnesium halides and corresponded roughly to the efficacy of the resulting isomerization catalyst (*vide infra*). Similar color changes were observed on addition of hydrocarbon solutions of *n*-butyllithium, *sec*-butyllithium, and *tert*-butyllithium. Grignard reagents without β -hydrogens (methyl, benzyl) also produced the same color change. The observed color changes were reversible in that work-up using dilute HCl regenerated the polymers **1**.

Addition of a terminal alkene to the catalyst prepared from **1** resulted in isomerization of the starting alkene (eq 2). This isomer-



ization was successful with 1-butene, 1-pentene, and 1-octene. Internal alkenes such as 2-octene apparently do not further react. Similar behavior was noted in hydrogenation reactions. In long term experiments in which fresh charges of Grignard reagent and starting alkene were periodically added, turnover numbers (total

mmol of alkene isomerized/mmol Ti) as high as 1000 were achieved. Less than 15% of the starting alkene formed alkane in contrast to the homogeneous reaction which produced 70% octane (by protonation on aqueous HCl work-up of an intermediate *n*-octylmagnesium halide) when 1-octene was allowed to react with isopropylmagnesium bromide and bis(cyclopentadienyl)titanium chloride under Finkbeiner and Cooper's conditions.^{5,6}

The relative efficacy of various Grignard reagents for promoting the isomerization was *tert*-butyl > isopropyl > *n*-butyl. Methylmagnesium bromide and benzylmagnesium chloride were ineffective at generating an alkene isomerization catalyst. Typical product mixtures from these isomerizations were: 1-octene; E-2-octene (84%), Z-2-octene (12%), *n*-octane (4%); 1-butene; E-2-butene (72%), Z-2-butene (28%); 1-pentene; E-2-pentene (85%), Z-2-pentene (15%). The starting 1-alkenes were typically completely consumed. Alkyl lithium reagents were ineffective at producing this isomerization catalyst although they are known to form alkene hydrogenation catalysts.⁴

Although we typically used gas chromatography to follow these reactions we also found ¹³C NMR of sealed tubes containing substrate alkene, a Grignard reagent, and polymeric catalyst could conveniently be used to follow these reactions (Figure 1). In these experiments, the polymer and catalyst were not detectable because of the line broadening characteristic of solids. The substrate and product alkenes were, however, easily seen and their absorptions could be easily assigned. We believe this procedure could be generally useful in studying reactions involving polymer-bound catalysts so long as all substrates and reagents can be contained in an NMR tube and as long as the spectral analysis is simple.

The mechanism of alkene isomerization was not studied explicitly. However, a mechanism involving oxidative addition of an allylic C-H bond to titanium may be involved. This process is likely occur-




Figure 1. ^{13}C NMR spectrum of isomerization reaction. Arrows indicate position of resonances characteristic of starting 1-alkenes.


ring to a very limited extent for cyclohexene to give the small amount of benzene observed in attempted cyclohexene hydrogenations. Similar processes have been described for other metals.^{7,8} The alternative process involving "Ti-H" addition and β elimination may also occur. A similar hydride addition-elimination seems likely to be occurring in the metalation reactions employing soluble Cp_2TiCl_2 , isopropylmagnesium bromide and alkenes.⁶

Heterogeneous species such as the polymer-supported catalysts described above are not readily characterized except by analogy to homogeneous systems. Thus, we have been unable to further define the actual catalyst. Nevertheless, our studies show that immobilization of homogeneous catalysts can achieve not only activation and experimental simplicity but also modified catalyst selectivity.

Polymeric Cofactors for Homogeneous Catalysts. A novel application of polymers in catalysis would be to use them to activate a conventional homogeneously catalyzed reaction. Instead of binding the catalyst to the polymer, the polymer in this operation would be used to selectively remove a catalyst inhibitor or poison. For many homogeneous transition metal catalysts the most active form of the catalyst results from a dissociation reaction like eq 3. The most active



form of the catalyst in such a case, ML_{n-1} , is thus formed by an unfavorable equilibrium. Active catalysts may also be generated by oxidative addition of H_2 followed by reductive elimination of HX (eq 4). Decreasing the concentration of either L (eq 3) or HX (eq 4) would increase the concentration of the active catalyst species



ML_{n-1} or ML according to Le Chatelier's Principle. Thus it might be possible to observe increased hydrogenation rates if suitable procedures for removing L or HX from a catalytically active system could be devised.

Our initial goal was to design a functionalized polymer to absorb a ligand L such as triphenylphosphine from an active catalytic system, in hope that the catalysis would then go more rapidly. The metal containing polymers described below meet this objective and do indeed accelerate alkene hydrogenations using some conventional homo-

geneous catalysts.

Other attempts to shift equilibria 3 and 4 have been made. One of the reasons $\text{RhCl}(\text{PPh}_3)_3$ was first supported on a polymer was in the hope of shifting equilibrium 3.^{2,11} It was hoped that the concentration of ML_{n-1} could be increased. Several methods for removal of triorganophosphines from solution have also been tried. In 1970, an increased hydrogenation rate was claimed to be the result of reaction of a dissociated triorganophosphine, causing an equilibrium shift. The claim was not well substantiated.¹² In 1973 Knoth, Gosser and Parshall found that forcing solutions of PPh_3 -containing platinum group compounds through a polyimide membrane separated the products of a catalysis and PPh_3 from the transition metal catalyst. This method requires pressures of several to 100 atmospheres and has also been effective for organometallic synthesis.^{13,14} Shriver and co-workers have used soluble Lewis acids to react with the Lewis base PPh_3 to induce a faster hydrogenation rate.¹⁵⁻¹⁷ However, the major rate increases seen were due to halide exchange or metal hydride formation. Many of the Lewis acids tried reacted with the catalyst and left it inactive. Shriver concluded that a ligand abstracting substance must be chosen with great care so that it will react with PPh_3 but not with the catalyst.

Our proposed scheme to remove triphenylphosphine ligand from a solution of a homogeneous catalyst required functionalized polymers. Specifically, we expected that Lewis acid containing polymers could be prepared which could discriminate between absorption of PPh_3 and a homogeneous transition metal catalyst. If a Lewis acid was supported on an insoluble polymer it would be expected to interact more rapidly with PPh_3 than with the larger transition metal catalyst because the pores in the polymer permit more rapid passage of smaller molecules. It is conceivable that the transition metal catalyst might not interact with the supported Lewis acid at all. The use of a polymer-supported cofactor to increase catalysis rates by absorption of free ligands is a novel application of polymers in transition metal chemis-

try. Such a ligand absorbing polymer would be expected to circumvent many of the side reactions encountered by Shriver.^{15,17}

A polymeric reagent designed to absorb PPh_3 should meet several criteria. The polymer must not be catalytically active of itself, nor should it interact with the active catalyst. It should be easily prepared, preferably from commercially available materials if wide use is ever anticipated. It should be easily handled and, if possible, it should be air stable. The ability to absorb more than one type of ligand would also be advantageous.

Preparation of a polymeric reagent with these characteristics was successfully accomplished using commercially available ion exchange resins which had been exchanged with various metal salts. The ion exchange resin chosen as a support for preparation of ligand-absorbing polymers was Amberlyst 15,¹⁸ a sulfonated polystyrene ($\text{PS-SO}_3\text{H}$). It is 20% crosslinked with divinylbenzene and has a pore size of 50-300Å. As a sphere of 20Å diameter may easily contain a molecule of PPh_3 , there should be little problem with access of PPh_3 to the supported metal ion. Functionalization of $\text{PS-SO}_3\text{H}$ with metal ions was performed by conventional ion exchange techniques, and the efficacy of the resulting polymers for PPh_3 absorption was determined by monitoring the disappearance of PPh_3 from a PPh_3 solution which contained the solid $\text{PS-SO}_3\text{M}$ by ^1H NMR. The observed data for absorption of PPh_3 by several different $\text{PS-SO}_3\text{M}$'s is displayed in Table I. Silver(I)-exchanged $\text{PS-SO}_3\text{H}$ ($\text{PS-SO}_3\text{Ag}$) was clearly the most effective at removing PPh_3 from a 0.04 M solution of PPh_3 in THF; all the detectable PPh_3 was gone after 1 h.

An initial objective of our studies was to prepare a polymer which would be selective and absorb PPh_3 but not an active transition metal catalyst. The ability of $\text{PS-SO}_3\text{Ag}$ to selectively absorb PPh_3 has been shown by UV-visible spectroscopy. In these experiments, a toluene solution of $\text{RhCl}(\text{PPh}_3)_3$ was allowed to react with $\text{PS-SO}_3\text{Ag}$ and $\text{PS-SO}_3\text{H}$. The relative rates of absorption of PPh_3 and $\text{RhCl}(\text{PPh}_3)_3$

Table I. Absorption of Triphenylphosphine (PPh_3) by Metal-ion-exchanged Ion Exchange Resin (Macroreticular Sulfonated Polystyrene, Amberlyst 15).^a

Metal Ion	% PPh_3 Absorbed
Co^{+3}	0
Co^{+2}	10
$\text{Ni}^{+2\text{b}}$	10
$\text{Cu}^{+2\text{b}}$	5
$\text{Cu}^{+1\text{c}}$	5
Ag^{+1}	100 (1h)

^aAbsorption occurring from a solution of PPh_3 , 0.04 M in THF, after 12 h. ^bThis resin also absorbed 10% of the $\text{P}(\text{n-Bu})_3$ which was present in a benzene solution which was initially 0.4 M in $\text{P}(\text{n-Bu})_3$. The solution turns yellow. ^cThe copper was present as the amine complex.

by $\text{PS-SO}_3\text{H}$ and $\text{PS-SO}_3\text{Ag}$ were then determined spectroscopically. All concentrations were typical of catalytic conditions. The PPh_3 absorption was monitored at 262 nm and the $\text{RhCl}(\text{PPh}_3)_3$ absorption was monitored at 417 nm.

The absorbance at 417 nm was found to decrease slowly in the presence of both $\text{PS-SO}_3\text{H}$ and $\text{PS-SO}_3\text{Ag}$; a more rapid decrease was found with $\text{PS-SO}_3\text{Ag}$. To an extent this decrease in absorbance reflects the differing ϵ for $\text{RhCl}(\text{PPh}_3)_3$ and $[\text{RhCl}(\text{PPh}_3)_2]_2$.¹⁹ The dimer $[\text{RhCl}(\text{PPh}_3)_2]_2$ is reported to form slowly in non-polar solvents

try. Such a ligand absorbing polymer would be expected to circumvent many of the side reactions encountered by Shriver.^{15,17}

A polymeric reagent designed to absorb PPh_3 should meet several criteria. The polymer must not be catalytically active of itself, nor should it interact with the active catalyst. It should be easily prepared, preferably from commercially available materials if wide use is ever anticipated. It should be easily handled and, if possible, it should be air stable. The ability to absorb more than one type of ligand would also be advantageous.

Preparation of a polymeric reagent with these characteristics was successfully accomplished using commercially available ion exchange resins which had been exchanged with various metal salts. The ion exchange resin chosen as a support for preparation of ligand-absorbing polymers was Amberlyst 15,¹⁸ a sulfonated polystyrene ($\text{PS-SO}_3\text{H}$). It is 20% crosslinked with divinylbenzene and has a pore size of 50-300Å. As a sphere of 20Å diameter may easily contain a molecule of PPh_3 , there should be little problem with access of PPh_3 to the supported metal ion. Functionalization of $\text{PS-SO}_3\text{H}$ with metal ions was performed by conventional ion exchange techniques, and the efficacy of the resulting polymers for PPh_3 absorption was determined by monitoring the disappearance of PPh_3 from a PPh_3 solution which contained the solid $\text{PS-SO}_3\text{M}$ by ^1H NMR. The observed data for absorption of PPh_3 by several different $\text{PS-SO}_3\text{M}$'s is displayed in Table I. Silver(I)-exchanged $\text{PS-SO}_3\text{H}$ ($\text{PS-SO}_3\text{Ag}$) was clearly the most effective at removing PPh_3 from a 0.04 M solution of PPh_3 in THF; all the detectable PPh_3 was gone after 1 h.

An initial objective of our studies was to prepare a polymer which would be selective and absorb PPh_3 but not an active transition metal catalyst. The ability of $\text{PS-SO}_3\text{Ag}$ to selectively absorb PPh_3 has been shown by UV-visible spectroscopy. In these experiments, a toluene solution of $\text{RhCl}(\text{PPh}_3)_3$ was allowed to react with $\text{PS-SO}_3\text{Ag}$ and $\text{PS-SO}_3\text{H}$. The relative rates of absorption of PPh_3 and $\text{RhCl}(\text{PPh}_3)_3$

Table I. Absorption of Triphenylphosphine (PPh_3) by Metal-ion-exchanged Ion Exchange Resin (Macroreticular Sulfonated Polystyrene, Amberlyst 15).^a

Metal Ion	% PPh_3 Absorbed
Co^{+3}	0
Co^{+2}	10
$\text{Ni}^{+2\text{b}}$	10
$\text{Cu}^{+2\text{b}}$	5
$\text{Cu}^{+1\text{c}}$	5
Ag^{+1}	100 (1h)

^aAbsorption occurring from a solution of PPh_3 , 0.04 M in THF, after 12 h. ^bThis resin also absorbed 10% of the $\text{P}(\text{n-Bu})_3$ which was present in a benzene solution which was initially 0.4 M in $\text{P}(\text{n-Bu})_3$. The solution turns yellow. ^cThe copper was present as the amine complex.

by $\text{PS-SO}_3\text{H}$ and $\text{PS-SO}_3\text{Ag}$ were then determined spectroscopically. All concentrations were typical of catalytic conditions. The PPh_3 absorption was monitored at 262 nm and the $\text{RhCl}(\text{PPh}_3)_3$ absorption was monitored at 417 nm.

The absorbance at 417 nm was found to decrease slowly in the presence of both $\text{PS-SO}_3\text{H}$ and $\text{PS-SO}_3\text{Ag}$; a more rapid decrease was found with $\text{PS-SO}_3\text{Ag}$. To an extent this decrease in absorbance reflects the differing ϵ for $\text{RhCl}(\text{PPh}_3)_3$ and $[\text{RhCl}(\text{PPh}_3)_2]_2$.¹⁹ The dimer $[\text{RhCl}(\text{PPh}_3)_2]_2$ is reported to form slowly in non-polar solvents

at these concentrations.¹⁹⁻²² It was expected that dimer formation would be accelerated by PS-SO₃Ag if this polymer were removing PPh₃. The determination of relative rates for rhodium(I) absorption versus PPh₃ absorption was therefore complicated by this dimerization reaction which had the effect of reducing the apparent selectivity of PS-SO₃Ag since dimer formation was not distinguished from RhCl(PPh₃)₃ absorption by UV-visible spectroscopy. Nevertheless, it was possible to estimate that PPh₃ absorption was at least 10-100 times faster than rhodium(I) absorption by this technique.

Having prepared the desired type of functionalized polymer, we next set out to use it to accelerate catalytic reactions. However, reactions in which RhCl(PPh₃)₃ only, RhCl(PPh₃)₃ and PS-SO₃H, or RhCl(PPh₃)₃ and PS-SO₃Ag were used to hydrogenate the alkenes 1-octene, 1-hexene, cyclohexene, styrene, ethylacrylate and 1,5-cyclooctadiene (1,5-cyclooctadiene was not hydrogenated) all occurred at the same rate (Table II). Reaction rates were obtained by monitoring the uptake of H₂ and plotting the data obtained as mmol of H₂ consumed/s. The rate was determined by fitting a straight line to this data and taking its slope. In the literature RhCl(PPh₃)₃-catalyzed alkene hydrogenation rates have been reported to be reproducible to $\pm 10\%$.^{10,21,23,24} James states that this reproducibility refers only to the same catalyst sample and solvent batch.²⁵ Because of these facts, a rate variation of $\pm 10\%$ was not considered significant unless it was consistently repeated. The ineffectiveness of PS-SO₃Ag under catalytic conditions apparently results from the inability of PS-SO₃Ag to compete with other species in solution for free PPh₃. However, PS-SO₃Ag is capable of PPh₃ absorption under hydrogenation conditions. This was shown by hydrogenation reactions run in the presence of excess PPh₃. When 0.01 to 0.02 M PPh₃ was present in hydrogenations of styrene, 1-octene, and cyclohexene, the rates of hydrogenation were from 1 to 13% of their uninhibited rates (cf. Figure 2). Addition of PS-SO₃Ag produced a dramatic increase in the rate of hydrogenation after an induction period of 350-1500 s; most

Table II. Hydrogenation of Alkenes by $\text{RhCl}(\text{PPh}_3)_3$ in the Presence of $\text{PS-SO}_3\text{Ag}$ and $\text{PS-SO}_3\text{H}$.^a

Alkene	Cofactor	$[\text{RhCl}(\text{PPh}_3)_3] \times 10^3, \text{ M}$	Rate $\times 10^3$ mmol H_2/s
Cyclohexene ^b	Nothing	2.23	1.50
Cyclohexene ^b	$\text{PS-SO}_3\text{H}$	2.11	1.30
Cyclohexene ^b	$\text{PS-SO}_3\text{Ag}$	2.21	1.48
1-Hexene ^c	nothing	3.01	1.00
1-Hexene ^c	$\text{PS-SO}_3\text{H}$	2.78	0.94
1-Hexene ^c	$\text{PS-SO}_3\text{Ag}$	2.89	1.04
Styrene ^d	Nothing	2.3	2.2
Styrene ^d	$\text{PS-SO}_3\text{Ag}$	2.3	2.2
Ethyl Acrylate ^e	Nothing	2.84	9.8
Ethyl Acrylate ^e	$\text{PS-SO}_3\text{Ag}$	2.73	9.8

^aRun in 10 mL of toluene at 25°C. 0.20 g of $\text{PS-SO}_3\text{H}$ or $\text{PS-SO}_3\text{Ag}$ was used except as noted. ^b0.47 M. ^c0.20 M. ^d0.11 M. ^e0.23 M.

induction periods were about 700–800 s. The final hydrogenation rate achieved was comparable to the rate observed in the absence of any added PPh_3 ; when more alkene was injected after all the alkene initially present was consumed (about 2000 s), the hydrogenation rate was the same as that observed in the absence of any added PPh_3 . By this time, virtually all of the excess PPh_3 had been adsorbed.

The polymeric reagent $\text{PS-SO}_3\text{Ag}$ clearly removes PPh_3 from organic solutions. The reason no catalyst activation was observed in Table II is apparently because $\text{PS-SO}_3\text{Ag}$ cannot compete with the other species in solution for free PPh_3 during a $\text{RhCl}(\text{PPh}_3)_3$ catalyzed hydrogenation. Therefore, if it were possible to "trick" $\text{RhCl}(\text{PPh}_3)_3$ into releasing PPh_3 while preventing dimerization, an increased hy-

drogenation rate might be observed.

This "trick" is indeed possible. The ethylene complex of $\text{RhCl}(\text{PPh}_3)_3$, $\text{RhCl}(\text{PPh}_3)_2(\text{C}_2\text{H}_4)$ is known to form reversibly on exposure






Figure 2. Hydrogenation of styrene (0.11 M) catalyzed by $2.3 \times 10^{-3} \text{ M}$ $\text{RhCl}(\text{PPh}_3)_3$ in toluene at 25.0° in the absence of both $\text{PS-SO}_3\text{Ag}$ and any excess PPh_3 (○), in the presence of $\text{PS-SO}_3\text{Ag}$ (●), and in the presence of excess PPh_3 ($1.6 \times 10^{-2} \text{ M}$ (⊗)). The effect of rate acceleration in the latter experiment reflects addition of $\text{PS-SO}_3\text{Ag}$ at 5400 s (arrow in Figure).

of toluene solutions of $\text{RhCl}(\text{PPh}_3)_3$ to ethylene (eq 5).²⁶ If $\text{RhCl}-$



$(\text{PPh}_3)_3$ were dissolved in the presence of $\text{PS-SO}_3\text{Ag}$ under an ethylene atmosphere, absorption of the PPh_3 released should drive eq 6 to the right, while still avoiding dimer formation. Subsequent addition of an alkene (after 900 s of mixing), removal of ethylene after 100 s by degassing, addition of hydrogen, and stirring then initiated the hydrogenation reaction. This procedure was found to have no effect on the rates of hydrogenation of styrene or ethyl acrylate; these alkenes were the most rapidly hydrogenated alkenes used in this study. However, 1-hexene, cyclohexene, and ethylene were observed to react



faster after ethylene pretreatment. Control experiments employing ethylene pretreatment in the absence of any cofactor, ethylene pretreatment in the presence of $\text{PS-SO}_3\text{H}$, and no ethylene pretreatment or cofactor present, all resulted in the same unactivated rates.

After an ethylene-pretreated hydrogenation of 1-hexene in the presence of $\text{PS-SO}_3\text{Ag}$, the $\text{PS-SO}_3\text{Ag}$ residue was extracted with MeOH/HNO_3 . In this experiment the residue was rinsed three times with CH_2Cl_2 to remove any PPh_3 which might be clinging to the surface before extraction by acidified MeOH . An absorbance maximum at 262 nm was observed by UV spectroscopy; this was the same as the maximum for PPh_3 in this solution. A spectrum of the acidified methanol-extract from a sample of $\text{PS-SO}_3\text{H}$ showed no absorbance in this region. Thus $\text{PS-SO}_3\text{Ag}$ does absorb PPh_3 when ethylene pretreatment is employed.

Further evidence that the concentration of $\text{RhCl}(\text{PPh}_3)_2(\text{alkene})$ has been increased by this ethylene pretreatment is found by comparing

the activations obtained in this study to work reported by Wilkinson in 1968.²⁷ The catalyst $\text{RhCl}(\text{PPh}_3)_n$ ($n = 2, 3$) was formed in situ by combination of varying ratios of PPh_3 and $[\text{RhCl}(\text{COD})]_2$ with alkene under H_2 . These reactions were run at rhodium and alkene concentrations comparable to those used in this work; a rate enhancement of 1.4 for hydrogenation of 1-hexene was observed for $n = 2$ as opposed to $n = 3$. This compares with the twofold rate enhancement seen in this work following ethylene activation in presence of $\text{PS-SO}_3\text{Ag}$, as opposed to an unmodified hydrogenation using $\text{RhCl}(\text{PPh}_3)_3$. The similarity of these results suggests that the concentration of $\text{RhCl}(\text{PPh}_3)_2$ is indeed increased when ethylene pretreatment is employed in presence of $\text{PS-SO}_3\text{Ag}$.

The idea of shifting equilibrium 3 or equilibrium 4 using a polymeric reagent should be generally applicable. As was described above $\text{PS-SO}_3\text{Ag}$ is a suitable species for absorption of PPh_3 . This polymer was therefore also examined as a means of activating other common homogeneous catalysts.

In the absence of any cofactors, $\text{RuCl}_2(\text{PPh}_3)_3$ is not a good alkene hydrogenation catalyst for 1-hexene or cyclohexene in toluene.¹⁰ Hydrogen uptake occurred very slowly, at a rate of about 4×10^{-4} mmol H_2/s . However, if $\text{PS-SO}_3\text{Ag}$, $\text{PS-SO}_3\text{H}$, neutral alumina, $\text{PS-CH}_2\text{NMe}_3(\text{OH})$ (Dowex 21K) or DMF were added to the solution containing $\text{RuCl}_2(\text{PPh}_3)_3$, the solution gradually turned from brown to red over a period of about 200 s. The hydrogenation rate increased dramatically; H_2 uptake occurred up to 15 times faster after the introduction of the cofactor (Table III). However, the use of polyethylene or macroreticular styrene did not lead to any rate enhancement or color change. Separate experiments showed each cofactor which produced rate enhancements to be capable of absorbing HCl . Only $\text{PS-SO}_3\text{Ag}$ absorbed PPh_3 .

The increase in the alkene hydrogenation rate which followed the addition of an HCl absorbing cofactor to a solution of $\text{RuCl}_2(\text{PPh}_3)_3$

Table III. Acceleration of 1-Hexene Hydrogenation by $\text{RuCl}_2(\text{PPh}_3)_3$ Using Various Cofactors.^a

Rate $\times 10^3$, mmol H_2/s	Cofactor
0.36	Nothing
5.6	$\text{PS-SO}_3\text{Ag}$
2.7	$\text{PS-SO}_3\text{H}$
2.1	$\text{PS-SO}_3\text{Na}$
1.2	$\text{PS-CH}_2\text{H}(\text{CH}_3)_3(\text{OH})$
5.5	Neutral alumina
1.8	Polyethylene
0.24	Polystyrene,
0.26	macroreticular

^aIn toluene, 10 mL, at 25°; [1-hexene] = 0.73 M; $[\text{RuCl}_2(\text{PPh}_3)_3] = 3.6 \times 10^{-3}$ M. The rates were determined from data collected from 200-800 s.

and 1-hexene under H_2 is consistent with conversion of at least some of the $\text{RuCl}_2(\text{PPh}_3)_3$ present to $\text{RuHCl}(\text{PPh}_3)_3$ by absorption of HCl (HX in eq 4). The hydrogenation rates obtained in this work are almost an order of magnitude smaller than those obtained for dilute ($<10^{-3}$ M) solutions of $\text{RuHCl}(\text{PPh}_3)_3$.^{10,29} The relatively depressed hydrogenation rates observed in this work are consistent with the incomplete conversion of $\text{RuCl}_2(\text{PPh}_3)_3$ to $\text{RuHCl}(\text{PPh}_3)_3$ as was shown by subsequent ^{31}P [^1H] NMR experiments.

When $\text{PS-SO}_3\text{Ag}$ was present in a solution of $\text{RuCl}_2(\text{PPh}_3)_3$ and 1-hexene, the initial hydrogenation rate was about two and a half times faster than when $\text{PS-SO}_3\text{H}$ was present. Although these rates tended to decrease after about 1000 s, this may be due to alkene consumption rather than catalyst deactivation. Hydrogenation rates

in the presence of $\text{PS-SO}_3\text{H}$ and $\text{RuCl}_2(\text{PPh}_3)_3$ also tended to decrease after about 3.8 mmol (48%) of the 1-hexene initially present was consumed. This rate differential suggests that $\text{PS-SO}_3\text{Ag}$ is a more efficient HCl absorber than $\text{PS-SO}_3\text{H}$, or that the PPh_3 absorption occurring in the $\text{PS-SO}_3\text{Ag}$ experiment may be important. It is thus possible that the presence of $\text{PS-SO}_3\text{Ag}$ may activate $\text{RuCl}_2(\text{PPh}_3)_3$ by absorption of both HCl and PPh_3 . This conclusion was supported by the results of GC analysis of a 1-hexene hydrogenation using $\text{RuCl}_2(\text{PPh}_3)_3$ as the catalyst precursor and either $\text{PS-SO}_3\text{Ag}$ or $\text{PS-SO}_3\text{H}$ as the cofactor which was stopped after 500 s. The reaction mixture was analyzed for hexane and hexenes. When $\text{PS-SO}_3\text{H}$ was the cofactor, 83.3% of the 1-hexene initially added had been converted to hexane; 15.6% was still 1-hexene, 0.01% was 3- or trans 2-hexene, and 1.1% was cis 2-hexene. However, when $\text{PS-SO}_3\text{Ag}$ was the cofactor, more isomerization had occurred. The six-carbon species were distributed: hexane, 79.1%; 1-hexene, 17.0%; 3- and trans 2-hexene, 2.0%, and cis 2-hexene, 1.8%. While isomerization is slight in each case, there is approximately three times as much isomerization when $\text{PS-SO}_3\text{Ag}$ is used as the cofactor. The great selectivity exhibited by $\text{RuHCl}(\text{PPh}_3)_3$ for 1-alkene hydrogenation is considered to be a result of steric hindrance from the three PPh_3 's associated with the molecule. This also makes isomerization difficult as there is not enough room to accommodate an internal alkene easily. Thus, removal of PPh_3 from solution would be predicted to increase the isomerizing ability of this catalyst, as it appears to have done. A related experiment was the hydrogenation of an internal olefin, cyclohexene, in presence of $\text{RuCl}_2(\text{PPh}_3)_3$ and $\text{PS-SO}_3\text{H}$ or $\text{PS-SO}_3\text{Ag}$. As stated above, $\text{RuHCl}(\text{PPh}_3)_3$ is not an efficient hydrogenation catalyst for internal olefins.²⁹ Formation of $\text{RuHCl}(\text{PPh}_3)_2$ might be expected to result in a faster rate for cyclohexene hydrogenation. The addition of $\text{PS-SO}_3\text{H}$ to a solution of $\text{RuCl}_2(\text{PPh}_3)_3$ and cyclohexene under H_2 led to a five-fold rate increase for the extremely slow hydrogenation. The addition of $\text{PS-SO}_3\text{Ag}$ resulted in a rate nine times faster than that observed in the absence

of any cofactor; nonetheless, the rate was still extremely slow. Although there are numerous other catalysts which are more efficient for this type of hydrogenation, absorption of PPh_3 by $\text{PS-SO}_3\text{Ag}$ does seem to be significant in inducing a faster hydrogenation rate for an internal olefin.

Dissociation of phosphine is regarded as essential to the mechanism generally accepted for many homogeneous hydrogenations of alkenes which are catalyzed by phosphine-containing transition metal compounds.¹⁰ Thus several other common hydrogenation catalysts were surveyed to look for possible activation in the presence of $\text{PS-SO}_3\text{Ag}$. A brief account of these reactions is provided below.

Hydrogenations using $\text{RhH(CO)(PPh}_3)_3$ ³⁰⁻³² were run as follows: the catalyst was dissolved in toluene or THF in presence of $\text{PS-SO}_3\text{Ag}$ and stirred for a total of 1200 s, at which point alkene was injected. Rates of hydrogenation of 1-hexene, 0.31 M, were the same regardless of the presence of $\text{PS-SO}_3\text{Ag}$. This was true at catalyst concentrations of both 1.3×10^{-3} M and 6.0×10^{-3} M. However, styrene (0.2 or 0.6 M) is hydrogenated about 10 times more slowly than 1-hexene.³³ It was hydrogenated about 30-60% faster in presence of $\text{PS-SO}_3\text{Ag}$ when $\text{RhH(CO)(PPh}_3)_3$ concentrations of 3×10^{-3} M to 10×10^{-3} M were employed; however, at 0.9×10^{-3} M in $\text{RhH(CO)(PPh}_3)_3$, no rate enhancement was observed.


The major interaction of $\text{RuCl}_2(\text{PPh}_3)_3$ with $\text{PS-SO}_3\text{Ag}$ and $\text{PS-SO}_3\text{H}$ under homogeneous hydrogenation conditions was HCl absorption by the polymer (vide supra). A similar system, $\text{RuH(OAc)(PPh}_3)_3$, which was not susceptible to this reductive elimination, was therefore chosen so that PPh_3 absorption could be studied.³⁴

Hydrogenations of 1-hexene in THF were observed to occur 1.4 times more quickly in presence of $\text{PS-SO}_3\text{Ag}$. Addition of $\text{PS-SO}_3\text{H}$ has been observed to result in a slower rate of hydrogenation, 0.2 times the rate observed in the absence of any cofactor. This deactivation may have resulted from attack on the carboxyl group by the supported



acid. The question was not pursued. The presence of PS-SO₃Ag also led to an increase in the rate of styrene hydrogenation catalyzed by RhH(OAc)(PPh₃)₃ in THF, by a factor of 1.5 to 1.8. Cyclohexene was not detectably hydrogenated by RhH(OAc)(PPh₃)₃ in the presence or the absence of PS-SO₃Ag.

In summary, the polymer PS-SO₃Ag has been shown to be effective at inducing a rate acceleration for alkene hydrogenation in many PPh₃-containing systems. It was not effective at inducing a rate acceleration in all hydrogenations studied. Selective absorption of either L or HX in eq 3 and 4 was demonstrated. It seems likely that other suitably designed polymers could also be prepared which might demonstrate these effects. Alteration of the polymer's functionality or structure could increase the activity and selectivity of these polymeric cofactors.

Dispersed Metals on Polymers. In recent work we have succeeded in developing routes to a variety of organometallic polymers containing reactive groups.^{35,36} These organometallic polymers or those developed by other groups provide a way of preparing new organometallic polymers from the catalytically active Group VIII metals according to eq 7. When reactive organometallic polymers such as lithiated



polystyrene are allowed to react with palladium(II) salts in reactions like 8 and 9, organometallic polymers are expected to be formed which are functionalized with highly dispersed palladium complexes. Subsequent thermolysis or reduction can transform these palladium(II) species into palladium(0) which can then be used as active alkene and alkyne hydrogenation catalysts. Palladium(0) aggregate size variations and surface areas for the resulting heterogeneous catalysts may



be determined from electron micrographs. Two representative micrographs are shown in Figure 3.

The activity of these heterogeneous catalysts toward terminal and internal alkenes is comparable to palladium-on-carbon catalysts as shown by the data in Table IV. Greater selectivity than Pd/C for terminal versus internal alkene hydrogenation was seen. Alkynes were




Figure 3. Electron micrographs of polystyrene-palladium(0). Metal aggregates range in size from 10-40Å and are localized along the polymer pore surface.

preferentially hydrogenated in the presence of alkenes, and as a result, alkenes could be isolated in high yield after the addition of one equiv of hydrogen. Alkene isomerization was competitive with hydrogenation in cases where more substituted alkenes could be formed. For example, after the partial hydrogenation of 1-hexene, the reaction solution contained 1-, 2-, and 3-hexene and n-hexane.

Table IV. Hydrogenation Reactions Using Polystyrene-Palladium(0).^a

Catalyst ^b	Hydrogenation Rate (mmol H ₂ /min-mmol Pd)	Substrate
(<u>n</u> -Bu ₃)PdCl ₂ (0.4% Pd)	2.4	1-Octene
"	0.4	Cyclohexene
(COD)PdCl ₂ (1.26% Pd)	6.1 ^c	1-Octene
"	6.7	Cyclohexene
"	34.7	1-Octyne
1% Pd/C	26.8	1-Octene
"	20.8	Cyclohexene
"	22.8	1-Octyne
5% Pd/C	35.8	1-Octene
"	14.0	Cyclohexene
"	79.0	1-Octyne

^aHydrogenations were performed at 1 atm pressure and room temperature with 0.05-0.5 g of catalyst and 10 mL of THF which was 0.2 M in substrate. Rates were followed using a gas buret. ^bThe indicated palladium complexes were allowed to react with lithiated polystyrene and reduced. ^cWhole polymer beads. Ground beads hydrogenate octene at 48.7 mmol H₂/min-mmol Pd.

Acknowledgement. We thank the Office of Naval Research, the

Department of Energy, and the Texas A&M Center for Energy and Mineral Resources for support of this research.

1. James, B. R. Adv. Organomet. Chem., 1979, 17, 319-405.
2. Chauvin, Y.; Commereuc, D.; Dawans, F. Prog. Polym. Sci., 1977, 5, 95-226.
3. Perkins, P.; Vollhardt, K. P. C. J. Am. Chem. Soc., 1979, 101, 3985.
4. Grubbs, R. H.; Gibbons, C.; Kroll, L. C.; Bonds, Jr., W. D.; and Brubaker, Jr., C. H. J. Am. Chem. Soc., 1973, 95, 2373; Bonds, Jr., W. D.; Brubaker, Jr., C. H.; Chandrasekaran, E. S.; Gibbons, C.; Grubbs, R. H.; and Kroll, L. C. J. Am. Chem. Soc., 1975, 97, 2128.
5. Cooper, G. D.; Finkbeiner, H. L. J. Org. Chem., 1962, 27, 1493.
6. Finkbeiner, H. L.; Cooper, G. D. J. Org. Chem., 1962, 27, 3395.
7. Ashby, E. C.; Noding, S. A. J. Org. Chem., 1979, 44, 4364.
8. Bingham, D.; Hudson, B.; Webster, D.; Wells, D. B. J. Chem. Soc., Dalton, 1974, 1521.
9. Harrod, J. F.; Chalk, A. J. J. Am. Chem. Soc., 1966, 88, 3491.
10. James, B. R. "Homogeneous Hydrogenation;" John Wiley and Sons: New York, 1973.
11. Pittman, Jr., C. U.; Wilemon, G. Ann. N. Y. Acad. Sci., 1980, 333, 67-73, and references therein.
12. Hidai, M.; Kuse, T.; Hidita, T.; Uchida, Y.; Misono, A. Tetrahedron Lett., 1970, 1715-7.
13. Gosser, L. W.; Knoth, W. H.; Parshall, G. W. J. Am. Chem. Soc., 1973, 95, 3436-7.
14. Gosser, L. W.; Knoth, W. H.; Parshall, G. W. J. Mol. Catal., 1977, 2, 253-63.
15. Strauss, S. H.; Shriver, D. F. Inorg. Chem., 1978, 17, 3069-74.
16. Strauss, S. H.; Diamond, S. E.; Mares, F.; Shriver, D. H. Inorg.

- Chem., 1978, 17, 3064-8.
17. Porter, R. A.; Shriver, D. F. J. Organometal. Chem., 1975, 90, 41-7.
 18. Chemical Dynamics Company, "Chemalog" Catalog. South Plainfield, N. J., 1977/78; p. 387.
 19. Arai, H.; Halpern, J. J. Chem. Soc., Chem. Commun., 1971, 1571-2. The ϵ given for the compound " $\text{RhCl}(\text{PPh}_3)_2$ " is one half the value it should be as the species is really $[\text{RhCl}(\text{PPh}_3)_2]_2$.
 20. Halpern, J.; Wong, C. S. J. Chem. Soc., Chem. Commun., 1973, 629-30.
 21. Tolman, C. A.; Meakin, P. Z.; Lindner, D. L.; Jesson, J. P. J. Am. Chem. Soc., 1974, 96, 2742-4.
 22. Ohtani, Y.; Fujimoto, M.; Yamagishi, A. Bull. Chem. Soc. Jpn., 1976, 49, 1871-3.
 23. Osborn, J. A.; Jardine, F. H.; Young, J. F.; Wilkinson, G. J. Chem. Soc., (A) 1966, 1711-32.
 24. Candlin, J. P.; Oldham, A. R. Faraday Discuss. Chem. Soc., 1968, 46, 60-71.
 25. Ref. 10, p. 208.
 26. Meakin, P.; Jesson, J. P.; Tolman, C. A. J. Am. Chem. Soc., 1972, 94, 3240-2.
 27. Montelatici, S.; van der Ent, A.; Osborn, J. A.; Wilkinson, G. J. Chem. Soc., (A) 1968, 1054-8.
 28. The actual active catalyst is $\text{RuHCl}(\text{PPh}_3)_3$ in such hydrogenations (cf. ref. 29).
 29. Hallman, P. S.; McGarvey, B. R.; Wilkinson, G. J. Chem. Soc., (A) 1968, 3143-50.
 30. Evans, D.; Yagupsky, G.; Wilkinson, G. J. Chem. Soc., (A) 1968, 2660-5.

31. Dewhirst, K. C.; Kein, W.; Reilly, C. A. Inorg. Chem., 1968, 7, 546-51.
32. Yagupsky, M.; Wilkinson, G. J. Chem. Soc., (A) 1970, 941-4.
33. O'Connor, C.; Wilkinson, G. J. Chem. Soc., (A) 1968, 2665-71.
34. Hoffman, P. R.; Caulton, K. G. J. Am. Chem. Soc., 1975, 97, 4121-8.
35. Bergbreiter, D. E.; Killough, J. M. J. Chem. Soc., Chem. Commun., 1980, 319.
36. Organometallic polymers derived from polyethylene like those described in reference 35 may also be prepared. Y. F. Liang, unpublished results.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 356B

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. T. C. Williams Union Carbide Corporation Chemical and Plastics Tarrytown Technical Center Tarrytown, New York	1	Douglas Aircraft Company Attn: Technical Library CI 290/36-84 AUTO-Sutton 3855 Lakewood Boulevard Long Beach, California 90846	1
Dr. R. Soulen Contract Research Department Pennwalt Corporation 900 First Avenue King of Prussia, Pennsylvania 19406	1	NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135	1
Dr. A. G. MacDiarmid University of Pennsylvania Department of Chemistry Philadelphia, Pennsylvania 19174	1	Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375	1
Dr. C. Pittman University of Alabama Department of Chemistry University, Alabama 35486	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. H. Allcock Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1	Dr. E. Fischer, Code 2853 Naval Ship Research and Development Center Annapolis Division Annapolis, Maryland 21402	1
Dr. M. Kenney Case-Western University Department of Chemistry Cleveland, Ohio 44106	1	Dr. Martin H. Kaufman, Head Materials Research Branch (Code 4542) Naval Weapons Center China Lake, California 93555	1
Dr. R. Lenz University of Massachusetts Department of Chemistry Amherst, Massachusetts 01002	1	Dr. J. Magill University of Pittsburg Metallurgical and Materials Engineering Pittsburg, Pennsylvania 22230	1
Dr. M. David Curtis University of Michigan Department of Chemistry Ann Arbor, Michigan 48105	1	Dr. C. Allen University of Vermont Department of Chemistry Burlington, Vermont 05401	1
Dr. M. Good Division of Engineering Research Louisiana State University Baton Rouge, Louisiana 70803	1	Dr. D. Bergbreiter Texas A&M University Department of Chemistry College Station, Texas 77843	1

TECHNICAL REPORT DISTRIBUTION LIST, 356BNo.
Copies

Professor R. Drago
Department of Chemistry
University of Illinois
Urbana, Illinois 61801

1

Dr. F. Brinkman
Chemical Stability & Corrosion
Division
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

1

Professor H. A. Titus
Department of Electrical Engineering
Naval Postgraduate School
Monterey, California 93940

1

COL R. W. Bowles, Code 100M
Office of Naval Research
800 N. Quincy Street
Arlington, Virginia 22217

1

Professor T. Katz
Department of Chemistry
Columbia University
New York, New York 10027

1

Professor James Chien
Department of Chemistry
University of Massachusetts
Amherst, Massachusetts 01002

1

Professor Malcolm B. Polk
Department of Chemistry
Atlanta University
Atlanta, Georgia 30314

1

58

