

AD-A092 386

MARYLAND UNIV. BALTIMORE DEPT OF MATHEMATICS
DETECTION, ESTIMATION, AND CONTROL ON GROUP MANIFOLDS.(U)
JUL 80 J T LO

F/G 12/1

UNCLASSIFIED

AFOSR-TR-80-1055

F49620-79-C-0186
NL

110
100
100



END
DATE
FILMED
1-81
DTIC

AFOSR-TR-80-1055

LEVEL

12

AD A092386

Final Scientific Report, 1 Jan 79-31 Jan 80
on
Detection, Estimation, and Control
on Group Manifolds,

DTIC
SELECTED
DEC 3 1980

TO: Air Force Office of Scientific Research
Air Force Systems Command
USAF

FROM: James T. Lo
Department of Mathematics
University of Maryland Baltimore County
Baltimore, Maryland 21228

SUBJECT: Contract No. F49620-79-C-0186
Detection, Estimation, and Control
on Group Manifolds

covering the period:
July 1, 1979 - July 31, 1980

DDC FILE COPY

Approved for public release;
distribution unlimited.

80 1201 082

I. General

This final report covers work carried out by the principal investigator and a graduate research assistant at the Department of Mathematics during the 13 months period from July 1, 1979 to July 31, 1980 under Contract F49620-79-C-0186.

The progress has resulted in three technical papers listed in Section II, which will be submitted for publication as soon as completed.

During the report period, the following people contributed to the project: Professor James T. Lo and Mr. Sze-Kui Ng.

II. Publications

- (1) Transition Probabilities of Homogeneous Markov Processes on Compact Lie Groups, to appear.
- (2) Projecting Homogeneous Markov Processes onto Compact Homogeneous Spaces, to appear.
- (3) Approximation of Brownian Motion Densities on the Three Dimensional Rotation Group, to appear.

III. Summary of Progress

An optimal estimation scheme for continuous-time rotational processes with one degree of freedom was obtained in [1]. The estimation problem for continuous-time rotational processes with three degrees of freedom turns out to be rather difficult. Many attempts were made by the P.I. in the past few years on the problem. While these attempts produced minimal results, many crucial issues have been placed in

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

perspective. Because of the theoretical and practical importance of the problem, a concentrated effort was finally made during the report period to study these issues.

(1) Transition Probabilities of Homogeneous Markov Processes on Compact Lie Groups:

A signal process on $SO(3)$ (the 3-dimensional rotation group) can be constructed by injecting a 3-dimensional Brownian motion into $SO(3)$ [2]. It is known that such a signal process is governed by a bilinear stochastic differential equation. But how do we compute the transition density for this process? In the $SO(2)$ case, such a transition density has an explicit expression called the folded normal density. It is also called a theta function. Then what is a general folded normal density or theta function? This question is answered in a much more general context as follows:

Let $\{T_g^{(n)}\}$ be a sequence of finite dimensional irreducible unitary representations of a k -dimensional compact Lie group G where $T_g^{(1)} = 1$. The infinitesimal operators $A_i^{(n)}$ of $T_g^{(n)}$ is defined as

$$A_i^{(n)} = \lim_{t \rightarrow 0} \frac{1}{t} \left(T_{g_i(t)}^{(n)} - I_n \right), \quad i = 1, \dots, k$$

where $\{g_i\}$ are k one-parameter subgroups of which the tangent vectors at the group identity e are orthogonal. There is a neighborhood of e in which a parameterization $x_1(g), \dots, x_k(g)$ satisfies

$$T_g^{(n)} = \exp \left\{ \sum_{i=1}^k x_i(g) A_i^{(n)} \right\}, \quad \forall g \in G.$$

These local coordinates $\{x_i(g)\}$ can be extended continuously to the

Accession No.	
NTIS GRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability	
Dist	Avail and Special
A	

entire G such that they do not vanish simultaneously except at e .

Let X_i denote the left invariant vector fields corresponding to g_i .

Hung [3] proved that for a smooth function f and the transition

probability $P(t, g, E)$ of a homogeneous Markov process, the function

$u(t, g) = \int_G f(\sigma) P(t, g, d\sigma)$ satisfies the Kolmogorov backward equation, with $P(0, e, E)$ concentrates at e .

$$\frac{\partial u(t, g)}{\partial t} = \sum_{i,j=1}^k a_{ij} X_i X_j u(t, g) + \sum_{i=1}^k a_i X_i u(t, g) \\ + \int_{G \setminus \{e\}} \left[u(t, gh) - u(t, g) - \sum_{i=1}^k x_i(h) X_i u(t, g) \right] dF(h),$$

for a real vector $[a_1, \dots, a_k]$, a non-negative definite matrix

$[a_{ij}]_{i,j=1}^k$, and a positive measure F on $G \setminus \{e\}$ such that

$$\int_{G \setminus \{e\}} \sum_{i=1}^k x_i^2(g) dF(g) < \infty.$$

Our result is that the transition probability $P(t, g, E)$ can be calculated by using the following fact:

$$\text{Let } C_n(t) = \exp t \left\{ \sum_{i,j=1}^k a_{ij} A_i^{(n)} A_j^{(n)} + \sum_{i=1}^k a_i A_i^{(n)} \right. \\ \left. + \int_{G \setminus \{e\}} \left[T_g^{(n)} - I_n - \sum_{i=1}^k x_i(g) A_i^{(n)} \right] dF(g) \right\}$$

and $\rho_n = (\dim. \text{ of } T_g^{(n)}) / (\text{the volume of } G)$.

Under certain conditions, there exists a sequence of functions

$$f_r(t, g, \sigma) = \sum_{n=0}^{N(r)} \rho_n \lambda_n^r \operatorname{tr} \left(T_g^{(n)*} T_\sigma^{(n)} C_n^*(t) \right)$$

such that for $h \in C(G)$,

$$\lim_{r \rightarrow \infty} \int h(\sigma) f_r(t, g, \sigma) d\sigma = \int h(\sigma) P(t, g, d\sigma)$$

where $|\lambda_n^r| \leq 1$, $\lambda_0^r = 1$, $\lim_{r \rightarrow \infty} \lambda_n^r = 1$, $n = 0, 1, \dots$ and $\lim_{r \rightarrow \infty} N(r) = \infty$.

For example, let us consider $G = SO(3)$ and the Brownian motion governed by the bilinear Ito equation,

$$dX(t) = \begin{bmatrix} -dt & -dw_3 & dw_2 \\ dw_3 & -dt & -dw_1 \\ -dw_2 & dw_1 & -dt \end{bmatrix} X(t), X(0) = I$$

where $[w_1, w_2, w_3]$ is the standard Brownian motion in R^3 . The transition density $p(t, g, \sigma)$ of the process satisfies

$$\begin{aligned} \frac{\partial p}{\partial t}(t, g, \sigma) &= \frac{1}{\sin \beta} \frac{\partial}{\partial \beta} \left\{ \sin \beta \frac{\partial p}{\partial \beta}(t, g, \sigma) \right\} + \frac{1}{\sin^2 \beta} \left\{ \frac{\partial^2 p}{\partial \lambda^2}(t, g, \sigma) \right. \\ &\quad \left. - 2 \cos \beta \frac{\partial^2 p}{\partial \alpha \partial \gamma}(t, g, \sigma) + \frac{\partial^2 p}{\partial \alpha^2}(t, g, \sigma) \right\} \end{aligned}$$

$$p(0, e, \sigma) = \delta(\sigma)$$

where $g = (\alpha', \beta', \gamma')$ and $\sigma = (\alpha, \beta, \gamma)$. It can be explicitly written as

$$\begin{aligned} p(t, g, \sigma) &= \sum_{j=0}^{\infty} \frac{2j+1}{8\pi^2} e^{-j(j-1)t} \sum_{m, n=-j}^j e^{im\alpha' - j} p_{mn}^j(\cos \beta') e^{in\gamma'} e^{-im\alpha} \\ &\quad \cdot p_{mn}^j(\cos \beta) e^{-in\gamma} \end{aligned}$$

where $e^{-im\alpha'} p_{mn}^j(\cos \beta') e^{-in\gamma}$ is the generalized spherical functions of order j .

(2) Projecting Homogeneous Markov Processes onto Compact Homogeneous Spaces:

The 2-sphere S^2 , the surface of a ball, is a natural state space for the directional processes. It is isomorphic to the quotient space $SO(3)/SO(2)$. A signal process on S^2 can be induced by projecting a signal process on $SO(3)$ onto S^2 . If the signal process on $SO(3)$ is Brownian, then what is the transition density of the induced Brownian motion on S^2 ? This question is also answered in a general context as follows:

Let H be a closed subgroup of the Lie group G . Corresponding to a projection $\Pi: g \rightarrow Hg$ from G onto the homogeneous space $H \backslash G$, there exists a C^∞ mapping s such that $\Pi \circ s = 1$. For each s , an element $g \in G$ is expressed uniquely as $s_1(x)h$ for some $x \in H \backslash G$ and $h \in H$. A homogeneous process on $H \backslash G$ can be induced by projecting through Π a homogeneous process on G .

The transition probability $P(t, x, B)$ of the resulting process on $H \backslash G$ can be calculated by using the following fact:

Under certain conditions, there exists a sequence of functions

$$f_r(t, x, y) = \sum_{n=0}^{N(r)} \rho_n \lambda_n^r \operatorname{tr} \left[T_{s(x)}^{(n)*} \int_H T_g^{(n)} dg T_{s(y)}^{(n)} C_n^*(t) \right]$$

such that for $h \in C(H \backslash G)$,

$$\lim_{r \rightarrow \infty} \int_{H \backslash G} h(y) f_r(t, x, y) d\mu(y) = \int_{H \backslash G} h(y) P(t, x, dy)$$

where μ is an invariant measure on $H \backslash G$ and the other symbols are defined as in (1).

For example, the transition density of the homogeneous process that is the projection of the example in (1) satisfies

$$\begin{aligned} \frac{\partial p}{\partial t}(t, x, y) = & \frac{1}{\sin \beta} \frac{\partial}{\partial \beta} \left[\sin \beta \frac{\partial p}{\partial \beta}(t, x, y) \right] \\ & + \frac{1}{\sin^2 \beta} \frac{\partial^2 p}{\partial \gamma^2}(t, x, y) \end{aligned}$$

$$p(0, 0, y) = \delta(y)$$

where $x = (\alpha', \beta')$ and $y = (\alpha, \beta)$, the spherical coordinates. It can be explicitly written as

$$p(t, x, y) = \sum_{k=1}^{\infty} \frac{2k+1}{4\pi} e^{-k(k+1)t} \sum_{n=-k}^k \bar{Y}_k^n\left(\frac{\pi}{2} - \gamma', \beta'\right) Y_k^n\left(\frac{\pi}{2} - \gamma, \beta\right)$$

where $Y_k^n\left(\frac{\pi}{2} - \gamma, \beta\right)$, $-k \leq n \leq k$ are the unnormalized spherical functions of the k -th order.

(3) Approximation of Brownian Motion Densities on the Three Dimensional Rotation Group:

The transition densities obtained above are very complicated functions, as illustrated by the explicit expressions for the transition densities of the simplest possible Brownian motions on $SO(3)$ and S^2 . In order to use them for estimation, a natural question is whether they can be approximated closely by simpler closed-form functions.

Let us restrict our attention to the density obtained in the example in (1):

$$\begin{aligned} p_0(\alpha, \beta, \gamma) = & p(t, e, (\alpha, \beta, \gamma)) \\ = & \sum_{l=0}^{\infty} \frac{2l+1}{8\pi} e^{-\sigma l(l+1)t} \sum_{m=-l}^l e^{-im(\alpha + \gamma)} P_{lm}^l(\cos \beta). \end{aligned}$$

It can be approximated closely by

$$p_k(\alpha, \beta, \gamma) = \frac{1}{2\pi C_0(k)C_1(k)} e^{k \cos \beta + k \cos (\alpha + \gamma)}$$

$$C_0(k) = \int_0^{2\pi} e^{k \cos \alpha} d\alpha$$

$$C_1(k) = \int_0^\pi e^{k \cos \beta} d\beta$$

where k is determined by $\coth k - \frac{1}{k} = e^{-2\sigma}$. Let the Fourier series expansion of p_k be

$$p_k(\alpha, \beta, \gamma) = \sum_{\ell=0}^{\infty} \frac{2\ell+1}{8\pi^2} \operatorname{tr} \left(T_g^{(\ell)} C_\ell^*(k) \right)$$

where $C_\ell(k) = \left(C_{nm}^\ell(k) \right)$ is a $2\ell+1$ dimensional coefficient matrix.

It is proven that

(a) For each m, ℓ such that $-\ell \leq m \leq \ell$,

$$C_{mm}^\ell(k) = e^{-\sigma\ell(\ell+1)} + O\left(\frac{1}{k^2}\right)$$

$$C_{nm}^\ell(k) = 0, \quad \text{for } n \neq m$$

(b) Given any $\epsilon > 0$ and positive integer N , there exists a number $K(\epsilon, N)$ such that, for

$$k \geq K(\epsilon, N), \quad \ell^2 \leq Nk, \quad \text{and } -\ell \leq m \leq \ell,$$

$$\left| C_{mm}^\ell(k) - e^{-\sigma\ell(\ell+1)} \right| \leq \epsilon.$$

It is noted that (a) is analogous to those for spheres and circles obtained in [4] and [5]. (b) is new and stronger than (a). (b) for spheres and circles is also obtained in our work.

References

- [1] J. T. Lo and A. S. Willsky, "Estimation for Rotational Processes with One Degree of Freedom I: Introduction and Continuous-Time Processes," IEEE Transactions on Automatic Control, AC-20-1, February 1975.
- [2] J. T. Lo, "Representation of Continuous Curves on the 3-dimensional Rotation Group," Mathematical System Theory, 8-4, pp. 368-375, 1975.
- [3] G. A. Hunt, "Semi-groups of Measures on Lie Groups," Trans. Amer. Math. Soc., 81, pp. 264-293, 1956.
- [4] P. H. Roberts and H. D. Ursell, "Random Walk on a Sphere and on a Riemannian Manifold," Philos. Trans. A, 252, pp. 317-56, 1960.
- [5] M. A. Stevens, "Random Walk on a Circle," Biometrika 50, 3 and 4, pp. 385-390, 1963.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 80 - 1055	2. GOVT ACCESSION NO. AD-A092 386	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DETECTION, ESTIMATION, AND CONTROL ON GROUP MANIFOLDS	5. TYPE OF REPORT & PERIOD COVERED Final	
7. AUTHOR(s) J. T. Lo	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Maryland Baltimore County Department of Mathematics Catonsville, Maryland 21228	8. CONTRACT OR GRANT NUMBER(s) F49620-79-C-0186	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, D. C. 20332	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/A1	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE July 1980	
	13. NUMBER OF PAGES 9	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An optimal estimation scheme for continuous-time rotational processes with one degree of freedom was obtained, in (1). The estimation problem for continuous-time rotational processes with three degrees of freedom turns out to be rather difficult. Many attempts were made by the P.I. in the past few years on the problem. While these attempts produced minimal results, many crucial issues have been placed in perspective. Because of the theoretical and practical importance of the problem, a concentrated effort was finally made during the report period to study these issues.		

IED
8