

AD-A094 120

AIR FORCE GEOPHYSICS LAB HANSCOM AFB MA
BEHAVIOR OF HIGH-LATITUDE IRREGULARITIES DURING GEOMAGNETIC DIS--ETC(U)
JUN 80 Z HOUMINER, J AARONS

F/G 4/1

UNCLASSIFIED

AFGL-TR-80-0194

NL

1-1

20

6-10-77

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1

1


END

DATE

FORMED

2 81

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A094120

Behavior of High-Latitude Magnetosphere
During Geomagnetic Disturbances

ZWISCHNER
JULIA ARONS

24 June 1964

12 21

Approved for public release; distribution unlimited

SPACE PHYSICS DIVISION PROJECT 64-23
AIR FORCE GEOPHYSICS LABORATORY
HANCOM AFB, MASSACHUSETTS 01931


AIR FORCE SYSTEMS COMMAND, USAF

FILE COPY

This report has been reviewed by the DOD and is
releasable to the National Technical Information Service.

This technical report has been reviewed and
is approved for publication.

FOR THE COMMANDER


Chief Scientist

Qualified requestors may obtain additional copies from the
Defense Technical Information Center. All others should contact the
National Technical Information Service.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFGL-TR-80-0194	2. GOVT ACCESSION NO. AD-A094120	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) BEHAVIOR OF HIGH-LATITUDE IRREGULARITIES DURING GEOMAGNETIC DISTURBANCES		5. TYPE OF REPORT & PERIOD COVERED Scientific. Interim.
7. AUTHOR(s) Zwi Houminer* Jules Aarons		6. PERFORMING ORG. REPORT NUMBER ERP No. 709 ✓
9. PERFORMING ORGANIZATION NAME AND ADDRESS Air Force Geophysics Laboratory (PHP) ✓ Hanscom AFB Massachusetts 01731		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory (PHP) Hanscom AFB Massachusetts 01731		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62101F 46430505
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 24 June 1980
		13. NUMBER OF PAGES 19
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES *NRC Resident Research Associate		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ionospheric scintillation Mean scintillation pattern Auroral irregularity Plasmapause irregularity Magnetic storms		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) ✓ Scintillation observations of VHF and UHF transmissions from geostationary satellites at Goose Bay have been used to study the average characteristics of the high latitude irregularity region. The paper describes the average time development and mean diurnal pattern of irregularities during 58 magnetic storms in 1971-1976. The diurnal variation at Goose Bay shows two peaks of scintillation activity. One peak occurs during the afternoon hours, while the second occurs during the night. The average diurnal pattern is independent		

DD FORM 1 JAN 79 1473 EDITION OF 1 NOV 68 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. Abstract (Continued)

of type of storm. Diurnal and seasonal effects appear only in the first day of storm commencement and not in the following days which show a very similar diurnal picture.

X

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Accession For	
DTIC	☒
DTIC TAB	☐
Unannounced	☐
Justification	
Distribution/	
Availability Codes	
Data Source or	
Special	
A	

Contents

1. INTRODUCTION	5
2. DATA ANALYSIS	5
3. SCINTILLATION STORM PATTERNS	8
4. CONCLUSION	17
REFERENCES	19

Illustrations

1. 15-Minute Scintillation Indices for ATS-3 Signal Recorded at Goose Bay During October 30–November 3, 1972	7
2. 15-Minute Scintillation Indices for ATS-3 Signal Recorded at Goose Bay During August 3–7, 1972	8
3. Average Diurnal Variation Pattern SD(t) for 58 Magnetic Storms During 1971–1976	9
4. Average Diurnal Variation Pattern SD(t) for 26 Sudden Commencement Storms During 1971–1976	10
5. Average Diurnal Variation Pattern SD(t) for 32 Gradual Commencement Storms During 1971–1976	10
6. Average Diurnal Variation Pattern SD(t) for 35 Storms Which Started During Daytime	11

Illustrations

7. Average Diurnal Variation Pattern SD(t) for 23 Storms Which Started During Nighttime	11
8. Average Diurnal Variation Pattern SD(t) for 15 Storms Occurring in Spring	12
9. Average Diurnal Variation Pattern SD(t) for 17 Storms Occurring in Summer	12
10. Average Diurnal Variation Pattern SD(t) for 18 Storms Occurring in Autumn	13
11. Average Diurnal Variation Pattern SD(t) for Eight Storms Occurring in Winter	13
12. Average Storm Commencement Time Pattern Dst(t) for 58 Magnetic Storms in 1971-1976	14
13. Average Storm Commencement Time Pattern Dst(t) for 26 SSC Storms in 1971-1976	15
14. Average Storm Commencement Time Pattern Dst(t) for 32 GSC Storms in 1971-1976	15
15. Average Storm Commencement Time Pattern Dst(t) for 35 Daytime Storms	16
16. Average Storm Commencement Time Pattern Dst(t) for 23 Nighttime Storms	16
17. The Average Diurnal Pattern of the H Component of the Magnetic Field, Recorded at St. John's, for Eight Magnetic Storms in 1972	17

Table

1. Storm Periods	6
------------------	---

Behavior of High - Latitude Irregularities During Geomagnetic Disturbances

1. INTRODUCTION

Scintillation observations of VHF and UHF transmissions from geostationary satellites at high-latitude stations can be used to study the average characteristics of the irregularity region during magnetic storms. Houminer et al¹ described the average time development and mean diurnal pattern of irregularities during 12 magnetic storms in 1971-1972, based on observations taken at Goose Bay and Sagamore Hill. This paper will present results of the average behavior of irregularities in the auroral region, using scintillation observations of geostationary satellites at Goose Bay during 58 magnetic storms in 1971-1976.

2. DATA ANALYSIS

The data base used in the present analysis is a continuous series of scintillation measurements taken at Goose Bay (GB) during periods of geomagnetic activity. The GB invariant latitude of the 350 km ionospheric intersection to the satellite is 60° . Table 1 lists the storm commencement periods. The storms were of the

(Received for publication 20 June 1980)

1. Houminer, Z., Aarons, J., and MacKenzie, E. (1980) Plasmopause and auroral oval irregularities during magnetic storms, URSI-IEEE Symposium, June 2-6, 1980, Quebec.

Table 1. Storm Periods

Number	Date	S. C. Time (U. T.)	Station Name	Max. Value of K_p	Max. Value of A_p
1	17 December 1971	SSC = 14:18	FRED	6	67
2	21 January 1972	SSC = 11:51	FRED	5	32
3	24 February 1972	SSC = 06:42	SJUA	5	33
4	6 March 1972	SSC = 21:08	SJUA	7	45
5	29 April 1972	GSC = 03:--	FRED	6	42
6	15 May 1972	SSC = 18:49	FRED	6	38
7	17 June 1972	SSC = 13:12	FRED	8	126
8	24 July 1972	GSC = 19:--	FRED	6	33
9	4 August 1972	SSC = 01:19	FRED	9	182
10	8 August 1972	SSC = 23:54	FRED	6	74
11	13 September 1972	SSC = 12:40	FRED	8	54
12	18 October 1972	SSC = 17:46	FRED	5	34
13	31 October 1972	SSC = 16:58	FRED	7	98
14	15 November 1972	GSC = 09:--	NEWP	5	31
15	12 December 1972	SSC = 21:45	BOUL	5	30
16	27 January 1973	GSC = 04:--	FRED	5	33
17	21 February 1973	GSC = 16:--	BOUL	6	54
18	1 March 1973	GSC = 15:--	TUCS	5	40
19	18 March 1973	GSC = 12:--	FRED	6	82
20	31 March 1973	GSC = 13:--	BOUL	6	91
21	13 April 1973	SSC = 04:34	FRED	6	57
22	26 April 1973	GSC = 12:--	FRED	5	59
23	13 May 1973	GSC = 17:--	FRED	7	80
24	21 May 1973	SSC = 02:54	FRED	6	46
25	10 June 1973	GSC = 15:--	FRED	5	34
26	17 June 1973	GSC = 19:--	FRED	5	38
27	28 June 1973	GSC = 21:--	FRED	5	41
28	14 July 1973	GSC = 15:--	FRED	5	32
29	26 July 1973	GSC = 03:--	FRED	5	36
30	23 August 1973	GSC = 12:--	FRED	6	46
31	9 September 1973	GSC = 08:--	TUCS	5	44
32	22 September 1973	GSC = 21:--	FRED	6	63
33	2 October 1973	GSC = 05:--	FRED	6	48
34	16 October 1973	SSC = 05:20	TUCS	5	30
35	21 October 1973	GSC = 01:--	FRED	5	37
36	28 October 1973	GSC = 07:--	FRED	6	86
37	24 November 1973	GSC = 13:--	FRED	6	42

SSC (Sudden Storm Commencements) and GSC (Gradual Storm Commencements) types, and were chosen according to the criteria of Mendillo.²

For each storm, five days of scintillation data (a day before storm commencement, the day of the storm, and three days after) were used to determine the average storm pattern. The data consisted of 15-minute scintillation indices reduced by the method of Whitney et al.³ Examples of the five-day scintillation data are shown in Figures 1 and 2.

For each storm, the Storm Enhancement Factor (S) was calculated to eliminate the quiet day diurnal pattern. This factor S:

$$S = F(t) - \bar{F}$$

where $F(t)$ is the observed scintillation index (SI) in dB for the 5-day period, and $\bar{F}$ is the average scintillation index in dB determined for magnetically quiet days. $\bar{F}$ is determined as shown in Aarons et al.⁴ and varies with local time, season and solar flux for each station.

This average storm enhancement factor was then analyzed using two different time approaches. The first involves construction of average storm time (Dst). This represents the storm induced variations calculated at hourly intervals from the time of commencement of each storm. The second method is the average local time behavior SD(t). In this method, the first day of a storm is defined to be the remaining portion of the day after the Sudden Commencement. Following this definition of the first day, each hourly value is set into its correct local time for the averaging process. (Mendillo and Klobuchar,⁵ Mendillo²)

It should be noted that scintillations are rather low for Goose Bay during quiet magnetic conditions. The mean scintillation index ($\bar{F}$) is zero for 75 percent of the time, and is greater than 4 dB for 3 percent of the time. It does not exceed 5 dB.

3. SCINTILLATION STORM PATTERNS

The average local time storm pattern SD(t) for 58 storms during 1971-1976 is shown in Figure 3. It can be seen that the diurnal pattern of scintillation at Goose Bay shows two peaks of scintillation activity. One peak occurs during the afternoon at about 17-18 hours local time, especially on the days following the storm

2. Mendillo, M. (1978) Behavior of the Ionospheric F-Region During Geomagnetic Storms, AFGL-TR-78-0092 (II), AD A056 978.
3. Whitney, H. E., Aarons, J., and Malik, C. (1969) A proposed index for measuring ionospheric scintillations, Planet. Space Sci. 17:1069.
4. Aarons, J., MacKenzie, E., and Bhavnani, K. (1980) High-latitude analytical formulas for scintillation levels, Radio Sci. 15:115.
5. Mendillo, M. and Klobuchar, J. A. (1974) An Atlas of the Mid-Latitude F-Region Response to Geomagnetic Storms, AFCRL-TR-74-0065, AD 778 069.




Figure 1. 15-Minute Scintillation Indices for ATS-3 Signal Recorded at Goose Bay During October 30–November 3, 1972. The geomagnetic storm commencement is indicated with an arrow.




Figure 2. 15-Minute Scintillation Indices for ATS-3 Signal Recorded at Goose Bay During August 3–7, 1972. The storm commencement is indicated with an arrow.




Figure 3. Average Diurnal Variation Pattern SD(t) for 58 Magnetic Storms During 1971-1976

commencement. The second peak occurs during the night and is centered at about local midnight.

Figure 4 shows SD(t) for 26 sudden commencement storms (type 1) while Figure 5 gives SD(t) for 32 gradual commencement storms (type 2). Figures 4 and 5 show large similarity in the diurnal pattern with two peaks in scintillation activity. Thus, the diurnal pattern does not depend on the type of storm.

Next, the storms were grouped according to the time of storm commencement. Figure 6 gives SD(t) for storms which commenced during the day, while Figure 7 gives SD(t) for nighttime storm commencement. Comparison of Figure 6 to Figure 7 shows a large difference in the diurnal pattern at the day of the storm commencement. However, the diurnal pattern during the following days is very similar for day or nighttime storms.

The seasonal effect on SD(t) is shown in Figures 8 to 11, where storms were grouped according to season. Again, there are some differences in the diurnal pattern at the day of storm commencement. However, for the following days, the diurnal pattern is very similar, showing the double hump characteristics.

The average storm commencement time pattern Dst(t) for 58 storms in 1971-1976 is shown in Figure 12. The Dst(t) pattern shows an immediate response to the storm commencement. However, this increase is small in comparison to the scintillation increase during the main phase of the storm. The Dst(t) patterns for




Figure 4. Average Diurnal Variation Pattern $SD(t)$ for 26 Sudden Commencement Storms During 1971-1976




Figure 5. Average Diurnal Variation Pattern $SD(t)$ for 32 Gradual Commencement Storms During 1971-1976




Figure 6. Average Diurnal Variation Pattern SD(t) for 35 Storms Which Started During Daytime




Figure 7. Average Diurnal Variation Pattern SD(t) for 23 Storms Which Started During Nighttime




Figure 8. Average Diurnal Variation Pattern SD(t) for 15 Storms Occurring in Spring




Figure 9. Average Diurnal Variation Pattern SD(t) for 17 Storms Occurring in Summer




Figure 10. Average Diurnal Variation Pattern SD(t) for 18 Storms Occurring in Autumn




Figure 11. Average Diurnal Variation Pattern SD(t) for Eight Storms Occurring in Winter




Figure 12. Average Storm Commencement Time Pattern $Dst(t)$ for 58 Magnetic Storms in 1971-1976

types 1 and 2 storms are shown in Figures 13 and 14. The patterns indicate that the immediate increase in scintillation occurs only for type 1 storms, that is, sudden storm commencement. Figures 15 and 16, which give $Dst(t)$ for day and nighttime storms, indicate that the sudden scintillation response is a nighttime phenomena.




Figure 13. Average Storm Commencement Time Pattern $Dst(t)$ for 26 SSC storms in 1971-1976




Figure 14. Average Storm Commencement Time Pattern $Dst(t)$ for 32 GSC Storms in 1971-1976




Figure 15. Average Storm Commencement Time Pattern $Dst(t)$ for 35 Daytime Storms




Figure 16. Average Storm Commencement Time Pattern $Dst(t)$ for 23 Nighttime Storms

4. CONCLUSION

Scintillation observations of VHF and UHF transmissions from geostationary satellites at Goose Bay show a two-peak diurnal behavior of scintillation during magnetic storms. One peak occurs during the afternoon hours, while the second occurs during the night. Houminer et al¹ have shown that the afternoon scintillations at Goose Bay occur when the southern wall of the electron density trough, or the plasmopause, passes through the line of sight to the satellite. The nighttime scintillation is associated with auroral oval electron density irregularities.

Figure 17 shows the average behavior of the H component of the magnetic field during eight magnetic storms in 1972. The magnetic data were recorded in St. John's, which is roughly at the same longitude and latitude as the subionospheric point to the satellite from Goose Bay.




Figure 17. The Average Diurnal Pattern of the H Component of the Magnetic Field, Recorded at St. John's, for Eight Magnetic Storms in 1972

The average H behavior shows a positive increase in H during the afternoon hours, and a negative excursion at local midnight. The similarity in time indicates that the afternoon increase is correlated with the plasmopause scintillation, while the midnight peak is correlated with the auroral irregularities.

The results presented here show that the average diurnal pattern of scintillation is not dependent on type of storm. The commencement time of the storm affects the diurnal behavior during the first day; however, the following days show a very similar diurnal picture. Seasonal effects also appear in the first day of storm commencement and not in the following days.

The immediate response of scintillation enhancement to storm commencement seems to occur only for sudden commencement storms, and only for storms with SSC at night.

References

1. Houminer, Z., Aarons, J., and MacKenzie, E. (1980) Plasmopause and auroral oval irregularities during magnetic storms, URSI-IEEE Symposium, June 2-6, 1980, Quebec.
2. Mendillo, M. (1978) Behavior of the Ionospheric F-Region During Geomagnetic Storms, AFGL-TR-78-0092 (II), AD A056 978.
3. Whitney, H.E., Aarons, J., and Malik, C. (1969) A proposed index for measuring ionospheric scintillations, Planet. Space Sci. 17:1069.
4. Aarons, J., MacKenzie, E., and Bhavnani, K. (1980) High-latitude analytical formulas for scintillation levels, Radio Sci. 15:115.
5. Mendillo, M. and Klobuchar, J.A. (1974) An Atlas of the Mid-Latitude F-Region Response to Geomagnetic Storms, AFCRL-TR-74-0065, AD 778 069.

DATE
FILMED
8