

AD-A095 248

SCRIPPS INSTITUTION OF OCEANOGRAPHY LA JOLLA CA

F/G 8/1

PHYSICS

I EXPEDITION, 19--ETC

UNCLASSIFIED

SIO-REF-74-20

ML

1 of 1
ADA
C95248

AD A
C95.2.8

END
DATE
FILMED
3-81
DTIC

AD A095248

DEC FILE COPY

Ordered
4/7/80

LEVEL

Br⁰2
=

UNIVERSITY OF CALIFORNIA SCRIPPS INSTITUTION OF OCEANOGRAPHY

DTIC
ELECTE
FEB 18 1981
D
C

data report

PHYSICAL, CHEMICAL AND BIOLOGICAL DATA

CLIMAX I EXPEDITION

19 September - 28 September 1968

N00014-69-A-0200-6049

SIO Reference 74-20
1 September 1974

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

80 9 3 052

7 October 1974

ERRATA

The ship's name was inadvertantly
entered RV ARGO. It should be
replaced with RV HORIZON.

(1)

UNIVERSITY OF CALIFORNIA
SCRIPPS INSTITUTION OF OCEANOGRAPHY

PHYSICAL, CHEMICAL AND
BIOLOGICAL DATA,

CLIMAX I EXPEDITION

19 September - 28 September 1968

Cruise Sponsored by:

National Science Foundation
Marine Research Committee

Data Processing and Analysis Sponsored by:

Office of Naval Research
Sandia Corporation
Marine Research Committee

SIO-Reference 74-20

Approved for distribution:

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

W. A. Nierenberg
W. A. Nierenberg, Director

CONTENTS

Introduction	3
Literature Cited	11
List of Figures and Tables	12
Personnel	14
Tabulated Data	
Hydrographic	17
Biological	29
Distribution List	

INTRODUCTION

Many of the earlier biological studies of the Central North Pacific Water mass have been of an exploratory nature. Expeditions NORTHERN HOLIDAY (1951), TRANS-PAC (1954), NORPAC-POFI (1955), URSA MAJOR (1964), and ZETES (1966) have all surveyed this area. One of the results of these surveys was the discovery of very abrupt latitudinal faunal discontinuities in mid-ocean. The fauna and flora of the North Pacific Water mass was found to differ strongly from that of the Subarctic Water mass and from the transition domain. There are also important differences from the Equatorial Water mass. Further, in samples taken within the gyre-like circulation of the Central Water mass the homogeneity of species content of macrozooplankton, phytoplankton and neuston is great relative to other planktonic environments (McGowan 1971). The standing crops of organisms here are always low and do not appear to vary greatly, either spatially or temporally (McGowan and Williams, 1973). The dimensions of this area are approximately 1200 by 4500 nautical miles, thus it represents a significant fraction of the Pacific Ocean.

The temperature and salinity structure of this area, as determined from the early cruises listed above, also indicates a certain monotony of hydrographic conditions as do measurements of phosphate, silicate and oxygen. Although there are seasonal variations in these properties in the upper layer, the range is comparatively small and there is little spatial heterogeneity for large parts of the year.

Because of the regularity of the distributions of both biological and physical variables and the lack of evidence of any large-scale horizontal advection of either biological or physical-chemical properties, the central gyre of the North Pacific was selected for a long-term study of community structure and function. On the basis of biogeographic studies the central gyre has been determined to be an independent faunal province. An area in the vicinity of 28°N 155°W was selected as the "center" of the eastern half of the gyre. At this locale we planned an intensive series of biological and physical measurements designed to reveal aspects of the structure and function of the community. Our assumption was that samples taken here would be representative of a much larger area within the gyre itself.

→ The first cruise of this series was Climax I and its specific purpose was to investigate the vertical distribution of phyto-

Handwritten notes and a stamp on the bottom right of the page:

✓
41-182-101 or 102
389-81 or 101
DIST. 101
EXPERIMENT 101
DATE 101
A

^{cat} plankton and macrozooplankton species, microzooplankton biomass, temperature, salinity, oxygen, nutrients, chlorophyll-a, primary productivity and light. The tactics used were to follow a pair of parachute drogues for a period of about eleven days while taking replicate measurements of these properties. Two drogues were used in order to ensure that complicated circulation patterns were not present and that small scale eddies would not disperse the properties being measured. If the drogues remained together and followed the "same" track as they drifted, it was taken as evidence that the sampling, done between the drogues, was in a reasonably coherent body of water. ↗

DAILY ROUTINE

The daily routine of sampling while following the drogues consisted of Bongo net tows taken at 6 or 8 depth ranges throughout the (24 hour) day. The maximum depth reached was 600m. Microzooplankton pumping from 5 depth ranges was done mainly in the early morning, late morning and early evening hours. The maximum depth reached was 200m. Samples of water from the pumping device were taken for nutrient, chlorophyll-a and phaeo-pigment analyses. Continuous salinity, temperature, depth profiles (S/T/D) were taken in the late morning, near noon and early evening hours. Bottle casts for nutrients, chlorophyll-a and phaeo-pigment analyses were made in the late morning and early evening. Submarine photometer and secchi disc lowerings were made near local apparent noon. Simulated in situ productivity measurements were done daily from local apparent noon until sunset. A few continuous traces of phosphate, using a flow of water from the microzooplankton pump, were made with an autoanalyzer. Continuous tracings of in situ "chlorophyll" fluorescence were also made. Echo sounding was done during the early morning and evening hours for records of the deep scattering layers. Visual sightings of birds, mammals and fish schools were made. Weather and sea state were recorded.

THE DROGUES

The original plan was to place three parachute drogues in a triangular configuration, 5 miles on each side. The parachutes were set at 10m depth. These three drogues were set in place on the afternoon of 18 September, 1968. At 0540 on 19 September the position of the drogues was determined by celestial navigation for #1 drogue (27°00'N, 155°18'W) and by radar bearings from #1 to #2 and #3 (Fig. 2, inset).

These three drogues moved together in a westerly direction at

speeds up to 1 knot. However, on the morning of 21 September they slowed down, and changed direction to the southwest. During the time of the direction change (between 0040 and 0550) #3 drogue sank. Shortly thereafter #2 drogue sank so that by 1300 on 21 September we were following only one drogue. This drogue (#1) changed direction again and began to move slowly to the northwest during the afternoon of that day. It continued to move slowly that night. During the morning of 22 September we were able to install a fourth drogue about 4 nautical miles to the east of #1. These two drogues (#1 and #4) moved as a pair in a northwesterly direction until 28 September when we terminated our study and purposely sank the drogues (Fig. 2).

Thus the sampling, which began on 19 September, was done at first in the center of the triangle, then for a short distance near one drogue only, and finally, between a pair of drogues. Because of the very short distance traveled between the time #2 drogue sank and the time of insertion of #4 drogue, this may be considered a continuous series of samples, with little or no dispersal of the water the drogues were tracking. The distance traveled by #1 drogue was 186.6 nautical miles. Table 1 shows wind directions and speeds during the entire eleven day period. Table 2 shows distance traveled and speed of the drogues during the same period.

METHODS

Macrozooplankton

The macrozooplankton was sampled with Bongo nets (McGowan and Brown, 1966) of 505 μ mesh. This is a paired net with each net of the pair having an area of 0.396m². The depth range of the samples was from the surface to 600m. The nets were deployed in sets of four frames which were intended to sample depths of 0-25m, 25-50m, 50-75m and 75-100m simultaneously. This was followed immediately by a set of four frames which sampled depths of 100-225m, 225-350m, 350-475m and 475-600m. Thus, eight depths were sampled by these paired nets. One complete series yielded 16 macrozooplankton samples. One such series was done near noon and the other near midnight on each day. We followed this routine from 19 September to 1630 on 24 September. However, it became apparent by then that some of the Bongo frame mechanisms were frequently malfunctioning and several of our nets were torn beyond repair. We ceased sampling for about 36 hours and began a new series using only two frames per cast beginning at 1337 on 26 September.

This new series sampled depths of 0-25m, 25-50m, 50-75m, 75-100m,

100-350m and 350-600m. Thus, sampling at only six depths per series was done for the remainder of the expedition. A further change in the routine was that these new series were done near noon, sunset, midnight and sunrise. This routine was followed until the afternoon of 27 September.

The maximum depth of tow on many early tows of these series was determined by a Benthos depth-telemetering pinger. Since the seas were calm, a relationship between wire angle and tow depth was soon established and on the remaining series, depths were determined from the wire angle.

The flow "control" meters of the 4 Bongo frames (#2, #3, #4, and #6) were calibrated against T.S.K. flow meters in the mouths of the net. Five replicate calibration runs were done on each frame. On the basis of these data the volumes of water filtered by the nets was estimated. Not all of these nets "worked" properly. Table 3 shows the dates, times and depths of sampling as well as the displacement volumes of the catches and comments on the performance of each net and on the reliability of the samples. Figure 3 shows the distribution of macrozooplankton sampling effort with time and depth. All times shown in the Tables are local.

Microzooplankton

The biomass of microzooplankton was estimated from samples taken by means of a pumping system described by Beers, et al (1967). In this system, water is pumped from depths to a series of nested "deck" filters aboard ship. These filters were of 363 μ , 103 μ and 35 μ mesh, nylon screen. After filtration these "deck" filters are washed into a container with 840ml of water from the ship's sea water system. An 84ml aliquote from this concentrate is further filtered on tared Millipore filters. These latter filters are then dried and reweighed. The weight gained is a measure of the material present in the original volume of sea water filtered on the nylon mesh "deck" filters. These values, however, must be "corrected" by subtraction of a "blank". The blank is a tared Millipore filter through which an aliquote from an equal volume of wash water (840ml) from the ship's sea water system has been filtered. Thus, the wash water contained microzooplankton and/or detritus from the upper 1 meter (the approximate depth of the intake for the sea water system) and probably detritus from the plumbing of the ship (the RV HORIZON was over 20 years old at the time). In most cases these blanks were equal to or greater than the values of the samples themselves. Therefore in only one case, the 2148-2300 (local time) series on 22 September 1968, was the blank low enough to yield useable sample values. The sampling technique used for this series was as

follows: the pump was raised slowly from 200m at the rate of 5m/min. to 125m. At this time the deck filters were removed and washed. The pumping rate during the ascent was 133 l/min. The same procedure was followed for the depths 125-75m, 75-50m, 50-25m and 25-0m. In both the depth measurements and volume calculations residence time of the water in the overboard pumping system was accounted for.

Photosynthetic Rate

The uptake of radioactive carbon by natural populations was determined by the procedure outlined by Strickland and Parsons (1968; Sec. V.3).

The vertical distribution of light in the water column was measured with a submarine photometer (Austin and Lauder milk, 1968), supplemented with secchi disc measurements to estimate the depth of the 1% light level.

Water samples were collected with Lexan bottles from depths reached by a specific percent of surface radiation. Subsamples of 250ml were inoculated with 20-25 μ c of C^{14} as bicarbonate. These samples were placed in deck incubators provided with neutral density filters to simulate in situ light intensities at six depths and cooled with surface sea water (Owen and Zeitzschel, 1970). At each intensity there were duplicate light bottles and a dark control. Samples were generally incubated for 6 hours from approximately noon. Samples were subsequently filtered onto Millipore filters, washed, dried, and their radioactivity determined with a Geiger counter. The production values presented in this report are the means of the two replicate determinations and have been corrected for dark "uptake".

Pigments

Chlorophyll-a and phaeophytin were determined from 550ml water samples according to the fluorometric procedure outlined by Strickland and Parsons (1968; Sec. IV. 3). The fluorometer was a Turner model #111, with a red sensitive photomultiplier and a blue lamp (Turner T-5 lamp #110-853). The instrument was calibrated against a spectrophotometer, using near surface populations and correcting the SCOR/UNESCO trichromatic equations for the presence of phaeophytin.

HYDROGRAPHIC DATA

These physical and chemical data were collected in part and processed by the Data Collection and Processing Group (DCPG,

MLR), Scripps Institution of Oceanography, University of California at San Diego.

Five Nansen bottle casts with 18 or fewer bottles were lowered to 1500 meters or less. The Nansen bottles contained paired protected reversing thermometers; unprotected reversing thermometers were included in 12 of the bottles.

Fifteen STD lowerings to 500m and 2 STD lowerings to 100m or less were made.

Water samples from the microzooplankton pumping device were taken for nutrient, chlorophyll-a and phaeo-pigment analyses at 4 "pump" stations.

Water samples for chemical measurements were obtained from the Nansen bottles. Salinity was determined by inductive salinometer. Dissolved oxygen was determined by the Winkler method as revised by J. H. Carpenter (1965). Phosphate, silicate, and nitrate determinations were made with the Technicon Autoanalyzer by methods suggested by Strickland and Parsons (1968).

TABULATED DATA

Data presented in this report were obtained by Nansen bottle casts, by analysis of water samples obtained by a pumping device, and by the in situ Salinity/Temperature/Depth Monitoring and Recording System (STD).^{1/} The data appear in three forms:

1. Data from the Nansen bottle casts are tabulated with values at observed depths on the left side of the page and with interpolated and computed values at standard depths on the right.

2. Data from stations with only nutrient and pigment values are tabulated at standard depths and centered on the page.

3. For each STD lowering, temperature and salinity values are read at standard depths only and appear with computed values on the right side of the page. Corrections may have been applied to the temperature or salinity values from continuing comparison of sample bottle data and STD data collected on the same station.

The time listed under "messenger time" for STD lowerings is "start down" time. Fathometer readings were not recorded except for the first hydrographic cast.

^{1/} In situ Salinity/Temperature/Depth Monitoring and Recording System, Model 9006, Tech. Rep. No. 102, HYTECH Marine Products, The Bissett-Berman Corporation.

The column headings from the computer are explained as follows:

Z	Depth	meters
T	Temperature	°C
S	Salinity	‰
O2	Oxygen	ml/L
PO4	Phosphate	µg at/L
SiO3	Silicate	µg at/L
NO3	Nitrate	µg at/L
DT	δT	cl/ton
SIGT	σt	g/L
DD	ΔD	dyn. m
CHLA	Chlorophyll-a	mg/m ³
PHAE	Phaeophytin	mg/m ³

STANDARD PROCEDURES

In situ Salinity/Temperature/Depth Recorder

The manufacturer of the STD claims an accuracy of $\pm 0.05^{\circ}\text{C}$ with repeatability of $\pm 0.01^{\circ}\text{C}$ for temperature and an accuracy of $\pm 0.03\text{‰}$ with repeatability of $\pm 0.01\text{‰}$ for salinity.

The data were digitized at Standard depths from the analog recording. Temperature data from the STD lowerings and Nansen bottle temperature data agreed closely so that no correction to the STD temperature data was necessary. No correction to the STD salinity data shallower than 150m was necessary. At depths greater than 150m a correction of 0.04‰ was applied to the STD salinity data.

Hydrographic Casts

The observed data have been plotted and then evaluated using the method described by Klein.^{1/} This involves consideration of their variation as functions of density or depth and their relations to each other, and comparison with previous or adjacent observations.

To indicate degree of accuracy, temperature is recorded in hundredths of a degree. Salinity, when determined by salinometer, is recorded to three decimal places, provided it meets accepted standards. The values are recorded to two decimal places when only one determination per sample was obtained, or where there is doubt

^{1/} Klein, Hans T. A new technique for processing physical oceanographic data. SIO Ref. 73-14.

concerning the accuracy of a particular sample, or of all samples on a station. Due to inexperienced personnel operating the salinometer during this leg of the cruise only the values for Cast 6 were considered to have the usual accuracy.

FOOTNOTES

In addition to footnotes, one special notation is used without a footnote because the meaning is always the same. Values which seem to be in error without apparent reason are indicated by the following notation:

u: uncertain value

LITERATURE CITED

- Austin, R. W. and R. W. Loudermilk, 1968. An oceanographic illuminometer for light penetration and reflection studies. SIO Ref. 68-11, Univ. of Calif., Scripps Inst. of Oceanog.
- Beers, J. R., G. L. Stewart and J. D. H. Strickland, 1967. A pumping system for sampling small plankton. J. Fish. Res. Board of Canada 24(8): 1811-1818.
- McGowan, J. A. and D. M. Brown, 1966. A new opening-closing paired zooplankton net. SIO Ref. 66-23, Univ. of Calif., Scripps Inst. of Oceanog.
- McGowan, J. A., 1971. Oceanic biogeography of the Pacific. Section A. In The Micropalaeontology of Oceans, edited by B. M. Funnell and W. R. Riedel. Cambridge University Press. Cambridge. pp 3-75.
- McGowan, J. A. and P. L. Williams, 1973. Oceanic habitat differences in the North Pacific. J. Exp. Mar. Biol. Ecol., Vol. 12 pp 187-217.
- Owen, R. W. and B. Zeitzschel, 1970. Phytoplankton production: seasonal change in the Oceanic Eastern tropical Pacific. Marine Biology Vol. 7 pp 32-36.
- Strickland, J. D. H. and T. R. Parsons, 1968. A practical handbook of seawater analysis. Fish Res. Board of Canada, Bull No. 167 311pp.

FIGURES

1. Station positions
2. Parachute drogue tracks
3. Bongo net tows

TABLES

1. Wind Velocity
2. Drogue data
3. Bongo tow and macrozooplankton biomass
4. Water Transparency
5. Primary productivity
6. Chlorophyll-a and Phaeophytin
7. Chlorophyll profile data
8. Bird and fish sightings
9. Fish catches




FIGURE 1

PERSONNEL

SHIP'S CAPTAIN

Ferris, Noel, RV Argo

PERSONNEL PARTICIPATING IN THE COLLECTION OF DATA

RV Argo

McGowan, J. A., Chief Scientist, Assoc. Prof. S.I.O.
Barnett, A., 2nd year Graduate Student
Clutter, R. I., Assoc. Prof., Univ. of Hawaii
Gopalakrishnan, K., 1st year Graduate Student
Haury, L., 1st year Graduate Student
Hurley, A., 1st year Graduate Student
Judkins, D., 2nd year Graduate Student
Kamykowski, D., 1st year Graduate Student
Klapow, L., 2nd year Graduate Student
Mauck, W., Marine Technician
Renz, G., 1st year Graduate Student
Rosendahl, R., Marine Technician
Smith, G., 1st year Graduate Student
Venrick, E., Post Grad. Res. Oceanog. S.I.O.
Wormuth, J., 2nd year Graduate Student
Yoshioka, P., 1st year Graduate Student

HYDROGRAPHIC DATA

RV ARGO				CLIMAX I EXPEDITION										STD 1			
LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES	
26 58.0N		155 24.0W		09/19/68		2330		GMT		090		15KT		1		030 04	
Z	T	S	O2	PO4	SI03	NO2	NO3	DT	Z	T	S	O2	SIGT	DT	DN		
									0	26.93	34.96		22.731	513.1			
									10	26.82	34.98		22.781	508.3	.051		
									20	26.82	35.07		22.849	501.8	.102		
									30	26.57	35.09		22.943	492.8	.151		
									50	23.73	35.13		23.840	407.2	.242		
									75	21.50	35.14		24.483	345.9	.317		
									100	20.49	35.07		24.704	324.8	.421		
									125	19.39	35.02		24.955	300.9	.500		
									150	18.38	34.91		25.128	284.5	.575		
									200	16.33	34.73		25.483	250.7	.711		
									250	14.04	34.44		25.765	223.9	.833		
									300	12.26	34.28		26.000	201.6	.941		
									400	9.61	34.11		26.343	169.0	1.136		
									500	7.39	34.02		26.614	143.9	1.300		

RV ARGO		CLIMAX I EXPEDITION										PUMP 1	
LATITUDE	LONGITUDE	MO/DAY/YR	MESSENGER	TIME	BOTTOM	WIND	SPEED	WEATHER	DOMINANT WAVES				
26 59.0N	155 31.5W	09/20/68	0315	GMT		100	17KT	I	100 03				
	Z	PO4	SI03	NO2	NO3	NH4	CHLA	PHAE					
	0A	.10	5.										
	10	.04	5.										
	25	.04	6.										
	35	.06	5.										
	50	.06	5.										
	60	.06	6.										
	75	.06	5.										
	100	.08	5.										
	125	.10	6.										
	150	.26	9.										
	200	.34	11.										

RV ARGO					CLIMAX I EXPEDITION										STD 2			
LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER		TIME	BOTTOM	WIND	SPEED	WEATHER	DOMINANT WAVES					
26 59.0N		155 31.5W		09/20/68		0510		GMT		100	14KT	1	110 04					
Z	T	S	O2	PO4	SI03	NO2	NO3	DT	Z	T	S	O2	SIGT	DT	DN			
									0	26.94	34.98		22.743	511.9	0			
									10	26.98	35.09		22.813	505.3	.051			
									20	26.64	35.10		22.929	494.2	.101			
									30	26.00	35.17		23.183	470.0	.149			
									50	23.18	35.11		23.985	393.4	.236			
									75	21.58	35.11		24.438	350.2	.329			
									100	20.60	35.11		24.705	324.8	.415			
									125	19.43	35.04		24.960	300.5	.494			
									150	18.31	34.94		25.168	280.7	.568			
									200	15.94	34.64		25.503	248.8	.713			
									250	13.43	34.41		25.868	214.1	.822			
									300	12.04	34.28		26.042	197.6	.928			
									400	9.51	34.14		26.383	165.2	1.117			
									500	7.22	34.02		26.638	141.0	1.278			

RV ARGO			CLIMAX I EXPEDITION											CAST 1		
LATITUDE 26 56.0N		LONGITUDE 155 36.0W		MO/DAY/YR 09/20/68		MESSENGER 1945		TIME GMT	BOTTOM 5661M	WIND 120	SPEED 04KT	WEATHER 1	DOMINANT WAVES 080 06			
Z	T	S	O2	PO4	SI03	NO2	NO3	DT	Z	T	S	O2	SIGT	DT	DN	
2		34.99	4.55	.64	5.		.3		0			4.55				
11		34.99	4.71	.12	4.		.5		10			4.70				
22		35.04	4.77	.34	4.		.3		20			4.77				
33		35.07	4.90U	.08	4.		.2		30			4.63				
51		35.11	5.31U	.19	5.		.3		50			4.38				
76		35.12	4.25	.08	5.		.3		75			4.25				
100	20.15	35.10	5.07	.08	6.		.8	314.1	100	20.15	35.10	5.07	24.817	314.1		
125	19.12	35.04	4.83	.22	8.		1.9	292.9	125	19.12	35.04	4.83	25.040	292.9		
150	18.05	34.94	4.60U	.27	10.		2.8	274.6	150	18.05	34.94	4.76	25.232	274.6		
199	16.26	34.69	4.61	.27	12.		4.1	252.1	200	16.22	34.69	4.62	25.474	251.6		
247	14.40	34.51	4.89	.56	16.		6.6	226.0	250	14.28	34.50	4.89	25.758	224.5		
296	12.59	34.33	4.81	.89	21.		9.4	204.0	300	12.47	34.32	4.81	25.990	202.5		
395	10.08	34.16	4.71	1.40	31.		14.3	172.8	400	9.95	34.15	4.66	26.319	171.3		
492	7.62	34.03	3.61	2.85	48.			145.7	500	7.44	34.03	3.57	26.611	143.5		
590	5.81	34.04	3.20					121.8	600	5.70	34.05	3.13	26.860	119.9		
788	4.59	34.26		3.02	85.			91.8	700	4.90	34.15	2.42	27.039	103.0		
987	3.91	34.42	.62		91.			72.9	800	4.54	34.27	1.76	27.172	90.4		
1481	2.80	34.56	1.30U	3.47	102.			52.3	1000	3.87	34.43	.64	27.367	71.9		
									1200	3.28	34.52	.40	27.497	59.5		
									1500	2.79	34.56	1.30	27.575	52.2		

A) THESE PHOSPHATE AND SILICATE SAMPLES WERE DRAWN FROM MICROZOOPLANKTON PUMP WATER.

RV ARGO

CLIMAX I EXPEDITION

STD 3

LATITUDE 26 55.5N			LONGITUDE 155 36.0W			MO/DAY/YR 09/20/68		MESSENGER TIME 2105			BOTTOM	WIND 150	SPEED 05KT	WEATHER 1	DOMINANT WAVES 080 05		
Z	T	S	O2	P04	S103	N02	N03	DT	Z	T	S	O2	SIGT	DT	DD		
									0	26.82	35.01			22.804	506.1	0	
									10	26.76	35.01			22.823	504.3	.051	
									20	26.76	35.04			22.846	502.2	.101	
									30	26.77	35.09			22.880	498.9	.151	
									50	24.12	35.10			23.702	420.4	.243	
									75	21.64	35.09			24.406	353.2	.341	
									100	20.19	35.05			24.769	318.7	.425	
									125	19.19	35.02			25.007	296.0	.503	
									150	17.96	34.90			25.224	275.4	.576	
									200	16.08	34.60			25.441	254.8	.711	
									250	14.09	34.44			25.755	224.9	.834	
									300	12.34	34.30			26.000	201.6	.945	
									400	9.87	34.14			26.323	170.9	1.139	
									500	7.42	34.03			26.618	142.9	1.304	

RV ARGO

CLIMAX I EXPEDITION

STD 4

LATITUDE 26 55.5N			LONGITUDE 155 36.0W			MO/DAY/YR 09/21/68		MESSENGER TIME 0700			BOTTOM	WIND 110	SPEED 12KT	WEATHER 1	DOMINANT WAVES 080 04		
Z	T	S	O2	P04	S103	N02	N03	DT	Z	T	S	O2	SIGT	DT	DD		
									0	26.85	35.04			22.817	504.9	0	
									10	26.80	35.05			22.840	502.7	.050	
									20	26.80	35.09			22.871	499.8	.101	
									30	26.76	35.10			22.891	497.8	.151	
									50	23.88	35.11			23.781	412.9	.242	
									75	21.70	35.11			24.405	353.4	.338	
									100	20.57	35.13			24.728	322.5	.424	
									125	19.81	35.09			24.899	306.3	.503	
									150	18.72	34.96			25.080	289.0	.579	
									200	16.18	34.65			25.456	253.3	.717	
									250	13.65	34.36			25.785	222.1	.839	
									300	12.27	34.28			25.998	201.8	.949	
									400	9.68	34.13			26.347	168.6	1.142	
									500	7.77	34.04			26.575	147.0	1.308	

RV ARGO

CLIMAX I EXPEDITION

STD 5

LATITUDE 27 00.0N			LONGITUDE 155 50.0W			MO/DAY/YR 09/21/68		MESSENGER TIME 1803			BOTTOM	WIND 120	SPEED 12KT	WEATHER 1	DOMINANT WAVES 110 03		
Z	T	S	O2	P04	S103	N02	N03	DT	Z	T	S	O2	SIGT	DT	DD		
									0	26.80	35.02			22.818	504.8	0	
									10	26.80	35.02			22.818	504.8	.051	
									20	26.85	35.10			22.862	500.6	.101	
									30	26.83	35.10			22.869	500.0	.151	
									50	24.70	35.06			23.498	439.8	.245	
									75	21.73	35.08			24.374	356.3	.345	
									100	20.53	35.10			24.716	323.7	.431	
									125	19.27	35.01			24.978	298.7	.510	
									150	17.91	34.90			25.236	274.2	.583	
									200	16.15	34.70			25.501	249.0	.716	
									250	13.88	34.44			25.799	220.7	.837	
									300	12.51	34.36			26.013	200.3	.946	
									400	9.57	34.12			26.357	167.6	1.138	
									500	7.57	34.01			26.581	146.5	1.303	

RV ARGO

CLIMAX I EXPEDITION

STD 6

LATITUDE 27 05.3N			LONGITUDE 155 50.0W			MO/DAY/YR 09/22/68		MESSENGER TIME 0652			BOTTOM	WIND 150	SPEED 06KT	WEATHER 1	DOMINANT WAVES 120 03		
Z	T	S	O2	P04	S103	N02	N03	DT	Z	T	S	O2	SIGT	DT	DD		
									0	27.35	34.97			22.604	525.2	0	
									10	26.96	35.00			22.752	511.1	.052	
									20	26.89	35.05			22.812	505.4	.103	
									30	26.92	35.09			22.832	503.4	.153	
									50	25.65	35.21			23.321	456.7	.250	
									75	22.17	35.17			24.319	361.5	.352	
									100	20.59	35.09			24.693	325.9	.439	
									125	19.50	35.04			24.942	302.2	.519	
									150	18.27	34.94			25.178	279.7	.593	
									200	16.37	34.73			25.473	251.6	.728	
									250	13.72	34.46			25.847	216.1	.849	
									300	11.98	34.26			26.038	198.0	.956	
									400	9.55	34.12			26.361	167.3	1.146	
									500	7.43	34.03			26.616	143.1	1.309	

RV ARGO		CLIMAX I EXPEDITION							PUMP 2	
LATITUDE	LONGITUDE	MO/DAY/YR	MESSENGER TIME		BOTTOM	WIND	SPEED	WEATHER	DOMINANT WAVES	
27 05.5N	155 50.0W	09/22/68	0723	GMT		150	05KT	1	02	
	Z	P04	S103	NO2	NO3	NH4	CHLA	PHAE		
	0A	.05	5.				.03	.01		
	10	.05	6.				.02	.03		
	25	.05	5.				.02	.01		
	35	.05	5.				.03	.02		
	50	.06	7.				.04	.02		
	60	.07	5.				.05	.03		
	75	.03	6.				.07	.03		
	100	.06	8.				.12	.16		
	125	.23	11.				.12	.17		
	150	.16	8.				.04	.07		
	200	.33	13.				.01	.02		

RV ARGO			CLIMAX I EXPEDITION										STD 7					
LATITUDE 27 03.7N			LONGITUDE 155 56.8W		MO/DAY/YR 09/22/68		MESSENGER TIME 2157		BOTTOM		WIND 120		SPEED 04KT		WEATHER 1		DOMINANT WAVES 080 03	
Z	T	S	O2	P04	S103	NO2	NO3	CT	Z	T	S	O2	SIGT	DT	DD			
									0	27.18	35.03		22.704	515.7	0			
									10	26.92	35.04		22.795	507.0	.051			
									20	26.93	35.07		22.814	505.2	.102			
									30	26.92	35.08		22.825	504.1	.152			
									50	25.75	35.07		23.185	469.7	.250			
									75	21.88	35.12		24.362	357.4	.354			
									100	20.44	35.13		24.763	319.2	.439			
									125	19.23	35.03		25.004	296.3	.517			
									150	18.20	34.94		25.195	278.1	.590			
									200	16.38	34.70		25.448	254.0	.726			
									250	14.15	34.46		25.757	224.6	.849			
									300	12.38	34.33		26.015	200.1	.959			
									400	9.71	34.12		26.334	169.9	1.152			
									500	7.50	34.02		26.598	144.8	1.317			

RV ARGO			CLIMAX I EXPEDITION											STD 8	
LATITUDE 27 12.0N		LONGITUDE 156 05.0W		MO/DAY/YR 09/23/68		MESSENGER 0610		TIME GMT	BOTTOM	WIND 130	SPEED 10KT	WEATHER 1	DOMINANT WAVES 100 02		
Z	T	S	O2	P04	S103	NO2	NO3	DT	Z	T	S	O2	SIGT	DT	DD
									0	27.49	35.00		22.582	527.4	0
									10	26.96	35.03		22.774	509.0	.052
									20	26.94	35.03		22.781	508.4	.103
									30	26.92	35.06		22.810	505.6	.154
									50	26.15	35.07		23.060	481.6	.253
									75	22.64	35.07		24.110	381.4	.361
									100	20.80	35.10		24.644	330.6	.451
									125	19.60	35.04		24.916	304.7	.531
									150	18.57	34.94		25.103	286.9	.606
									200	16.25	34.69		25.471	251.9	.744
									250	13.78	34.35		25.750	225.3	.867
									300	12.12	34.28		26.027	199.0	.976
									400	9.47	34.10		26.358	167.6	1.167
									500	7.52	34.02		26.596	145.0	1.331

RV ARGO		CLIMAX I EXPEDITION							PUMP 3	
LATITUDE	LONGITUDE	MO/DAY/YR	MESSENGER TIME		BOTTOM	WIND	SPEED	WEATHER	DOMINANT WAVES	
27 12.5N	156 04.5W	09/23/68	0951	GMT		090	06KT	1	080 03	
	Z	P04	S103	NO2	NO3	NH4	CHLA	PHAE		
	0A	.11	4.				.05	.02		
	10	.09	4.				.02	.02		
	25	.09	4.				.02	.02		
	35	.09	4.				.06	.07		
	50	.07	5.							
	60	.05	5.							
	75	.07	6.				.06	.04		
	100	.12	6.				.14	.27		
	110	.14	7.				.18	.21		
	125	.16	7.				.11	.21		
	150	.28	10.				.02	.05		
	200	.31	13.				.01	.02		

A) THESE CHLOROPHYLL A AND PHAEOPHYTIN SAMPLES WERE DRAWN FROM MICROZOOPLANKTON PUMP WATER.

ST010

STD12

CAST 3

ST013

RV ARGO					CLIMAX I EXPEDITION										STD14			
LATITUDE 27 33.0N			LONGITUDE 156 14.0W		MO/DAY/YR 09/24/68		MESSENGER 1650		TIME GMT		BOTTOM	WIND 030	SPEED 06KT	WEATHER 6	DOMINANT WAVES 360 06			
Z	T	S	02	PD4	SD3	NC2	NC3	DT	Z	T	S	02	SIGT	DT	DD			
									0	27.18	35.06		22.727	513.5	0			
									10	27.04	35.08		22.787	507.8	.051			
									20	26.97	35.08		22.809	505.7	.102			
									30	26.96	35.10		22.827	503.9	.152			
									50	26.25	35.10		23.052	482.5	.251			
									75	21.73	35.10		24.389	354.9	.357			
									100	20.56	35.12		24.723	323.0	.442			
									125	19.55	35.06		24.944	302.0	.521			
									150	18.38	34.97		25.173	280.2	.595			
									200	16.10	34.67		25.490	250.1	.731			
									250	13.83	34.47		25.832	217.5	.851			
									300	12.34	34.31		26.008	200.9	.959			
									400	9.88	34.16		26.337	169.6	1.152			
									500	7.39	34.04		26.630	141.8	1.316			

RV ARGO					CLIMAX I EXPEDITION										STD16		
LATITUDE 27 40.0N			LONGITUDE 156 21.0W		MO/DAY/YR 09/24/68		MESSENGER 2323		TIME GMT	BCTDM	WIND 090	SPEED 08KT	WEATHER 1	DOMINANT WAVES 350 06			
Z	T	S	02	PD4	SD03	NC2	NC3	DT	Z	T	S	02	SIGT	DT	DD		
									0	27.32	34.87		22.539	531.5	0		
									10	27.08	35.07		22.766	509.8	.052		
									20	26.99	35.09		22.810	505.6	.103		
									30	26.97	35.11		22.831	503.5	.153		
									50	25.07	35.12		23.432	446.2	.249		
									75	22.01	35.13		24.334	360.1	.350		
									100	20.88	35.12		24.637	331.2	.437		
									125	19.78	35.10		24.915	304.8	.518		
									150	18.58	35.00		25.146	282.7	.593		
									200	16.30	34.68		25.451	253.7	.729		
									250	13.58	34.46		25.876	213.4	.849		
									300	11.92	34.32		26.096	192.5	.954		
									400	9.32	34.12		26.398	163.8	1.140		
									500	7.51	34.06		26.628	141.9	1.301		

RV ARGO					CLIMAX I EXPEDITION										CAST 3					
LATITUDE 27 41.0N			LONGITUDE 156 25.5W		MO/DAY/YR 09/25/68		MESSENGER 0708		TIME GMT		BOTTOM		WIND 070		SPEED 09KT		WEATHER 2		DOMINANT WAVES 360 06	
Z	T	S	02	PD4	SD3	N02	N03	DT	Z	T	S	02	SIGT	DT	DD					
1	27.01	34.54	4.80U	.22	4.		.3	545.7	0	27.01	34.54		22.390	545.7		0				
11	26.96	34.98	4.79U	.17	4.		.2	512.6	10	26.97	34.95		22.714	514.7		.053				
21	26.89	35.05	4.42	.19	4.		.3	505.4	20	26.90	35.05	4.42	22.809	505.6		.104				
31		35.06	4.77	.17	4.		.2		30	26.72	35.07	4.72	22.882	498.6		.154				
50	25.49	35.08	5.33		4.		.3	461.3	50	25.49	35.08	5.33	23.273	461.3		.251				
75	21.69	35.19	5.13	.31	5.		.3	347.3	75	21.69	35.19	5.13	24.468	347.3		.352				
93	20.47	35.10	5.07		6.		.4	322.2	100	20.42	35.10	5.06	24.745	320.9		.437				
124	19.38	35.12	4.76	.79	7.		1.0	293.4	125	19.34	35.12	4.76	25.043	292.6		.514				
149	18.40	35.06	4.67	.29	9.		3.1	274.1	150	18.36	35.06	4.67	25.246	273.3		.586				
198	16.50	34.97	4.95	.48	11.		3.1	237.0	200	16.43	34.96	4.95	25.638	236.0		.717				
246	14.64	34.73	5.00	.51	15.		5.3	214.8	250	14.46	34.71	4.99	25.885	212.5		.832				
295	12.45	34.52	4.83	.68	20.		9.9	187.4	300	12.28	34.51	4.83	26.171	185.3		.935				
394	9.67	34.30	4.75	1.22	32.		16.8	155.9	400	9.52	34.29	4.71	26.496	154.5		1.113				
492	7.47	34.12	3.71	2.02	49.		23.5	136.9	500	7.31	34.11	3.56	26.696	135.5		1.266				
589	5.79	34.04	1.89	2.96	66.		29.2	121.6	600	5.68	34.04	1.76	26.855	120.4		1.401				
787	4.62	34.06	.64	3.30	83.		31.2	107.1	700	4.92	34.06	.90	26.947	111.7		1.525				
984	3.95	34.07	.79	4.03U	91.		31.4	99.6	800	4.57	34.06	.65	27.002	106.5		1.642				
1473	2.84	34.04	1.58	3.43	102.		32.0	91.9	1000	3.90	34.07	.82	27.079	99.2		1.863				
									1200	3.30	34.05	1.17	27.122	95.1		2.074				
									1500	2.83	34.04	1.59	27.157	91.8		2.380				

RV ARGO CLIMAX I EXPEDITION STD17

LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES	
27 56.0N		156 30.5W		09/25/68		1637		GMT		060		12KT				360 05	
Z	T	S	O2	PD4	SI03	NO2	NO3	DT	Z	T	S	O2	SIGT	DT	DD		
									0	26.98	34.84		22.625	523.2	0		
									10	26.98	34.84		22.625	523.2	.052		
									20	27.03	34.95		22.692	516.9	.104		
									30	27.01	35.10		22.811	505.5	.156		
									50	25.30	35.14		23.376	451.5	.252		
									75	22.25	35.10		24.243	368.7	.355		
									100	20.75	35.16		24.703	325.0	.442		
									125	19.52	35.10		24.982	298.3	.521		
									150	18.38	35.00		25.196	278.0	.594		
									200	16.19	34.72		25.507	248.4	.729		
									250	13.80	34.47		25.838	216.9	.848		
									300	12.12	34.33		26.065	195.4	.955		
									400	9.73	34.18		26.378	165.7	1.143		
									500	7.48	34.06		26.633	141.5	1.305		

RV ARGO CLIMAX I EXPEDITION STD18

LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES	
28 05.0N		156 36.0W		09/26/68		0458		GMT		060		09KT		1		010 05	
Z	T	S	O2	PD4	SI03	NO2	NO3	DT	Z	T	S	O2	SIGT	DT	DD		
									0	27.17	34.78		22.519	533.4	0		
									10	27.10	34.78		22.542	531.2	.053		
									20	27.05	34.91		22.655	520.3	.106		
									30	27.02	35.00		22.733	513.0	.158		
									50	26.72	35.04		22.858	500.9	.259		
									75	21.87	35.05		24.312	362.2	.368		
									100	20.60	35.09		24.690	326.2	.455		
									125	19.55	35.02		24.914	304.9	.535		
									150	18.40	34.92		25.130	284.3	.609		
									200	16.27	34.66		25.443	254.5	.747		
									250	14.35	34.46		25.715	228.7	.871		
									300	12.45	34.26		25.948	206.6	.983		
									400	9.72	34.12		26.332	170.0	1.180		
									500	7.53	34.00		26.578	146.7	1.346		

RV ARGO CLIMAX II EXPEDITION CAST 4

LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES	
28 05.0N		156 36.0W		09/26/68		0545		GMT		050		12KT		1		110	
Z		PD4	SI03	NO2	NO3	NH4	CHLA	PHAE									
	0	.09	4.		.1												
	20	.10	5.		.1						.03		.01				
	40	.10	4.		.1						.03		.01				
	60	.10	5.		.0						.04		.02				
	80	.12	5.		.0						.05		.05				
	90	.14	7.		.0						.05		.10				
	100	.17	7.		.3						.14		.18				
	110	.22	7.		1.0						.14		.20				
	120	.24	8.		2.0						.10		.14				
	140	.29	9.		2.9												
	160	.31	10.		3.1						.02		.02				
	180	.36	10.		3.3						.01		.02				
	200	.32	11.		2.9						.01		.02				
	230	.39	14.		4.5						.01		.01				
	260	.63	18.		7.7						.00		.01				
	290	.73	20.		9.1						.00		.01				
	320	.92	24.		11.2						.00		.01				
	350	1.07	27.		13.3						.00		.00				

RV ARGO										CLIMAX I EXPEDITION										STD19			
LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES							
28 17.0N		157 12.0W		09/26/68		1802				060		12KT		1		040 04							
Z	T	S	O2	PD4	S103	NC2	NC3	DT	Z	T	S	O2	S1.T	DT	DT								
									0	27.01	34.79			22.578	527.7		0						
									10	27.01	34.79			22.578	527.7		.053						
									20	27.04	34.91			22.659	520.0		.105						
									30	26.88	35.01			22.785	508.0		.157						
									50	24.60	35.06			23.528	436.9		.252						
									75	21.58	35.05			24.392	354.5		.351						
									100	20.54	35.06			24.683	326.8		.437						
									125	19.37	35.00			24.945	301.9		.517						
									150	18.35	34.88			25.112	286.0		.591						
									200	16.53	34.70			25.413	257.3		.730						
									250	14.24	34.41			25.700	230.1		.855						
									300	12.00	34.22			26.003	201.3		.967						
									400	9.69	34.10			26.322	171.0		1.160						
									500	7.18	33.97			26.604	144.2		1.326						

RV ARGON				CLIMAX I EXPEDITION										CAST 5									
LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES							
28 17.0N		157 12.0W		09/26/68		2057				060		04KT		2		050 05 08							
Z	T	S	O2	PD4	S103	NC2	NC3	DT	Z	T	S	O2	S1GT	DT	DD								
1	27.02	34.85						523.7	0	27.02	34.85			22.620	523.7	0							
11	26.93	34.84						521.7	10	26.94	34.84			22.639	521.9	.052							
21	27.04	35.02						512.1	20	27.03	34.99			22.726	513.6	.104							
31		35.07							30	26.11	35.07			23.076	480.2	.154							
50	23.18	35.10						394.1	50	23.18	35.10			23.978	394.1	.242							
75	20.87	35.14						329.5	75	20.87	35.14			24.655	329.5	.333							
98	19.80	35.08						306.7	100	19.71	35.07			24.915	304.7	.413							
123	18.69	35.01						284.7	125	18.62	35.00			25.139	283.4	.487							
148	17.86	34.92						271.6	150	17.78	34.91			25.275	270.5	.558							
197	15.84	34.65						245.9	200	15.70	34.63			25.552	244.1	.689							
246	13.59	34.41						217.2	250	13.42	34.39			25.858	215.1	.807							
294	11.76	34.26						194.0	300	11.61	34.25			26.102	191.9	.912							
393	9.78	34.18						166.5	400	9.62	34.17			26.386	164.9	1.098							
491	7.66	34.04						145.5	500	7.48	34.04			26.614	143.3	1.260							
589	5.94	34.05						122.6	600	5.81	34.06			26.854	120.5	1.400							
787	4.50	34.24						92.4	700	4.92	34.15			27.032	103.6	1.520							
985	3.85	34.40						73.8	800	4.44	34.25			27.167	90.9	1.625							
1474	2.82	34.55							53.2	1000	3.80	34.40			27.351	73.4	1.805						
										1200	3.25	34.43			27.425	66.4	1.962						
										1500	2.81	34.58			27.589	50.9	2.165						

RV ARGO										CLIMAX I EXPEDITION										STD20			
LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER TIME		BOTTOM		WIND		SPEED		WEATHER		DOMINANT WAVES						DD	
28 36.0N		157 52.0W		09/28/68		0024				070		05KT		1		030 04							
Z	T	S	O2	PD4	S103	NC2	NC3	DT	Z	T	S	O2	S1GT	DT	DT								
									0	27.33	34.83			22.505	534.7		0						
									10	27.02	34.84			22.612	524.5		.053						
									20	27.00	34.85			22.626	523.1		.105						
									30	27.09	35.03			22.733	512.9		.157						
									50	25.25	35.10			23.362	452.9		.254						
									75	21.35	35.05			24.456	348.5		.355						
									100	20.04	35.04			24.801	315.6		.439						
									125	18.79	34.96			25.063	290.7		.516						
									150	17.93	34.87			25.208	276.8		.588						
									200	16.03	34.65			25.490	250.0		.722						
									250	13.57	34.36			25.801	220.5		.843						
									300	11.73	34.18			26.023	199.4		.951						
									400	9.34	34.03			26.324	170.7		1.144						
									500	7.56	33.98			26.558	148.6		1.312						

RV ARGO

CLIMAX I EXPEDITION

CAST 6

LATITUDE		LONGITUDE		MO/DAY/YR		MESSENGER		TIME	BOTTOM		WIND	SPEED	WEATHER	DOMINANT WAVES	
28 33.0N		156 52.0W		09/28/68		0457		GMT			070	06KT	1	030 04	
Z	T	S	02	PD4	SID3	NO2	NO3	DT	Z	T	S	02	SIGT	DT	CC
1	27.14	34.907	4.83	.10	4.		.3	523.3	0	27.14	34.907	4.83	22.624	523.3	0
10	26.98	35.002	4.75	.10	5.		.1	511.6	10	26.98	35.002	4.75	22.747	511.6	.052
19	26.92	35.007	4.71	.10	5.		.1	509.4	20	26.84	35.012	4.70	22.798	506.7	.101
28		35.077	5.25	.17	5.		.2		30	25.92	35.068	5.33	23.131	474.9	.152
46	23.78	35.187	5.66	.12	6.		.1	404.5	50	23.23	35.206	5.64	24.043	387.8	.239
70	20.84	35.253	5.33	.17	6.		.1	320.6	75	20.53	35.226	5.29	24.813	314.5	.327
92	19.72	35.096	5.13	.18	7.		.1	303.6	100	19.22	35.051	4.95	25.024	294.4	.404
115	18.35	34.985	4.67	.22	9.		.9	278.4	125	18.02	34.950	4.71	25.248	273.1	.476
139	17.66	34.906	4.76	.22	11.		1.5	267.9	150	17.36	34.881	4.90	25.356	262.8	.544
184	16.25	34.768	5.20	.18	12.		0.7	246.2	200	15.47	34.647	5.01	25.615	238.2	.672
228	14.05	34.446	4.61	.45	18.		4.7	223.7	250	13.16	34.380	4.61	25.899	211.2	.787
273	12.36	34.341	4.62	.65	23.		6.7	198.9	300	11.64	34.283	4.69	26.120	190.1	.891
361	10.21	34.159	4.84	1.05	32.		11.7	175.0	400	9.09	34.096	4.40	26.416	162.1	1.074
434	8.18	34.058	3.93	1.68	46.		17.2	151.4	500	7.06	34.029	3.27	26.660	138.3	1.232
530	6.69	34.027	2.99	2.48	60.		21.4	133.6	600	5.85	34.064	2.13	26.854	120.5	1.369
725	4.77	34.181	.87	3.42	86.		25.1	99.6	700	4.94	34.154	1.10	27.033	103.5	1.489
921	4.03	34.365	.68	3.76U	97.		26.1	78.2	800	4.35	34.252	.80	27.177	89.9	1.593

RV ARGO

CLIMAX I EXPEDITION

PUMP 4

LATITUDE	LONGITUDE	MO/DAY/YR	MESSENGER	TIME	BOTTOM	WIND	SPEED	WEATHER	DOMINANT WAVES
28 43.0N	156 49.0W	09/28/68	1020	GMT		070	04KT	1	
	Z	PD4	SID3	NO2	NO3	NH4	CHLA	PHAE	
	0	.12	5.				.03	.05	
	10	.38	5.				.03	.01	
	25	.12	6.				.03	.02	
	35	.12	4.				.03	.03	
	50	.05	5.				.04	.02	
	60	.12	6.				.05	.03	
	75	.13	8.				.06	.03	
	100	.33	7.				.16	.21	
	125	.23	7.				.14	.21	
	150	.25	10.				.06	.11	
	200	.37	13.				.01	.02	

A) THESE CHLOROPHYLL A AND PHAEOPHYTIN SAMPLES WERE DRAWN FROM MICROZOOPLANKTON PUMP WATER.

BIOLOGICAL DATA





FIGURE 2



BONGO NET TOWS
CLIMAX I
SEPTEMBER 1968

Table 1

CLIMAX I WIND VELOCITY

Date 1968	Local Time	Direction degrees	Speed knots
Sept. 19	0600	095	18
	1200	100	15
	1800	075	13
	2400	100	12
Sept. 20	0600	115	13
	1200	150	5
	1800	110	10
	2400	120	15
Sept. 21	0600	105	8
	1200	145	8
	1800	120	5
	2400	145	5
Sept. 22	0600	120	4
	1200	120	4
	1800	125	5
	2400	090	6
Sept. 23	0600	090	8
	1200	120	6
	1800	-	0
	2400	040	2
Sept. 24	0600	030	6
	1200	045	5
	1800	070	8
	2400	090	14
Sept. 25	0600	060	12
	1200	050	10
	1800	060	9
	2400	090	12
Sept. 26	0600	090	14
	1200	090	8
	1800	065	12
	2400	090	8
Sept. 27	0600	070	5
	1200	045	5
	1800	070	6
	2400	070	8
Sept. 28	0600	060	4
	1200	045	6
	1800	115	8
	2400	120	11

Table 2

CLIMAX I DROGUE DATA

Date 1968	Local Time	Interval hours:min.	Distance naut. mi.	Speed knots
Sept. 19	0540	-	-	-
	1804	12:24	14.2	1.15
Sept. 20	0550	11:46	6.0	0.51
	1500	9:10	6.1	0.67
Sept. 21	0040	7:40	6.9	0.90
	1300	12:20	5.0	0.41
	1840	5:40	5.0	0.88
Sept. 22	0923	12:43	5.8	0.46
	1250	3:27	2.5	0.72
	1835	5:45	6.0	1.04
Sept. 23	0230	5:55	7.3	1.23
	1300	10:30	6.4	0.61
	1840	5:40	5.3	0.94
Sept. 24	0030	5:50	6.9	1.18
	1200	11:30	9.0	0.78
	1845	6:45	4.8	0.71
Sept. 25	0030	5:45	6.9	1.20
	0550	5:20	7.9	1.48
	1200	6:10	8.3	1.35
	1836	6:36	3.5	0.53
Sept. 26	0015	5:39	4.1	0.73
	0555	5:40	5.0	0.88
	1200	6:05	4.8	0.79
	1840	6:40	5.0	0.75
Sept. 27	0005	5:25	5.8	1.07
	0550	5:45	6.0	1.39
	1200	6:10	7.3	1.18
	1835	6:35	9.0	1.37
Sept. 28	0000	5:25	4.5	0.83
	0550	5:50	4.1	0.70
	1200	6:10	8.2	1.33

Table 3

CLIMAX I BONGO TOW AND MACROZOOPLANKTON BIOMASS

Date 1968	Local Time	Expected Depth (m) Fished	Side of Net R or L	Overall Reliability	Comments	Depth (m) if differs from exp'd.	Linear Meters Fished	Disp. Vol. (ml)	Biomass ³ ml/1000 m ³
Sept. 19	1446-1525	25-0	L	useful	Net did not close		1540		
		25-0	R	good			1010	20	50
		50-25	L	useful	Net closed at surface	50-0	1664	25	38
		50-25	R	useful	Net closed at surface	50-0	1664	20	30
		75-50	L	poor	Came up open	75-0	2157	25	29
		75-50	R	poor	Came up c/s'd	75-0	1012	5	12
		100-75	L	poor	#2 (25-50m) therefore not opening net #3 (50- 75m) and #4 (75-100m) at expected depth.	100-75	1010	17.5	44
		100-75	R	poor		100-75	1010	17.5	44
Sept. 20	0326-0417	25-0	L	good			1015	65	162
		25-0	R	good			1015	32.5	81
		50-25	L	poor	Did not close	50-0	2465	52.5	54
		50-25	R	poor	Did not close	50-0	2465	45	46
		75-50	L	poor	Did not close	75-0	2850	42.5	38
		75-50	R	-	Did not open	-	-	-	-
		100-75	L	poor	Nets came up open. Although no mes- senger hit this frame, nets apparently opened during ascent.	100-0	3389		
		100-75	R	poor		100-0	3389		
	1345-1358	25-0	L	good			1360	10	19
		25-0	R	good			1360	15	28
		50-25	L	good			1000	10	25
		50-25	R	good			1000	17.5	44
		75-50	L	good			1010	<5	<10
		75-50	R	good		73	1010	20	50
	1421-1441	100-75	L	good			1015	12.5	31
		100-75	R	good			1015	12.5	31
	1623-1644	225-100	L	good			1015	10	25
		225-100	R	good			1015	5	12
		350-225	L	good			1015	12.5	31
		350-225	R	good			1015	15	37
		475-350	L	good			1012	2.5	6
		475-350	R	good			1012	10	25
		600-475	L	good		626	1010	15	38
		600-475	R	poor	Hole in cod end		1010	<5	<10
Sept. 21	0250-0325	25-0	L	good			1015	40	100
		25-0	R	good			1015	30	75
		50-25	L	good			1015	12.5	31
		50-25	R	good			1015	25	62
		75-50	L	poor	Hole in cod end		1012	<5	<10
		75-50	R	good			1012	25	62
		100-75	L	good		108	1010	<5	<10
		100-75	R	good			1010	20	50
	0529-0600	225-100	L	good			1015	5	12
		225-100	R	good			1015	15	37
		350-225	L	poor	Net did not close	350-0	?	20	
		350-225	R	poor	Net did not close	350-0	?	30	
		475-350	L	good			1012	7.5	19
		475-350	R	good			1012	10	25
		600-475	L	-	Nets did not open	518	-	-	-
		600-475	R	-	Nets did not open	-	-	-	-
	1405-1431	25-0	L	good			1070	15	35
		25-0	R	good			1070	25	59
		50-25	L	useful	Nets did not close	50-0	2865	30	28
		50-25	R	useful	Nets did not close	50-0	2865	40	35

Date 1968	Local Time	Expected Depth (m) Fished	Side of Net R or L	Overall Reliability	Comments	Depth (m) if differs from exp'd.	Linear Meters Fished	Disp. Vol. (ml)	Biomass ml/1000 m ³
		75-50	L	useful	Nets apparently worked but note large difference in L/R volumes		1012	2.5	6
		75-50	R	useful			1012	20	50
		100-75	L	good		104	1015	<5	12
		100-75	R	good			1015	<5	12
	1638-1656	225-100	L	good			1015	7.5	19
		225-100	R	good			1015	7.5	19
		350-225	L	good			1012	5	12
		350-225	R	good			1012	5	12
		475-350	L	good			1010	5	12
		475-350	R	good			1010	10	25
		600-475	L	good			1510	10	17
		600-475	R	good			1510	10	17
Sept. 22	0214-0234	225-100	L	good			1015	15	37
		225-100	R	good			1015	10	25
		350-225	L	good			1400	10	18
		350-225	R	good			1400	10	18
		475-350	L	poor	Hole in cod end		1012	<5	<10
		475-350	R	good			1012	5	12
		600-475	L	poor	Hole in cod end		1010	<5	<10
		600-475	R	good		612	1010	2.5	6
	0408-0445	25-0	L	good			1015	30	75
		25-0	R	good			1015	22.5	56
		50-25	L	good			1400	35	63
		50-25	R	good			1400	5	9
		75-50	L	poor	Hole in cod end		1012	<5	<10
		75-50	R	good			1012	30	75
		100-75	L	poor	Hole in cod end		1010	<5	<10
		100-75	R	good		102	1010	22.5	56
	1345-1417	25	L	useful	This cast did not "start up" until 10 min. after the messenger time, thus these are essentially horizontal tows at the maximum depths.		1015	17.5	44
		25	R	useful			1015	15	37
		50	L	useful			1400	35	63
		50	R	useful			1400	30	54
		75	L	useful			1012	<5	<10
		75	R	useful			1012	12.5	31
		100	L				975	-	-
		100	R				975	<5	<10
	1543-1615	225-100	L	good	Nets apparently worked but note large difference in L/R volumes		1015	7.5	19
		225-100	R	good			1015	20	50
		350-225	L	good			1400	5	9
		350-225	R	good			1400	7.5	14
		475-350	L	good			1012	<5	<10
		475-350	R	good			1012	5	12
		600-475	L	good			1010	<5	<10
		600-475	R	good			1010	<5	<10
Sept. 23	0247-0330	25-0	L	good			1510	15	25
		25-0	R	good			1510	27.5	46
		50-25	L	good			940	22.5	60
		50-25	R	-	Sample lost		940	-	-
		75-50	L	poor	Post trip--nets fished shallower layer	?	?	55	
		75-50	R	poor		?	?	<5	
		100-75	L	useful		25-0	243	7.5	78
		100-75	R	useful		25-0	243	5	
	0557-0630	225-100	L		Wire clamp failed, net slid down wire	?		5	
		225-100	R		Wire clamp failed, net slid down wire	?			

Date 1968	Local Time	Expected Depth (m) Fished	Side of Net R or L	Overall Reliability	Comments	Depth (m) if differs from exp'd.	Linear Meters Fished	Disp. Vol. (ml)	Biomass ml/1000m ³
		350-225	L		Wire clamp failed, net slid down wire ?			10	
		350-225	R		Net failed to close; see above			20	
		475-350	L	?	May be useable but see above		1012	15	37
		475-350	R	?	May be useable but see above		1012	10	25
		600-475	L	?	May be useable but see above		1015	<5	<10
		600-475	R	?	May be useable but see above		1015	<5	<10
							1015	10	25
	1324-1345	25-0	L	good			1015	10.5	26
		25-0	R	good			1410	10	18
		50-25	L	poor	Net torn		1410	22.5	40
		50-25	R	good			1012		
		75-50	L	poor	Net torn		1012	10	25
		75-50	R	good					
		100-75	L	-	Net apparently worked but no sample	100-0	3882	7.5	5
		100-75	R	useful	Net came up open				
							1015	12.5	31
Sept. 24	0046-0111	225-100	L	good			1015	10	25
		225-100	R	good			1012	5	12
		350-225	L	good			1012	2.5	6
		350-225	R	good			1010	5	12
		475-350	L	good			1010		
		475-350	R	-	Net apparently worked but no sample				
	0413-0500	75-50	L	useful	Did not close	75-0	2773	30	27
		75-50	R	useful	Did not close	75-0	2773	25	23
		100-75	L	good	Nets apparently worked but note		1010	10	25
		100-75	R	good	large difference in L/R volumes		1010	<5	<10
							3420	50	37
	0507-0540	25-0	L	useful	Net did not close		3420	60	44
		25-0	R	useful	Net did not close		1015	10	25
		50-25	L	good			1015	12.5	31
		50-25	R	good					
							1015	20	50
	1546-1630	25-0	L	good			1015	32.5	81
		25-0	R	good			1400	<5	<10
		50-25	L	good			1400	5	9
		50-25	R	good			1012	2.5	6
		75-50	L	good	Nets apparently worked but note		1012	10	25
		75-50	R	good	large difference in L/R volumes		1010	15	37
		100-75	L	good	Nets apparently worked but note		1010	2.5	6
		100-75	R	good	large difference in L/R volumes				
							1015	12.5	31
Sept. 26	1337-1353	25-0	L	good			1015	5	12
		25-0	R	good			1010	10	25
		50-25	L	good			1010	17.5	44
		50-25	R	good					
							1010	17.5	44
	1444-1506	75-50	L	good			1010	15	38
		75-50	R	good			1015	5	12
		100-75	L	good			1015	10	25
		100-75	R	good					
							1015	-	-
	1553-1620	350-100	L	-	Net lost		1015	10	25
		350-100	R	good			1010	10	25
		600-350	L	good			1010	10	25
		600-350	R	good					
							1010	10	25
	1656-1721	25-0	L	good			1010	22.5	56
		25-0	R	good			1015	15	37
		50-25	L	good			1015	20	50
		50-25	R	good					

Date 1968	Local Time	Expected Depth (m) Fished	Side of Net R or L	Overall Reliability	Comments	Depth (m) if differs from exp'd.	Linear Meters Fished	Disp. Vol. (ml)	Biomass ₃ ml/1000m
	1758-1821	75-50	L	poor	Net did not close				
		75-50	R	poor	Net did not close	75-0		30	
		100-75	L	poor	Net did not close	75-0		10	
		100-75	R	poor	Net did not close	100-0		2.5	
						100-0		5	
	1859-1922	75-50	L	good			1015	15	37
		75-50	R	good			1015	15	37
		100-75	L	good			1010	12.5	31
		100-75	R	good			1010	10	25
	2010-2033	350-100	L	good					
		350-100	R	good			1015	10	25
		600-350	L	good			1015	10	25
		600-350	R	good			1010	5	12
							1010	7.5	19
	2118-2137	25-0	L	good					
		25-0	R	good			1010	45	113
		50-25	L	good			1010	35	81
		50-25	R	good			1015	15	37
							1015	17.5	44
	2206-2230	75-50	L	good					
		75-50	R	good			1015	15	37
		100-75	L	good	Nets apparently worked but note large difference in L/R volumes		1015	22.5	56
		100-75	R	good			1010	7.5	19
							1010	55	138
	2328-2351	350-100	L	good	Nets apparently worked but note large difference in L/R volumes				
		350-100	R	good			1010	2.5	6
		600-350	L	good			1010	20	50
		600-350	R	good			1015	2.5	6
							1015	<5	<10
Sept. 27	0040-0103	25-0	L	good					
		25-0	R	good			1015	15	37
		50-25	L	good	Nets apparently worked but note large difference in L/R volumes		1015	25	62
		50-25	R	good			1010	<5	<10
							1010	15	37
	0126-0156	75-50	L	good					
		75-50	R	good			1010	22.5	
		100-75	L	poor	Hole in cod end		1010	20	50
		100-75	R	good			1015	2.5	6
							1015	20	50
	0237-0255	350-100	L	good					
		350-100	R	good			1015	7.5	19
		600-350	L	good			1015	7.5	19
		600-350	R	good			1010	10	25
							1010	5	13
	0341-0404	25-0	L	good					
		25-0	R	good			1010	30	75
		50-25	L	good			1010	22.5	56
		50-25	R	good			1015	12.5	31
							1015	15	37
	0449-0500	75-50	L	useful	Net lost				
		75-50	R	useful			1015	10	25
		100-75	L	useful			1010		
		100-75	R	useful			1010	22.5	56
							1010	20	50
	0606-0633	350-100	L	good					
		350-100	R	good			1010	10	25
		600-350	L	good			1010	10	25
		600-350	R	good			1015	10	25
							1015	5	12

Date 1968	Local Time	Expected Depth (m) Fished	Side of Net R or L	Overall Reliability	Comments	Depth (m) if differs from exp'd.	Linear Meters Fished	Disp. Vol. (ml)	Biomass ₃ ml/1000m ³
0804-0838		75-50	L	poor	Hole in cod end		1015	15	38
		75-50	R	good			1015	10	25
		100-75	L	?			1010	<5	<10
		100-75	R	good			1010	25	62
0929-0953		350-100	L	good			1010	7.5	19
		350-100	R	good			1010	5	13
		600-350	L	good			1050	7.5	18
		600-350	R	good			1050	5	12
1044-1100		25-0	L	good	Nets apparently worked but note large difference in L/R volumes		985	7.5	19
		25-0	R	good			985	5	13
		50-25	L	good			1010	<5	<10
		50-25	R	good			1010	15	38
1133-1200		75-50	L	useful	Net did not close	75-0	1010	37.5	94
		75-50	R	useful	Net did not close	75-0	1010	30	75
		100-75	L	good	Net hung up on wire; note difference		1015	10	25
		100-75	R	useful	in L/R volumes		1015	<5	<10
1228-1250		75-50	L	good			1010	17.5	41
		75-50	R	good			1010	20	50
1332-1350		350-100	L	good			1015	2.5	6
		350-100	R	good			1015	5	12
		600-350	L	good			1010	<5	<10
		600-350	R	good			1010	10	25

Table 4

CLIMAX I WATER TRANSPARENCY

Date 1968	Local Time	Solar Elevation degrees	Wind dir(°) speed(kts)	Clouds ^{a)} type amount	Sea	Swell dir(°) hght(ft)	Secchi Disc Depth (m)	K ^{b)} (m ⁻¹)
Sept. 19	1125-1225	64	120 14	8,4,2 2	Moderate	090 4	32	.060
Depth (m) E₀^{c)} E_{dw}^{d)}/E₀^{e)} (%)								
0								
3.4								
7.9								
10.1								
13.4								
20.1								
21.3								
26.2								
32.0								
41.5								
50.3								
58.8								
74.7								
93.7								
113.4								
Sept. 21	1215-1240	65	170 10	8 4	Slight	120 4	34	.047
Depth (m) E₀ E_{dw}/E₀ (%)								
0								
9.5								
19.5								
40.0								
64.8								
49.1								
29.6								
18.9								
9.2								
Sept. 28	1210-1300	54	050 4	8 2	Slight	020 3	-	.048
Depth (m) E₀ E_{dw}/E₀ (%)								
0								
5.0								
10.4								
15.5								
20.8								
29.9								
39.6								
50.0								
60.8								
68.3								
50.0								
29.6								
10.7								

a) Cloud data are coded using the National Oceanographic Data Center (NODC) method.

b) K is defined as $I = I_0 e^{-Kz}$

c) E₀ = Incident light radiation in foot-candles (cosine collector on ship, above surface)

d) E_{dw} = Downwelling radiation at depth Z (cosine collector)

e) E_{dw}/E₀ at Z=0 gives loss of light at the air-sea interface

Table 5

CLIMAX I PRIMARY PRODUCTIVITY

Date 1968	Depth meters	Chlorophyll-a mg/m ³	Phaeophytin mg/m ³	Productivity mg C/m ³ /hr.	Productivity per unit Chloro-a mg C/hr./mg-Chl-a	Integrated Water Column Productivity mg C/m ² /12 hr. day
Sept. 19	0	.0359	.0347	.161	4.48	156.0
	23	.0442	.0269	.209	4.72	
	33	.0414	.0257	.261	6.30	
	49	.0529	.0290	.223	4.21	
	60	.0602	.0303	.116	1.92	
	74	.0694	.0387	-.002	-.028	
Sept. 20	0	.0366	.0238	.239	6.53	189.6
	23	.0320	.0159	.274	8.56	
	33	.0285	.0195	.261	9.15	
	49	.0489	.0244	.346	7.07	
	60	.0605	.0314	.185	3.05	
	74	.0695	.0442	.043	.618	
Sept. 21	0	.0205	.0156	.211	10.29	240.0
	23	.0330	.0146	.323	9.78	
	33	.0302	.0128	.393	11.35	
	49	.0400	.0165	.326	8.15	
	60	.0495	.0195	.221	4.26	
	74	.0585	.0262	.068	1.16	
Sept. 22	0	.0285	.0116	.212	7.43	180.0
	23	.0327	.0113	.268	8.19	
	33	.0338	.0102	.337	9.97	
	49	.0392	.0116	.329	8.39	
	60	.0534	.0200	.111	2.07	
	74	.0467	.0155	.010	.214	
Sept. 23	0	.0224	.0151	.215	9.59	180.0
	23	.0179	.0160	.260	14.52	
	33	.0249	.0132	.249	10.00	
	49	.0311	.0138	.254	8.16	
	60	.0383	.0214	.127	3.31	
	74	.0604	.0443	.014	.23	
Sept. 24	6	.0320	.0159	.322	10.06	228.0
	26	-	-	.268	-	
	38	.0377	.0146	.202	5.35	
	57	.0572	.0277	.209	3.65	
	70	.0692	.0287	.152	2.19	
	88	.0712	.0408	.030	.42	
Sept. 25	5	.0329	.0209	.241	7.32	192.0
	25	.0240	.0229	.225	9.37	
	36	.0169	.0124	.321	18.99	
	54	.0356	.0339	.198	5.56	
	66	.0196	.0156	.138	7.04	
	82	.0570	.0622	.006	.105	

Table 6

CLIMAX I CHLOROPHYLL-A AND PHAEOPHYTIN

Date 1968	Local Time	Depth meters	Chlorophyll-a μg/L	Phaeophytin μg/L	Date 1968	Local Time	Depth meters	Chlorophyll-a μg/L	Phaeophytin μg/L
Sept. 20	1030	0	.0161	.0134	Sept. 25	2000	20	.0247	.0106
		10	.0130	.0121			40	.0262	.0129
		20	.0074	.0078			60	.0395	.0153
		35	.0096	.0091			80	.0480	.0499
		50	.0021	.0239			90	.0542	.0981
		75	.0192	.0031			100	.1424	.1791
		100	.0053	.0334			110	.1376	.1945
		125	.0064	.0448			120	.0984	.1430
		150	.0010	.0271			160	.0154	.0244
		200	.0011	.0144			180	.0087	.0208
Sept. 21	1000	0	.0264	.0113	Sept. 27	1900	200	.0089	.0194
		10	.0181	.0272			230	.0069	.0127
		25	.0228	.0104			260	.0024	.0084
		35	.0325	.0230			290	.0019	.0050
		50	.0420	.0205			320	.0014	.0055
		60	.0503	.0317			350	.0028	.0027
		75	.0676	.0319					
		100	.1214	.1605			300	.0000	.0125
		125	.1206	.1742			400	.0014	.0050
		150	.0370	.0648			500	.0005	.0067
Sept. 22	-	200	.0070	.0152			600	.0010	.0048
							800	.0004	.0044
Sept. 22	-	150	.0436	.0729	Sept. 28	0100	1000	.0005	.0049
		200	.0021	.0128			0	.0286	.0477
Sept. 22	2400	0	.0500	.0239			10	.0311	.0133
		10	.0194	.0174			25	.0279	.0154
		25	.0215	.0221			35	.0267	.0257
		35	.0549	.0719			50	.0405	.0229
		75	.0559	.0391			60	.0490	.0246
		100	.1396	.2653			75	.0614	.0328
		110	.1757	.2175			100	.1631	.2137
		125	.1057	.2061			125	.1439	.2192
		150	.0237	.0502			150	.0626	.1059
		200	.0107	.0217			200	.0103	.0218
Sept. 24	0900	0	.0538	.0269					
		10	.0351	.0252					
		25	.0361	.0210					
		35	.0323	.0174					
		50	.0504	.0272					
		60	.0699	.0365					
		75	.0771	.0586					
		100	.1794	.2089					
		125	.0449	.0765					
		150	.0224	.0425					
		200	.0024	.0112					

Table 7

CLIMAX I CHLOROPHYLL PROFILE DATA

Date 1968	Local Time	Depth of Maximum Layer (m)	Estimated Depth Lag: Pump to Fluorometer (m)	Maximum Value (0-100 scale) Door 10	Surface Value (0-100 scale) Door 10	Character of Maximum Value	Depth Range of Sample	Comments
Sept. 18	2134	84	54	34.0	19.3	Single max. value	0-140	Poor depth Correlation
Sept. 19	1719	84	54	26.5	11.3	Possibly double	0-110	Short scale
Sept. 20	2136	-	-	-	8.8	-	Surface	Break in hose
Sept. 20	2312	90	54	30.1	8.0	Single	0-230	Short scale
Sept. 21	2123	-	-	-	10.5	-	Surface	Poor record
Sept. 21	2335	90	54	36.8	15.6	Single	0-230	Short scale
Sept. 22	2351	101	54	18.9	13.0	Single	0-230	Salinity sam- ples enabled good depth correlation
Sept. 23	2025	108	54	24.1	10.0	Single	0-156	
Sept. 24	0900	109	54	38.1	23.5	Single	0-176	
Sept. 25	0320	-	-	-	6.6	-	Surface	Poor record
Sept. 25	0800	101	54	35.2	15.5	Single	0-240	Severe hose angle
Sept. 26	0830	91	54	38.3	10.5	Single	0-225	Salinity sam- ples taken at 128-225m
Sept. 27	2400	99-105	54	34.0	8.0	Single	0-200	

Table 8

CLIMAX I BIRD AND FISH SIGHTINGS

Date 1968	Local Time	Description	Number
Sept. 19	1200	Red-tailed Tropic bird	1
	1830	White-tip shark, 5ft	1
Sept. 20	1200	White-tailed Tropic bird	1
	1340	White-tailed Tropic bird	1
	1345	White-tailed Tropic bird	1
	all day	Flights of 5-20 small birds heading South about 7-15ft above water; possibly Golden Plover	
Sept. 21	1130	White-tip shark, 6ft	1
	1310	Dolphinfish	5
	1315	Black-footed Albatross	1
	1345	Golden Plover	1
	1510	Skipjack	40-50
	1700	Dolphinfish	1
	1800	White-tailed Tropic bird	1
Sept. 22	0658	Golden Plover	1
	0830	Frigate birds	2
	0845	Sooty Shearwater	2
	0915	Dolphinfish	5
	1345	Golden Plover	2
	1710	Golden Plover	1
Sept. 23	0630	Golden Plover	2
	0715	Golden Plover	10
	0730	Golden Plover	12
	0915	White-tailed Tropic bird	1
	0930	Sooty Shearwater	2
	1008	Golden Plover	3
	1330	White-tailed Tropic bird	1
	1330	Golden Plover	4
	1425	Sooty Shearwater	1
	1500	Frigate bird	1
	1630	Red-footed Booby	1
Sept. 24	1210	Golden Plover	3
	1215	Golden Plover	9
	1430	Pelagic Triggerfish	5
	1600	White-tailed Tropic bird	1
Sept. 26	0610	Golden Plover	1
	1345	Red-tailed Tropic bird and White-tailed Tropic bird flying together	1 each
	1820	Golden Plover	2
Sept. 27	0200	Shark (White-tip?)	1
	0650	Golden Plover	1
	0730	Wedge-tailed Shearwater	2
	0940	Golden Plover	1

Table 9

CLIMAX I FISH CATCHES

Date 1968	Local Time	Species	Weight (lbs.)	Length (ins.)
Sept. 20	1500	Dolphinfish		
	2330	"	12	36
Sept. 21	1400	"	15	37
	1400	"	11	32
	1730	"	15	36
Sept. 22	0915	"	8	30
Sept. 23	0200	"	4	24
Sept. 24	1800	"	10	32
	2230	" , (four)	8	30
Sept. 25	1000	"	8	30
	1200	"	8	30
Sept. 26	0630	"	8	30
Sept. 28	1700	"	?	?

DISTRIBUTION LIST

Inter-American Tropical Tuna Commission
(c/o Scripps Institution of Oceanography)

Dr. James Joseph

National Marine Fisheries Service
(c/o Scripps Institution of Oceanography)

Dr. Alvariño de Leira
Mr. Ronald Lynn
Mr. Robert W. Owen, Jr.
Director's Office
Library (2)

Scripps Institution of Oceanography

Mr. Norman E. Anderson
Dr. Maurice Blackburn (2)
Dr. Edward Brinton
Dr. T. J. Chow
Mr. Edward H. Coughran
Dr. Richard W. Eppley
Dr. Abraham Fleminger
Mr. Jeffery D. Frautschy
Mr. John D. Isaacs
Miss Margaret D. Knight
Dr. John A. McGowan
Dr. Fred B. Phleger
Mr. Joseph L. Reid, Jr.
Dr. Richard H. Rosenblatt
Mr. Richard A. Schwartzlose
Mr. George H. Snyder
Dr. Elizabeth Venrick
Director's Office
Library, SIO, Archives
Library, SIO, Circulation (3)

CalCOFI Distribution List

Mr. William Allen, Jr.
1070 - 16th Place South
Edmonds, Wash. 98020

Mr. D. L. Alverson, Base Dir.
Northwest Fisheries Center
National Marine Fisheries
Service
2725 Montlake Blvd.
Seattle, Wash. 98102

Mr. William E. Batzler
Code 3185 C
Naval Undersea Center
San Diego, Calif. 92132

Director
Biological Station
Fisheries Research Board
of Canada
P. O. Box 100
Nanaimo, British Columbia,
Canada

British Museum
Dept. of Printed Books-SB
Stechert-Hafner, Inc.
London, W. C. 1, England

British Navy Staff
British Embassy
3100 Massachusetts Ave. N.W.
Washington, D. C. 20008
Attn: Scientific Info.
Officer

Dr. Dail W. Brown
Ofc. of Oceanography and
Limnology
Smithsonian Institution
Washington, D. C. 20560

Dr. Wayne V. Burt
Prof. of Oceanography
Department of Oceanography
Oregon State University
Corvallis, Ore. 97331

Library
California Academy of
Sciences
Golden Gate Park
San Francisco, Calif. 94118

University of California
Serials Department
General Library
Berkeley, Calif. 94720

Marine Technical Information
Center
Department of Fish & Game
350 Golden Shore
Long Beach, Calif. 90802

Mr. Charles R. Carry
Tuna Research Foundation
215 Cannery Street
Terminal Island, Calif. 90731

Capitan de Navio
Juan B. Torti
Servicio de Hidrografia Naval
Avenida Montes de Oca 2124
Buenos Aires, Argentina

Biblioteca
Instituto Nacional de Pesca
Centro de Promocion de Pesquera
Departamento Oceanografico
Apartado Postal 396,
Mazatlan, Sinaloa, Mexico

Director, Centro de Promocion
Pesquera, Apdo. No. 396
Mazatlan, Sinaloa, Mexico

Mr. Harold B. Clemens, Jr.
California State Fisheries Lab.
350 Golden Shore
Long Beach, Calif. 90802

Dr. Daniel M. Cohen
National Center for Systematics
Ichthyological Laboratory
U. S. National Museum
Washington, D. C. 20560

Dr. G. M. Cresswell
Tiburón Oceanographic Institute
Tiburón, Calif. 94920

Deutsche Akademie der
Wissenschaften zu Berlin
Institut für Meereskunde,
Seestr. 15
Rostock-Warnemünde (DDR-253)
Deutsche Demokratische Republik
(East Germany)

Deutsches Hydrographisches Institut
Bernhard-Nocht-Str. 78
2 Hamburg 4,
Bundesrepublik Deutschland
(West Germany)

Mr. Robert L. Eberhardt
Marine Occupations Program
7250 College Drive
San Diego, Calif. 92111

Environmental Sciences Division
Code 3150, Box 7
Pacific Missile Range
Point Mugu, Calif. 93041

Dr. David Farris
Department of Biology
San Diego State University
San Diego, Calif. 92115

Library
Oceanographic Group
Fisheries Research and Development
Agency
Pusan, Korea

Prof. James A. Gast
Division of Natural Resources
Humboldt State College
Arcata, Calif. 95521

Dr. Robert H. Gibbs, Jr.
Division of Fisheries
U. S. National Museum
Washington, D. C. 20560

Dr. Donn S. Gorsline
Department of Geology
University of Southern
California
Los Angeles, Calif. 90007

Hancock Library of Biology
and Oceanography
Allan Hancock Foundation
University of Southern
California
Los Angeles, Calif. 90007

Dr. William J. Hargis, Jr.,
Director
Virginia Inst. of Marine
Sciences
Gloucester Point, Va. 23062

Mr. Koji Hidaka
Ocean Research Institute
University of Tokyo
Nakano, Tokyo, Japan

Mr. T. Hirano
Tokai Regional Fisheries
Research Laboratory
5, Kachidoki, Chuo-ku
Tokyo, Japan

Director
Institute of Marine Science
University of Alaska
College, Alaska 99701

Librarian
Institute of Marine Science
University of Miami
10 Rickenbacker Causeway
Miami, Fla. 33149

Director, Inst. de
Geofisica
Torre de Ciencias, 3er Piso
Universidad Nacional
Autonoma de Mexico
Villa Obregon, D. F.,
Mexico

Biblioteca
Instituto Nacional de Pesca
Chiapas 121
Mexico 7 D. F.

Director
Instituto del Mar
Apartado Postal 3734
Lima, Peru

Director
Instituto Hidrografico de
la Armada de Chile
Departamento de Oceanografia
Casilla 324
Valparaiso, Chile

Director
Estacion de Biologia Marina
Instituto Tecnologico de
Veracruz
Heroica, Veracruz, Mexico

Intersea Research Corp.
P. O. Box 2389
La Jolla, Calif. 92037

Japan Meteorological Agency
Oceanographical Section
Tokyo, Japan

Japan Oceanographic Data
Center
Hydrographic Division
Maritime Safety Agency
No. 3-1, 5-Chome, Tsukiji,
Chuo-ku
Tokyo, Japan

Editor
Journal of Marine Research
Box 2161, Yale Station
New Haven, Conn. 06520

Mr. Larry Kiml, Director
Water and Natural Resource
Depts.
Calif. State Chamber of
Commerce
455 Capitol Mall, Suite 300
Sacramento, Calif. 95814

Dr. H. Kitamura
Oceanographic Section
Kobe Marine Observatory
Kobe, Japan

Dr. E. C. LaFond
Code 50401
Naval Undersea Center
San Diego, Calif. 92132

Mr. Owen S. Lee
Program Manager for Physical
Oceanography
Code 504
Naval Undersea Center
San Diego, Calif. 92132

Dr. Alfred J. Carsola
Lockheed Marine Laboratory
3380 N. Harbor Drive
San Diego, Calif. 92101

Dr. Yuan-Hui Li
Hoffman Laboratory
Harvard University
20 Oxford Street
Cambridge, Mass. 02138

Mr. Jotaro Masuzawa
Oceanographical Section
Japan Meteorological Agency
Chiyoda-ku, Tokyo, Japan

Dr. Giles W. Mead
Museum of Comparative Zoology
Harvard University
Cambridge, Mass. 02138

Librarian
Ministry of Agriculture, Fisheries
and Food
Fisheries Laboratory
Lowestoft, Suffolk, England

Library
Moss Landing
Marine Laboratories
P. O. Box 223
Moss Landing, Calif. 95039

Director
Nansei Laboratory
6-2, Sanbashi-dori, Kochi-shi
Kochi, Japan

Librarian
Narragansett Marine Laboratory
University of Rhode Island
Kingston, R. I. 02881

Director
U. S. Department of Commerce
Natl. Oceanic and Atmospheric Admin.
National Oceanographic Data Center
Washington, D. C. 20235

Librarian
Natl. Oceanic and Atmospheric Admin.
Washington Science Center
Rockville, Md. 20852

Director (6)
U. S. Department of Commerce
Natl. Oceanic & Atmospheric Admin.
World Data Center A, Oceanography
Washington, D. C. 20235

Dr. Kenneth S. Norris
Director, Coastal Marine Laboratory
University of California
Santa Cruz, Calif. 95064

Dr. Robert M. Norris
Department of Geology
University of California
Santa Barbara, Calif. 93106

Oceanographic Research Institute
Centenary Aquarium Bldgs.
2 West Street
Durban, Natal, South Africa

Library, Research Laboratory
Oregon Fish Commission
Route 2, Box 31A
Clackamas, Ore. 97015

Pacific Marine Fish. Commission
741 State Office Building
1400 S. W. Fifth Avenue
Portland, Ore. 97200

Dr. Robert G. Paquette
Dept. of Oceanography
U. S. Naval Postgraduate
School
Monterey, Calif. 93940

Director
San Diego Society of
Natural History
P. O. Box 1390
San Diego, Calif. 92112

Dr. G. L. Pickard
Institution of Oceanography
University of British
Columbia
Vancouver 8, B.C., Canada

Dr. D. W. Pritchard, Director
Chesapeake Bay Institute
The Johns Hopkins University
Oceanography Building
Baltimore, Md. 21218

Mr. D. W. Privett, Librarian
Inst. of Oceanographic
Science
Wormley, Near Godalming
Surrey, England

Publications Office (2)
101 University Hall
2200 University Avenue
Berkeley, Calif. 94720

Dr. Ricardo M. Pytkowicz
Department of Oceanography
Oregon State University
Corvallis, Ore. 97331

Mr. John Radovich, Head
Operations Research Branch
Department of Fish & Game
1416 Ninth Street
Sacramento, Calif. 95814

Mr. Gunnar I. Roden
Department of Oceanography
University of Washington
Seattle, Wash. 98195

Environmental Prediction
Research Facility
U. S. Naval Postgraduate
School
Monterey, Calif. 93940

Biblioteca
Unidad de Ciencias Marinas
Universidad Autonoma de
Baja Calif.
Apartado de Correos 453
Ensenada, B.C., Mexico

Librarian
Serials Department
San Diego State University
Library
San Diego, Calif. 92115

Librarian
Geology-Oceanography Dept.
San Fernando State University
Northridge, Calif. 91324

Mr. Daitaro Shoji
Oceanographic Section
Japanese Hydrographic Office
5-chome, Tsukiji, Chuo-ku
Tokyo, Japan

Mr. J. C. Simpson, Project
Manager
Fisheries Research and
Development Project
Apartado 2578
Caracas, Venezuela

Mr. Edmund H. Smith, Director
Pacific Marine Station
Dillon Beach, Calif. 94929

Dr. Reimer Simonsen
Institut fur Meeresforschung
285 Bremerhaven
Am Handelshafen 12
Bundesrepublik
Deutschland (West Germany)

Director
Institute of Marine Science
University of Miami
10 Rickenbacker Causeway
Miami, Fla. 33149

Library of the Division of
Systematic Biology
Stanford University
Stanford, Calif. 94305

Prof. Henry M. Stommel
Mass. Institute of Technology
Bldg. 54, Room 1416
Cambridge, Mass. 02139

Dr. Arthur D. Stump
Department of Chemistry
San Jose State University
San Jose, California 95100

Dr. Bruce A. Taft
Department of Oceanography
University of Washington
Seattle, Wash. 98195

Dr. Norman Tebble, Director
The Royal Scottish Museum
Chambers Street
Edinburgh, Scotland

Department of Oceanography
Texas A. and M. University
College Station, Texas 77843

Mr. A. J. Thomson
Official Secretary
New South Wales Government
Offices
56, Strand
London, W. C. 2, England

Dr. R. B. Tibby
Hancock Foundation
University of Southern Calif.
University Park
Los Angeles, Calif. 90007

Librarian
Nat'l. Oceanic and Atmospheric Admin.
Southeast Fisheries Center, NMFS
75 Virginia Beach Drive
Miami, Fla. 33149

Dr. M. Uda
College of Marine Science
and Technology
Tokai University
1000 Orido, Shimizu-shi
Shizuoka-ken, Japan

Commanding Officer
USCG Oceanographic Unit
Bldg. 159-E, Navy Yard Annex
Washington, D. C. 20390

Los Angeles Field Office
U. S. Coast and Geodetic Survey
417 S. Hill St., Room 535
Los Angeles, Calif. 90013

Librarian
Southwest Fisheries Center
P. O. Box 3830
Honolulu, Hawaii 96812

Chief, Branch of Marine Fisheries
U. S. Department of Commerce
Nat'l. Oceanic and Atmospheric Admin.
Washington, D. C. 20240

Tiburon Fisheries Laboratory
NMFS, NOAA
P. O. Box 98
Tiburon, Calif. 94920

Librarian
U. S. Naval Civil Engineering Lab.
Port Hueneme, Calif. 93041

Pacific Support Group
U. S. Naval Oceanographic Office
ASW School, Bldg. 6
San Diego, Calif. 92147

Commander (2)
U. S. Naval Oceanographic Office
Library Code A 43
Washington, D. C. 20390

Library (2)
U. S. Naval Electronics
Laboratory Center
San Diego, Calif. 92152

University of Washington (2)
Fisheries-Oceanography
Library
203 Fisheries Center
Seattle, Washington 98105

Dr. C. S. Wong, Head
Ocean Chemistry Division,
Pac. Region, Mar. Sci.
Branch, Dept. of the
Environment
1230 Government Street
Victoria, B. C., Canada
V8W 1Y4

Woods Hole Oceanographic
Inst.
Document Library LO-206
Woods Hole, Mass. 02543

Mr. Hajime Yamanaka
High Seas Fisheries
Research Lab.
Orido 1000, Shimizu
Shizuoka-ken, Japan

Dr. Kozo Yoshida
Geophysical Inst.
University of Tokyo
Hongo, Tokyo, Japan