

AD-A095 524

NEW YORK UNIV N Y DEPT OF CHEMISTRY

F/G 7/4

NON-ADIABATIC CAPTURE OF MOBILE ELECTRONS IN AQUEOUS SOLUTION.(U)

FEB 81 K VON BURG, P DELAHAY

N00014-75-C-0397

UNCLASSIFIED

TR-14

NL

1 OF 1

NO A
00000004

END

DATE
FILMED

9-81

DTIC

AD A095524

LEVEL

OFFICE OF NAVAL RESEARCH

Contract ⁵ N00014-75-C-0397

Task No. NR 051-258

⁹ TECHNICAL REPORT NO. 14 (New Series)

⁶ Non-adiabatic Capture of Mobile Electrons in Aqueous Solution •

by

¹⁰ Kathrin von Burg and Paul Delahay

Prepared for publication

in

Chemical Physics Letters

New York University
Department of Chemistry
New York, NY

¹¹ February 1981

RECEIVED
FEB 26 1981
C

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

406817

81 2 26 081

DEC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 (New Series)	2. GOVT ACCESSION NO. AD-A095	3. RECIPIENT'S CATALOG NUMBER 534
4. TITLE (and Subtitle) NON-ADIABATIC CAPTURE OF MOBILE ELECTRONS IN AQUEOUS SOLUTION		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0397
9. PERFORMING ORGANIZATION NAME AND ADDRESS New York University New York, NY 10003		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 051-258
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		12. REPORT DATE February 1981
		13. NUMBER OF PAGES 21
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted for publication in Chemical Physics Letters		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Aqueous solution Inorganic ions Mobile electrons Non-adiabatic capture Photoelectron spectroscopy		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Experimental evidence and theoretical arguments are presented for 11 effective and eight ineffective cations for non-adiabatic capture of mobile electrons in the photoionization (≤ 10.5 eV) of anions (Cl^- , ClO_4^- , SO_4^{2-}) in aqueous solution. This effect lowers yields and increases threshold energies for photoelectron emission by anions into the gas phase.		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

NON-ADIABATIC CAPTURE OF MOBILE ELECTRONS IN AQUEOUS SOLUTION

KATHRIN von BURG and PAUL DELAHAY

Department of Chemistry, New York University, 4 Washington Place, Room 514,
New York, NY 10003, U.S.A.

Experimental evidence and theoretical arguments are presented for 11 effective and eight ineffective cations for non-adiabatic capture of mobile electrons in the photoionization (< 10.5 eV) of anions (Cl^- , ClO_4^- , SO_4^{2-}) in aqueous solution. This effect lowers yields and increases threshold energies for photoelectron emission by anions into the gas phase.

1. Introduction

The yield for photoelectron emission by aqueous solutions of inorganic anions (Cl^- , SO_4^{2-}) was recently observed [1] to be lower in the presence of certain cations (e.g., H^+ , Fe^{2+}) than for the corresponding salts of the alkali metals. The effect is displayed in fig. 1. Curve A (2 M KCl) shows the emission by Cl^- (threshold energy $E_t = 8.77$ eV [2]), and B (1 M FeCl_2) displays the emission by Fe^{2+} ($E_t = 7.35$ eV [1]) and, at higher photon energies ($E \gtrsim 9$ eV), by Cl^- . Since the Cl^- concentration is the same for A and B, one would expect higher yields for B than A over the full range of photon energies. The opposite is observed for $E \gtrsim 9.0$ eV, that is, in the range in which emission by Cl^- becomes significant. Moreover, it was found [1] that the threshold energy E_t of Cl^- for the FeCl_2 solution is higher (~ 9.1 eV) than E_t of Cl^- for the KCl solution (8.77 eV [2]). The effect of cations on emission by anions was only referred to incidentally in [1] but was investigated in detail later. The results of this new work are discussed here.

2. Experimental results

Emission spectra displaying the yield Y against photon energy E (< 10.5 eV) were obtained by the methods reported in [3] and applied in


[1,2]. Two cases were distinguished in the analysis of data:

(i) The cation (e.g., H^+ , K^+) does not emit electrons for $E \leq 10.5$ eV. Plots of Y^n vs. E [4] were prepared with $n = 0.5$ [2] for the anions (Cl^- , ClO_4^- , SO_4^{2-}) studied here. Linearity of these plots was apparently not affected by the effect of cations on emission by anions. The threshold energy E_t of the anion was determined by extrapolation to $Y = 0$.

(ii) Both the cations C and anions A emit ($E_t(C) < E_t(A)$). Plots of Y^n vs. E were prepared in the range ($E < E_t(A)$) in which only cations emit ($Y \equiv Y(C)$), and $E_t(C)$ was obtained by extrapolation to $Y = 0$. Selection of n for cations is discussed in [1]. The resulting linear plots were extended to $E > E_t(A)$, and $Y(A) = Y - Y(C)$ was computed. The threshold energy $E_t(A)$ was obtained by extrapolation to $Y(A) = 0$ from the plot of $[Y(A)]^n$ vs. E ($n = 0.5$).

The following cases will be considered: (i) Only the anion emits electrons, and the cation either affects (H^+ , Ba^{2+}) emission by the anion or does not (e.g., K^+). (ii) Both the anion and cation (e.g., Fe^{2+}) display photoelectron emission, and the cation affects emission by the anion.

Results at constant anion concentration will be discussed first for $HCl + KCl$ and $H_2SO_4 + Li_2SO_4$ mixtures (no emission by cations). Since Y at a given E is proportional to the anion concentration [1-4], the squared slope of the $Y^{0.5}$ vs. E plot, $(dY^{0.5}/dE)^2$, should be proportional to the anion concentration. The latter is constant in the present experiments, and therefore the squared slope should be constant in the absence of a cation effect. Actually, the squared slope decreased linearly with increasing H^+ concentration, x , and the threshold energies of Cl^- and SO_4^{2-} shift to higher values (fig. 2).



Similar experiments were done with $\text{MCl}_2 + \text{KCl}$ mixtures at constant Cl^- concentration (2 M) for Mg^{2+} , Ca^{2+} , Sr^{2+} , Ba^{2+} (no emission by cations). Except for Ba^{2+} , no evidence was found for a cation effect on emission by Cl^- . The squared slope and threshold energy ($E_t = 8.77 \pm 0.05$ eV) were independent of M^{2+} ion concentration. Results for BaCl_2 were similar to those of fig. 2: ratio of squared slopes equal to 0.50 for 1 M BaCl_2 /2 M KCl , $\Delta E_t(1 \text{ M BaCl}_2) \approx 0.25$ eV. Thus, Ba^{2+} ions affect the emission by Cl^- . A similar conclusion was reached for emission by ClO_4^- (fig. 3) over a wider concentration range than allowed by BaCl_2 solubility.

The shift of threshold energy ($E_t = 8.77$ and 8.65 eV for Cl^- and SO_4^{2-} , respectively [2]) resulting from the cation effect was determined for several cations (1 M solutions) studied in [1]: $\Delta E_t(\text{Cl}^-) \approx 0.40$ eV for Cr^{2+} , Mn^{2+} , Co^{2+} , Ni^{2+} , V^{3+} , Cr^{3+} ; $\Delta E_t(\text{SO}_4^{2-}) \approx 0.34$ eV for V^{2+} , Cr^{2+} , Ni^{2+} , Cu^{2+} . The standard deviation for these values of ΔE_t varied from 0.05 to 0.15 eV. A more detailed study was made for $\text{FeCl}_2 + \text{KCl}$ mixtures at constant Cl^- concentration (2 M). Results were similar to those of fig. 2: ratio of squared slopes equal to 0.45 for 1M FeCl_2 /2 M KCl , $\Delta E_t(\text{Cl}^-) = 0.28$ eV for 1 M FeCl_2 .

The dependence of $Y(\text{Cl}^-)$ on concentration was also investigated with pure salts and pure HCl . The squared slope $(d[Y(\text{Cl}^-)]^{0.5}/dE)^2$ should be proportional to the salt (or HCl) concentration in the absence of cation effect [1-4]. Proportionality was verified for the alkali cations and the NH_4^+ ions (no cation effect) but was not observed for HCl and FeCl_2 . Curves with a maximum were observed in these two cases (fig. 4). Emission by Fe^{2+} was not affected in any way similar to emission by anions. The quantity $(d[Y(\text{Fe}^{2+})]^{0.4}/dE)^{2.5}$ ($n = 0.4$ for Fe^{2+} [1]) increased linearly with FeCl_2 concentration (fig. 4), and E_t (7.35 eV) was independent of this concentration. Similar observations to those for Fe^{2+} were made for the

cations studied in [1]. There was also no significant change of the threshold energies of Ag^+ , Tl^+ , Sn^{2+} , Pb^{2+} , Fe^{3+} when solutions of these cations were made highly acidic (up to 6 M).

3. Non-adiabatic capture of mobile electrons by cations

Two different approaches can be followed to interpret the results of sec. 2. (i) One supposes that the photoionization rate of anions is decreased by cations, e.g., because of complexation of anions or attenuation of the photon flux in solution. Such an interpretation could not be reconciled with experiment and was ruled out. (ii) One assumes that photoionization of anions is not affected by cations, but that some of the electrons generated by this process are captured by cations. This second interpretation is adopted here.

Electrons upon emission from solution into the gas phase exhibit a distribution over the full range of kinetic energies which is consistent with energy conservation for the prevailing threshold and photon energies [5,6]. A major decrease in yield would not be expected under these conditions if capture was restricted to thermalized and/or hydrated electrons. Yet major lowering in yield is observed, e.g., the ratio of squared slopes for HCl/LiCl is equal to 0.14 for 6 M solutions. We therefore conclude that lowering of the emission yield cannot be attributed to the loss of electrons which were thermalized in solution prior to capture by cations.

Capture by different cations M^{2+} will be examined by comparing the standard reduction potential E^0 for M^{2+}/M^+ or M^+/M with the energy level of mobile electrons. The E^0 's (in V) are [7]: ~ -3.5 (Na^+ , K^+ , Cs^+), -4.0 to -5.5 (Mg^{2+} , Ca^{2+} , Sr^{2+} , Ba^{2+}), -2.7 to -3.1 (Mn^{2+} , Fe^{2+} , Co^{2+} , Ni^{2+}). The bottom of the conduction band of mobile electrons in water is ~ 0.25 V more negative [8,9] in the hydrogen electrode scale than the standard potential

of the hydrated electron (-2.77 eV). Hence, one has $E^m \leq -3.0$ V for mobile electrons. On the basis of the E^0 and E^m values, one concludes that capture should be possible for the transition metal cations M^{2+} but impossible for Ba^{2+} . Yet Ba^{2+} ($E^0 \approx -4.6$ to -5.1 V) ions exhibit a cation effect whereas the alkali metal cations having lower $|E^0|$ values (~ -3.5 V) than Ba^{2+} do not. Thus, the foregoing analysis leads to contradictions.

The difficulties with the previous interpretation are removed if capture is considered in a time interval which is so short that only electronic polarization of the medium about the cation changes upon capture of a mobile electron. The species generated in such a non-adiabatic process (in the spectroscopic sense) should ultimately be reoxidized (e.g., by H^+) after relaxation of the orientation polarization. The difference $\Delta G_S^2 - \Delta G_S^1$ between the solvation energies of M^{2+} and M^+ appearing in the expression of E^0 for M^{2+}/M^+ [1,7] is smaller in the non-adiabatic case than for the adiabatic process. The difference $\Delta G_S^2 - \Delta G_S^1$ can be calculated, at least approximately, by means of the Born equation and by using the optical dielectric constant of water. One obtains (ionic radii from [10], other data from [11]) the potential (in V) $E^* = -3.9$ (Mg^{2+}), -2.2 (Ca^{2+}), -1.8 (Sr^{2+}), -1.4 (Ba^{2+}), -1.0 (V^{2+}), -0.8 (Fe^{2+}), -0.7 (Mn^{2+}), and less negative values for Cr^{2+} , Co^{2+} , Ni^{2+} , Cr^{3+} , H^+ . One also has $E^* \leq -2.5$ V for the alkali metal cations. These values of E^* are in general much less negative than the corresponding E^0 's for an adiabatic process.

The sequence of E^* values agrees with the results in sec. 2: cation effect for $E^* \geq -1.4$ V (Ba^{2+}) and no effect for $E^* \leq -1.8$ V (Sr^{2+}). These two limits for E^* cannot be compared directly with the value $E^m \leq -3.0$ V for mobile electrons. Thus, the orientation polarization of the medium about the anion emitting an electron does not change in the time interval

before capture of this electron by a cation. The electron therefore is in the potential well resulting from the orientation polarization about the anion. The potential energy is [12] $-(e^2/r)(\epsilon_0^{-1} - \epsilon_s^{-1})$, where e is the electronic charge, r the distance from the center of the well, ϵ_0 and ϵ_s the optical and static dielectric constants of water, respectively. If one assumes, for instance, a distance of 0.5 nm between anion and cation in solution, the potential energy is ~ -1.6 eV. Hence, the potential to be compared with the E^* values is not $E^m \sim -3.0$ V but rather $E \sim -1.4$ V. The limit of -1.4 V corresponds to $E^* = -1.4$ V for Ba^{2+} . A quantum [12] rather than classical calculation should be performed, and the uncertainty on energy on the scale of time being considered may not be negligible. Anyhow, our analysis does show that the process of non-adiabatic capture being proposed seems justified.

The foregoing analysis does not account for the absence of cation effect on the emission by cations (e.g., Fe^{2+} , sec. 2). Low-energy mobile electrons have a thermalization length (2 to 4 nm [9,13]) which exceeds the average distance between ions of the same sign (≈ 1.2 nm for 1 M). Capture of mobile electrons by cations prior to thermalization therefore seems possible in the photoionization of either cations or anions. The same conclusion is reached from consideration of the de Broglie wavelength of electrons [14] rather than their thermalization length. Actually, the threshold energies of cations are independent of their concentration (e.g., Fe^{2+}). Thus, the capture of mobile electrons must somehow be restricted to shorter distances than the thermalization length. This would be the case if capture occurs via tunneling [14,15]. Moreover, the average ion configurations are different for emission by anions and cations: emitting anions are surrounded by electron-capturing cations whereas emitting cations are shielded by anions from the nearest cations.

4. Kinetics of electron capture

The results of fig. 2 to 4 can be interpreted in a simple way by assuming that the generation of electrons and their capture proceed at rates which are proportional, respectively, to the anion (C_a) and cation (C_c) concentrations. If C_a is constant, the production rate is constant but the capture rate and consequently the emission yield decrease with increasing concentration C_c (fig. 2 and 3). If $C_a = C_c$ (HCl) or $C_a = 2C_c$ ($FeCl_2$), a maximum in the curve of the squared slope vs. C_c is observed (fig. 4) because capture overtakes production. The ratio of the squared slope to concentration for the data on HCl and $FeCl_2$ of fig. 4 decreases linearly with concentration up to ~ 1.5 M Cl^- concentration and more slowly at higher concentrations. A similar deviation from simple homogeneous kinetics is observed for the capture of mobile electrons in radiolysis studies of liquids [14]. The deviation can be interpreted as indicative of tunneling [14].

Acknowledgment

This work was supported by the Office of Naval Research and the National Science Foundation.

References

- [1] P. Delahay, K. von Burg and A. Dziedzic, Chem. Phys. Lett., accepted.
- [2] K. von Burg and P. Delahay, Chem. Phys. Lett., accepted.
- [3] I. Watanabe, J. B. Flanagan and P. Delahay, J. Chem. Phys. 73 (1980) 2057.
- [4] A. M. Brodsky and A. V. Tsarevsky, J. Chem. Soc. Faraday Trans. II 72 (1976) 1781.
- [5] L. Nemec, L. Chia and P. Delahay, J. Phys. Chem. 79 (1975) 2935.
- [6] L. Nemec, H. J. Gaehrs, L. Chia and P. Delahay, J. Chem. Phys. 66 (1977) 4450.

- [7] J. H. Baxendale and R. S. Dixon, Z. phys. Chem. N.F. 43 (1964) 161.
- [8] A. Henglein, Ber. Bunsenges. Phys. Chem. 78 (1974) 1078.
- [9] Yu. Ya. Gurevich, Yu. V. Pleskov and Z. A. Rotenberg, Photoelectrochemistry, translation by H. S. Wroblowa (Consultants Bureau, New York, 1980, pp. 91, 107-110).
- [10] R. M. Noyes, J. Am. Chem. Soc. 84 (1962) 513; 86 (1964) 971.
- [11] Selected Values of Chemical Thermodynamic Properties, Circular No. 500, National Bureau of Standards (U.S. Government Printing Office, Washington, D.C., 1952).
- [12] R. Platzman and J. Franck, Z. Phys. 138 (1954) 411.
- [13] H. Neff, J. K. Sass and H. J. Lawerenz, J. Phys. Chem. 84 (1980) 1135.
- [14] H. B. Steen, in Electron-solvent and anion-solvent interactions, L. Kevan and B. C. Webster, editors (Elsevier, Amsterdam, 1976, pp. 180-190).
- [15] A. Mozumder in ref [14], pp. 163-170.

Captions to Figures

Fig. 1. Photoelectron emission spectra of 2 M KCl (A) and 1 M FeCl_2 (B).

Fig. 2. Squared slope (top) vs. concentration x for x M HCl + $(2 - x)$ M KCl (A) and $x/2$ M H_2SO_4 + $(1 - x/2)$ M Li_2SO_4 (B). Shift ΔE_t of threshold energy (bottom) vs. x for emission by Cl^- (A) and SO_4^{2-} (B). $E_t = 8.77$ and 8.65 eV, respectively, for 2 M KCl and 1 M Li_2SO_4 .

Fig. 3. Squared slope (top) vs. concentration x for x M $\text{Ba}(\text{ClO}_4)_2$ + $(6 - 2x)$ M NaClO_4 and shift ΔE_t for ClO_4^- vs. x (bottom). $E_t = 8.45$ for ClO_4^- .

Fig. 4. Squared slope for emission by Cl^- ions vs. concentration of HCl (A) and FeCl_2 (B). Slope to the power 2.5 for emission by Fe^{2+} ions vs. concentration of FeCl_2 (C).




FIG. 1




FIG. 2




Fig. 3




FIG.4

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 44106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. M. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co., Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. F. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6G 2G2	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Rubv DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton SO9 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1	Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376	1
Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1	M. L. Robertson Manager, Electrochemical Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522	1
Dr. M. G. Sceats Department of Chemistry University of Rochester Rochester, New York 14627	1	Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720	1
Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1	Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709	1
Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1	Dr. Denton Elliott Air Force Office of Scientific Research Bldg. 104 Bolling AFB Washington, DC 20332	1
Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1		
Dr. A. Himy NAVSZA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	1		

TECHNICAL REPORT DISTRIBUTION LIST, 051B

	<u>No. Copies</u>		<u>No. Copies</u>
Professor K. Wilson Department of Chemistry, B-014 University of California, San Diego La Jolla, California 92093	1	Dr. B. Vonnegut State University of New York Earth Sciences Building 1400 Washington Avenue Albany, New York 12203	1
Professor C. A. Angell Department of Chemistry Purdue University West Lafayette, Indiana 47907	1	Dr. Hank Loos Laguna Research Laboratory 21421 Stans Lane Laguna Beach, California 92651	1
Professor P. Meijer Department of Physics Catholic University of America Washington, D.C. 20064	1	Dr. John Latham University of Manchester Institute of Science & Technology P.O. Box 88 Manchester, England M601QD	1
Dr. S. Greer Chemistry Department University of Maryland College Park, Maryland 20742	1		
Professor P. Delahay New York University 100 Washington Square East New York, New York 10003	1		
Dr. T. Ashworth Department of Physics South Dakota School of Mines & Technology Rapid City, South Dakota 57701	1		
Dr. G. Gross New Mexico Institute of Mining & Technology Socorro, New Mexico 87801	1		
Dr. J. Kassner Space Science Research Center University of Missouri - Rolla Rolla, Missouri 65401	1		
Dr. J. Telford University of Nevada System Desert Research Institute Lab of Atmospheric Physics Reno, Nevada 89507	1		

DATE
ILME