

AD-A096 640

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/6 12/1

ASYMPTOTIC PROPERTIES OF VOLTERRA EQUATIONS WITH NONINTEGRABLE --ETC(U)

DEC 80 S LONDON

DAAG29-80-C-0041

UNCLASSIFIED

MRC-TSR-2152

NL

For
AD-A
000640



END

DATE

FILED

4-81

DTIC

LEVEL II (2)

AD A 096640

MRC TECHNICAL SUMMARY REPORT #2152

ASYMPTOTIC PROPERTIES OF VOLTERRA
EQUATIONS WITH NONINTEGRABLE KERNELS

Stig-Olof Londen

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

DTIC
SELECTED
MAR 23 1981

December 1980

Received October 31, 1980

UFG FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

81 3 19 086

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

ASYMPTOTIC PROPERTIES OF VOLTERRA EQUATIONS WITH NONINTEGRABLE KERNELS

Stig-Olof Londen

Technical Summary Report #2152
December 1980

ABSTRACT

We study the asymptotic behavior of the solutions of the scalar Volterra integrodifferential equation

$$(E) \quad x'(t) + (a * g(x))(t) = f(t), \quad t > 0, \quad x(0) = x_0,$$

where $a, f : \mathbb{R}^+ \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ are given functions, $*$ denotes convolution and $x : \mathbb{R}^+ \rightarrow \mathbb{R}$ is the solution. We are in particular interested in the largely unsolved case when $a \notin L^1(\mathbb{R}^+)$ and f vanishes at infinity but does not belong to any $L^p(\mathbb{R}^+)$ space for $p < \infty$. The report examines both the linear ($g(x) \equiv x$) and the nonlinear ($g(x) \neq x$) version of (E).

AMS(MOS) Subject Classification: 45D05, 45M05, 45G10

Key Words: Volterra equations, nonlinear integral equations, asymptotic behavior, frequency domain methods

Work Unit Number 1 - Applied Analysis

Accession For	
THIS GRA&I	☒
THIS TAB	☐
Unreferred	☐
Justification	☐
By _____	
Distribution _____	
Availability _____	
Dist _____	
A	

SIGNIFICANCE AND EXPLANATION

In the construction of mathematical models of technical and physical systems one frequently ends up with equations in which the current rate of change $(= \frac{dx(t)}{dt})$ of the state of the system $(= x(t))$ is a function not only of $x(t)$ but also of $x(\tau)$ for past times $\tau < t$. Specifically, one obtains Volterra integrodifferential equations, exemplified by

$$(E) \quad \frac{dx}{dt} + \int_0^t a(t-s)g(x(s))ds = f(t), \quad x(0) = x_0, \quad t \geq 0.$$

Here $f(t)$ is the external input, $a(t)$ is the feedback kernel, $g(x)$ is an in general nonlinear function of x and $x(t)$ is the state of the system at time t .

The key problem concerning (E) is the behavior of $x(t)$ for large values of t . In particular one is interested in whether $x(t)$ tends to zero when $t \rightarrow \infty$ or whether the system keeps oscillating. The present report analyzes these questions. We are in particular interested in the case when the feedback kernel is large, that is when $a(t)$ is not integrable over the positive half-axis. Examples of such kernels often occur in applications where one encounters kernels behaving roughly as $t^{-\alpha}$, for some $0 < \alpha < 1$, for large t .

The second key aspect of this report is that we do allow large input functions $f(t)$. We only assume $f(t) \rightarrow 0$, as $t \rightarrow \infty$ but do not take $|f|^p$ to be integrable over the positive half-axis for any $p < \infty$.

Our main result (Theorem 1) concerns the nonlinear version of (E). We give conditions under which bounded solutions of (E) decay to zero as $t \rightarrow \infty$. We also give a result on the linear version of (E).

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

ASYMPTOTIC PROPERTIES OF VOLTERRA EQUATIONS WITH NONINTEGRABLE KERNELS

Stig-Olof Londen

INTRODUCTION

In this report we examine the asymptotic behavior of the solutions of the scalar Volterra equation

$$(1.1) \quad x'(t) + (a * g(x))(t) = f(t), \quad t > 0, \quad x(0) = x_0,$$

where a, g, f are given functions, x is the solution and where $*$ denotes convolution. Our first result (Theorem 1) concerns the nonlinear case with both a and f big; thus $a \notin L^1(\mathbb{R}^+)$ and f satisfying only (1.7) are not excluded. In Theorem 2 we examine the linear version of (1.1) under the same size conditions.

We begin by stating

Theorem 1. Assume

$$(1.2) \quad g \in C(\mathbb{R}),$$

$$(1.3) \quad a \in BV(\mathbb{R}^+), \quad a(\infty) = 0,$$

$$(1.4) \quad \operatorname{Re} \hat{a} > 0, \quad \omega \in \mathbb{R}, \quad \omega \neq 0$$

$$\text{with } \hat{a}(\omega) = 0 \text{ if } \omega \in \mathbb{Z} \stackrel{\text{def}}{=} \{\omega \mid \operatorname{Re} \hat{a}(\omega) = 0, \quad \omega \neq 0\},$$

$$(1.5) \quad \mathbb{Z} \text{ is countable.}$$

Suppose the differential resolvent $r_a(t)$ of $a(t)$ satisfies

$$(1.6) \quad \begin{aligned} & \text{(i)} \quad r_a \in L^1(\mathbb{R}^+) \\ & \text{(ii)} \quad r'_a \in L^1(\mathbb{R}^+) \\ & \text{(iii)} \quad r''_a \in L^1(\mathbb{R}^+). \end{aligned}$$

and let f be such that

$$(1.7) \quad f \in L^\infty(\mathbb{R}^+) , \quad \lim_{t \rightarrow \infty} f(t) = 0 .$$

Finally let

$$(1.8) \quad x \in (L^\infty \cap \text{LAC})(\mathbb{R}^+) \text{ be a solution of (1.1) a.e. on } \mathbb{R}^+ .$$

Then

$$(1.9) \quad \lim_{t \rightarrow \infty} g(x(t)) = 0 .$$

Above $\hat{a}(\omega) \stackrel{\text{def}}{=} \int_0^\infty e^{-i\omega t} a(t) dt$. By (1.3) $\hat{a}$ is well defined for $\omega \in \mathbb{R}$, $\omega \neq 0$. The differential resolvent $r_a(t)$ is defined as the solution of

$$(1.10) \quad r'_a(t) + (r_a * a)(t) = 0 , \quad r_a(0) = 1 .$$

The question naturally arises whether there exist kernels $a \notin L^1(\mathbb{R}^+)$ and of positive type (i.e. satisfying (1.4)) such that (1.6) holds. By a well-known result [7] the conditions $i\omega + \hat{a}(\omega) \neq 0$, $\omega \in \mathbb{R}$, and (1.11) $a(t)$ positive, nonincreasing and convex on $\mathbb{R}^+$, imply (1.6i). But under the same conditions one does in fact have (1.6ii) and if moreover $-a'(t)$ is convex then (1.6iii) is true. The two last statements are contained in Lemma 1 below which is proved in Section 3.

Lemma 1. Let $a(t)$ be nonnegative, nonincreasing and convex on $\mathbb{R}^+$ with $a(0) < \infty$, $a(\infty) = 0$, and assume that for every fixed $T > 0$ one has that $a(t)$ is not linear in all the intervals $[nT, (n+1)T]$; $n = 0, 1, \dots$. Then (1.4) with Z empty and (1.6i, ii) hold. If in addition $-a'(t)$ is convex then (1.6iii) is true.

Apart from being applicable to equations with kernels $a \notin L^1(\mathbb{R}^+)$ the above theorem also extends recent work [5], [9] done for equations having kernels $a \in L^1(\mathbb{R}^+)$ and a nonhomogeneous term f satisfying only (1.7). To see this we formulate Lemma 2 which follows upon an examination of the proof

of Theorem 1, and the fact that $a \in L^1(\mathbb{R}^+)$ together with (1.4) and $0 \notin \mathbb{Z}$ implies $r_a, r'_a \in L^1(\mathbb{R}^+)$.

Lemma 2. Let (1.2), (1.4), (1.5), (1.6iii), (1.7) and (1.8) hold. In addition assume $a \in L^1(\mathbb{R}^+)$, and $0 \notin \mathbb{Z}$. Then (1.9) is true.

Both [5] and [9] work with more general forms of the convolution term than (1.1). However, in addition to the condition (corresponding to) $a \in L^1(\mathbb{R}^+)$ the results of [9] do require a moment condition on the kernel to be satisfied and the results of [5] require g to be locally Lipschitzian. At the price of (1.6iii) all this has been removed.

The proof of Theorem 1 is essentially based on (1.4), on the size conditions (1.6), and on an asymptotic result obtained in [9]. Observe that the countability assumption (1.5) can be dropped (without the addition of any other hypothesis) if one applies the technique of [4] to the integrated version of (1.1). As this has not been explicitly done we prefer to use [9, Theorem 1b].

One of the main problems concerning the linear equation

$$(1.12) \quad x'(t) + (a * x)(t) = f(t), \quad t \geq 0, \quad x(0) = x_0,$$

is the formulation of hypotheses on the kernel $a(t)$ which imply

$r_a \in L^1(\mathbb{R}^+)$. The most frequently used approach to this hard problem is to give conditions on $a(t)$ which imply that $\hat{r}_a(\omega)$ is sufficiently smooth. But one also has (provided one at first demonstrates that in case f has compact support then $x \in L^\infty(\mathbb{R}^+)$) that $r_a \in L^1(\mathbb{R}^+)$ follows if the implication (1.13) is true [8].

$$(1.13) \quad \left\{ \begin{array}{l} x \in L^\infty(\mathbb{R}^+) \text{ satisfies (1.12)} \\ \text{with } f \text{ satisfying (1.7)} \end{array} \right\} \text{ implies } \left\{ \lim_{t \rightarrow \infty} x(t) = 0 \right\}$$

We show below that under reasonable conditions on the size of the derivatives of the kernel $a(t)$ one may weaken (1.13) to

$$(1.14) \quad \left\{ \begin{array}{l} x \in L^\infty(R^+) \text{ satisfies (1.12)} \\ \text{with } f \text{ satisfying (1.7)} \\ \text{and } \lim_{t \rightarrow \infty} x'(t) = 0 \end{array} \right\} \text{ implies } \left\{ \lim_{t \rightarrow \infty} x(t) = 0 \right\},$$

without altering the conclusion, namely that $r_a \in L^1(R^+)$ holds if the implication (1.14) is true.

A recent article by Gripenberg [2] analyzes the integral resolvent $R_a(t)$ of $a(t)$ by a related approach but under different hypotheses.

Theorem 2. Let $a(t) \in C^1[0, \infty)$, with $a(\infty) = 0$ satisfy

$$(1.15) \quad a' \in L^1(R^+)$$

$$(1.16) \quad a' \in BV(R^+)$$

$$(1.17) \quad \int_{R^+} t |da'(t)| < \infty$$

$$(1.18) \quad \left\{ \begin{array}{l} Z + \hat{a}(Z) \neq 0, \quad \operatorname{Re} Z > 0, \quad Z \neq 0 \\ \\ \lim_{\substack{|Z| \rightarrow 0, \\ \operatorname{Re} Z > 0}} \inf |Z + \hat{a}(Z)| > 0, \end{array} \right.$$

and suppose (1.14) holds. Then $r_a \in L^1(R^+)$.

2. PROOF OF THEOREM 1 .

Convolve (1.1) with r_a and use the fact that r_a satisfies (1.10).

This gives

$$(2.1) \quad (r_a * x')(t) - (r'_a * g(x))(t) = (r_a * f)(t) .$$

An integration of the first term on the left side of (2.1) by parts results in

$$(2.2) \quad x(t) - (r'_a * h(x))(t) = F(t)$$

where $h(x)$, $x \in R$; $F(t)$, $t \in R^+$; are defined by

$$(2.3) \quad h(x) = g(x) - x, \quad F(t) = (r_a * f)(t) + r_a(t)x_0 .$$

From (1.6i,ii), (1.7) and (1.10) follows

$$(2.4) \quad F \in (L^\infty \cap LAC)(R^+), \quad \lim_{t \rightarrow \infty} F(t) = 0$$

$$(2.5) \quad F' \in L^1(R^+), \quad \lim_{t \rightarrow \infty} F'(t) = 0 .$$

Thus

$$(2.6) \quad \lim_{t \rightarrow \infty} \int_0^t F'(\tau) d\tau = -F(0) = -x_0 .$$

Differentiate (2.2), use the fact that $r'_a(0) = 0$ and define $b(t) =$

$-r''_a(t)$. This gives

$$(2.7) \quad x'(t) + (b * h(x))(t) = F'(t) .$$

Then observe that $c(t) \stackrel{\text{def}}{=} \int_0^t b(s) ds = -r'_a(t)$ is such that

$$(2.8) \quad c(\infty) = 0, \quad \int_{R^+} c(t) dt = 1 .$$

From the fact that $b \in L^1(R^+)$ and from (1.10) we have

$$(2.9) \quad \hat{b}(\omega) = i\omega \hat{a}(\omega) [i\omega + \hat{a}(\omega)]^{-1}, \quad \omega \neq 0; \quad \hat{b}(0) = 0 .$$

Note that by (1.4) the transform condition $i\omega + \hat{a}(\omega) \neq 0$ is satisfied for

$\omega \neq 0$. A simple computation gives

$$(2.10) \quad \operatorname{Re} \hat{b}(\omega) = \omega^2 \operatorname{Re}\{\hat{a}(\omega)\} |i\omega + \hat{a}(\omega)|^{-2} \quad \omega \neq 0$$

$$(2.11) \quad \operatorname{Im} \hat{b}(\omega) = [\omega(\operatorname{Re} \hat{a})^2 + \omega^2 \operatorname{Im} \hat{a} + \omega(\operatorname{Im} \hat{a})^2] |i\omega + \hat{a}(\omega)|^{-2}, \quad \omega \neq 0,$$

and so, by (1.4), (2.9)-(2.11),

$$(2.12) \quad \left\{ \begin{array}{l} \operatorname{Re} \hat{b}(\omega) > 0, \quad \omega \in \mathbb{R}; \quad \operatorname{Re} \hat{b}(\omega) = 0 \quad \text{iff} \quad \omega \in \mathbb{Z} \cup \{0\}, \\ \hat{b}(\omega) = 0 \quad \text{if} \quad \operatorname{Re} \hat{b}(\omega) = 0. \end{array} \right.$$

By (1.5), (1.6) - in particular we need (1.6iii) - (1.8), (2.4), (2.5), (2.12)

and as $h \in C(\mathbb{R})$ we may apply [9, Theorem 1b] to get

$$(2.13) \quad \lim_{t \rightarrow \infty} [x(t) + h(x(t)) \int_{\mathbb{R}^+} c(\tau) d\tau] = 0.$$

But if the first part of (2.3) and the second part of (2.8) are used in (2.13) one gets (1.9).

3. PROOF OF LEMMA 1

It is well-known that under the stated hypotheses (1.4) holds with Z empty. See for example [10, p. 170], [3, p. 546]. From [7] we have that (1.6i) holds.

To show that (1.6ii) is satisfied one may at first observe that by (1.6i), (1.10), the monotonicity of $a(t)$ and as $a(0) < \infty$ one has

$r'_a \in L^\infty(\mathbb{R}^+)$. Thus we may Laplace transform r'_a for $\operatorname{Re} s > 0$ and obtain $\tilde{r}'_a(s) = -\tilde{a}(s)[s + \tilde{a}(s)]^{-1}$. (Note that $s + \tilde{a}(s) \neq 0$ for $\operatorname{Re} s > 0$). But by (1.3) $|\tilde{a}(s)| = O(|s|^{-1})$ as $|s| \rightarrow \infty$ and so we conclude by an application of [6, p. 368] that

$$(3.1) \quad r'_a \in L^2(\mathbb{R}^+), \quad \hat{r}'_a(\omega) = -\hat{a}(\omega)[i\omega + \hat{a}(\omega)]^{-1}, \quad \omega \neq 0.$$

To have $r'_a \in L^1(\mathbb{R}^+)$ it suffices to show that $\hat{r}'_a(\omega)$ is locally absolutely continuous on $\mathbb{R}$ and satisfies

$$(3.2) \quad \frac{d}{d\omega} \hat{r}'_a(\omega) \in L^1(\mathbb{R}).$$

But (3.2) follows after straightforward computations which make use of the monotonicity of a and the estimates in [7, Lemma 1].

To prove (1.6iii) one notes at first that the Laplace transform of $c(t) \stackrel{\text{def}}{=} t r''_a(t)$ is the analytic function $\tilde{c}(s) = [s^2 \frac{d\tilde{a}(s)}{ds} + [\tilde{a}(s)]^2][s + \tilde{a}(s)]^{-2}$, $\operatorname{Re} s > 0$. We assert that

$$(3.3) \quad \sup_{0 < x < \infty} \int_{\mathbb{R}} |\tilde{c}(x + iy)|^2 dy < \infty.$$

To see this one observes that

$$(3.4) \quad s^2 \frac{d\tilde{a}(s)}{ds} = -s\tilde{a}(s) - L\{ta''(t) - a'(t)\},$$

that $|\tilde{a}(s)| = O(|s|^{-1})$ and that by the monotonicity of $a(t)$ one has ta'' , $a' \in L^1(\mathbb{R}^+)$. Consequently

$$\sup_{\operatorname{Re} s > 0} |s^2 \frac{d\tilde{a}(s)}{ds}| < \infty \quad \text{and so} \quad \sup_{\operatorname{Re} s > 0} |\tilde{c}(s)| < \infty ,$$

which together with the asymptotic estimate for $\tilde{a}(s)$ gives (3.3). Thus

$c \in L^2(\mathbb{R}^+)$ and

$$(3.5) \quad \hat{c}(\omega) = \frac{[\hat{a}(\omega)]^2 + i\omega^2 \frac{d\hat{a}}{d\omega}}{[\hat{a}(\omega) + i\omega]^2}, \quad \omega \neq 0.$$

To have $c \in L^1(\mathbb{R}^+)$ we need to verify that $\hat{c}$ is locally absolutely continuous and satisfies

$$(3.6) \quad \frac{d\hat{c}(\omega)}{d\omega} \in L^1(\mathbb{R}).$$

From [1, p. 972] we have (under the assumption that $-a'$ is convex) that

$\hat{a}(\omega)$ is twice continuously differentiable and satisfies

$$(3.7) \quad \left| \frac{d^2 \hat{a}(\omega)}{d\omega^2} \right| \leq K \int_0^{1/|\omega|} t^2 a(t) dt, \quad \omega \neq 0$$

for some constant K . Estimating this upwards gives

$$(3.8) \quad \left| \frac{d^2 \hat{a}}{d\omega^2} \right| \leq K|\omega|^{-2} \int_0^{|\omega|^{-1}} a(t) dt \leq K_1 |\omega|^{-2} \hat{a}(\omega)$$

where the second inequality follows from [7, Lemma 1]. Note that we also have

$$(3.9) \quad \left| \frac{d\hat{a}(\omega)}{d\omega} \right| \leq K|\omega|^{-1} |\hat{a}(\omega)|.$$

It now takes some computations which use (3.5), (3.8), (3.9) and the asymptotic estimates

$$\left| \frac{d^{(k)} \hat{a}(\omega)}{d\omega^k} \right| = O(|\omega|^{-k}), \quad \omega \rightarrow \infty, \quad k = 0, 1, 2,$$

to arrive at (3.6).

4. PROOF OF THEOREM 2.

Let $\Gamma(x)$ denote the positive limit set of x , i.e.

$$(4.1) \quad \Gamma(x) = \{y \in L^\infty(R) \mid \text{there exists } t_k \rightarrow \infty \text{ such that } x(t+t_k) \rightarrow y(t) \text{ weak}^* \text{ in } L^\infty(R)\}.$$

We show that

$$(4.2) \quad \text{every } y \in \Gamma(x) \text{ is a constant.}$$

Take $c > 0$ and define Λ_c by

$$(4.3) \quad \Lambda_c(\omega) = \begin{cases} 1 & |\omega| < c \\ 0 & |\omega| > 2c \\ 1 + c^{-1}[c - |\omega|], & c \leq |\omega| \leq 2c \end{cases}$$

There exists $\delta_c(t)$ such that

$$(4.4) \quad \delta_c \in L^1(R), \quad \hat{\delta}_c(\omega) = \Lambda_c(\omega).$$

Define, for $n = 1, 2, \dots$, and $t \geq 0$,

$$(4.5) \quad f_n(t) = e^{-\frac{t}{n}} f(t),$$

thus

$$(4.6) \quad f_n \in L^1(R^+).$$

Let x_n satisfy

$$(4.7) \quad x'_n(t) + (a * x_n)(t) = f_n(t), \quad x_n(0) = x_0,$$

then

$$(4.8) \quad x_n(t) = x_0 r_a(t) + (r_a * f_n)(t), \quad t \geq 0$$

and

$$(4.9) \quad x_n \rightarrow x, \text{ as } n \rightarrow \infty, \text{ uniformly on compact sets of } R^+.$$

From (1.15), (1.16), (1.18) follows (much as in the proof of Lemma 1) that

$$(4.10) \quad r_a(t), \quad \frac{dr}{dt}^a \in L^2(\mathbb{R}^+) ,$$

and so

$$(4.11) \quad \lim_{t \rightarrow \infty} r_a(t) = 0 .$$

Define $x_n = x = f_n = f = 0$ for $t < 0$. By (4.6), (4.10) we may take Fourier transforms in (4.8) and obtain

$$(4.12) \quad \hat{x}_n(\omega) = x_0 \hat{r}_a(\omega) + [i\omega + \hat{a}(\omega)]^{-1} \hat{f}_n(\omega) , \quad \omega \neq 0 .$$

As $\hat{\delta}_c(\omega)[i\omega + \hat{a}(\omega)]^{-1} \in L^2(\mathbb{R})$ we have that there exists b_1 such that

$$(4.13) \quad b_1 \in L^2(\mathbb{R}) , \quad \hat{b}_1(\omega) = \frac{\hat{\delta}_c(\omega)}{i\omega + \hat{a}(\omega)} .$$

Suppose for the moment that there exists $b_2(t)$ satisfying

$$(4.14) \quad b_2 \in L^1(\mathbb{R}) , \quad \hat{b}_2(\omega) = [1 - \hat{\delta}_c(\omega)][i\omega + \hat{a}(\omega)]^{-1} .$$

Then

$$(4.15) \quad x_n(t) = x_0 r_a(t) + \int_0^\infty b_1(t-s) f_n(s) ds + \int_0^\infty b_2(t-s) f_n(s) ds .$$

As $b_1 + b_2 = r = 0$ for $t < 0$ we have $b_1(t) = -b_2(t)$, $t < 0$. But

$$(4.16) \quad \left| \int_t^\infty b_2(t-s) f_n(s) ds \right| < \|b_2\|_{L^1(\mathbb{R})} \|f\|_{L^\infty(\mathbb{R})} , \quad \forall t, n ,$$

and so

$$(4.17) \quad \sup_{t, n} \left| \int_t^\infty b_1(t-s) f_n(s) ds \right| = K < \infty .$$

Also

$$(4.18) \quad \lim_{n \rightarrow \infty} \int_0^\infty b_2(t-s) f_n(s) ds = \int_0^\infty b_2(t-s) f(s) ds ,$$

uniformly for $t \in \mathbb{R}$. Define $g_n(t)$, $g(t)$, $t \in \mathbb{R}$, by

$$(4.19) \quad g_n(t) = \int_0^\infty b_1(t-s)f_n(s)ds, \quad g(t) = x(t) - x_0 x_a(t) - \int_0^\infty b_2(t-s)f(s)ds.$$

Then

$$(4.20) \quad g_n \in L^2(R), \quad g \in L^\infty(R),$$

and so g_n, g represent tempered distributions.

We claim that

$$(4.21) \quad \lim_{n \rightarrow \infty} (g_n - g) = 0 \quad \text{weak}^* \text{ in } S'.$$

From (4.9), (4.15), (4.18), (4.19)

$$(4.22) \quad \lim_{n \rightarrow \infty} (g_n(t) - g(t)) = 0 \quad \text{uniformly on compact sets of } R,$$

and as, by (4.17)

$$(4.23) \quad \begin{cases} |g_n(t)| = \left| \int_0^t + \int_t^\infty \{b_1(t-s)f_n(s)ds\} \right| < t^{1/2} \|b_1\|_{L^2(R)} \|f\|_{L^\infty(R)} + K, & t > 0, \\ |g_n(t)| < K, & t < 0, \end{cases}$$

we conclude that (4.21) holds. But then

$$(4.24) \quad \lim_{n \rightarrow \infty} (\hat{g}_n - \hat{g}) = 0 \quad \text{weak}^* \text{ in } S'.$$

By (4.3), (4.4), (4.13), (4.19)

$$(4.25) \quad \text{supp } \hat{g}_n \subset [-2c, 2c].$$

From (4.24), (4.25) follows

$$(4.26) \quad \text{supp } \hat{g} \subset [-2c, 2c].$$

For $g \in L^\infty(R)$ we denote the spectrum of g (equivalently the support of the distribution Fourier transform) by $\sigma(g)$. The spectrum of a set A is defined as $\sigma(A) = \overline{\bigcup_{\varphi \in A} \sigma(\varphi)}$. For any $y \in L^\infty(R)$ we have $\sigma(\Gamma(y)) \subset \sigma(y)$ and therefore, by (4.26),

$$(4.27) \quad \sigma(\Gamma(g)) \subset \sigma(g) \subset [-2c, 2c].$$

But from (4.11) and the second part of (4.19) follows, as $b_2 \in L^1(\mathbb{R})$ and $f \rightarrow 0$ when $t \rightarrow \infty$,

$$\Gamma(g) = \Gamma(x)$$

and so $\sigma(\Gamma(g)) = \sigma(\Gamma(x))$. Hence $\sigma(\Gamma(x)) \subset [-2c, 2c]$. But c was arbitrary and therefore we conclude that $\sigma(\Gamma(x)) = 0$ which implies (4.2).

We return to the proof of (4.14). By (1.15), (1.18), (4.3), (4.4) we have that $[1 - \hat{\delta}_c(\omega)][i\omega + \hat{a}(\omega)]^{-1}$ is locally the transform of an L^1 -function. Thus we only need to check the behavior of $\hat{b}_2$ at infinity. But if one rewrites $\hat{a}$ as

$$\hat{a}(\omega) = a(0)[i\omega]^{-1} - a'(0)\omega^{-2} - (da')\omega^{-2}, \quad \omega \neq 0$$

and uses (1.17) it follows after some calculations that

$$\frac{d}{d\omega} ([i\omega + \hat{a}(\omega)]^{-1}) \in L^1((-\infty, -2c] \cup [2c, \infty))$$

and so (4.14) is true. The assertion (4.2) is hence valid.

From [8, Theorem 3.1] follows that the proof of Theorem 2 is complete provided we show that if f has compact support then

$$x_0 r_a(t) + (r_a * f)(t) \in L^\infty(\mathbb{R}^+).$$

This however is a consequence of (4.11).

References

1. R. W. Carr and K. B. Hannsgen, A nonhomogeneous integrodifferential equation in Hilbert space, SIAM J. Math. Anal., 10 (1979), 961-984.
2. G. Gripenberg, A Tauberian theorem for a Volterra integral operator. Preprint.
3. K. B. Hannsgen, Indirect Abelian theorems and a linear Volterra equation, Trans. A. M. S., 142 (1969), 539-555.
4. S-O. Londen, On a Volterra integrodifferential equation with L^∞ - perturbation and noncountable zero-set of the transformed kernel, J. Integral Eqs., 1 (1979), 275-280.
5. S-O. Londen, On an integral equation with L^∞ - perturbation. Preprint, Sonderforschungsbereich 123, Institut für Angewandte Mathematik, Universität Heidelberg.
6. W. Rudin, Real and Complex Analysis. McGraw-Hill, New York, 1966.
7. D. F. Shea and S. Wainger, Variants of the Wiener-Lévy theorem, with applications to stability problems for some Volterra integral equations, Amer. J. Math., 97 (1975), 312-343.
8. O. Staffans, On the integrability of the resolvent of a Volterra equation. Report HTKK-MAT-A103 (1977), Institute of Mathematics, Helsinki University of Technology.
9. O. Staffans, On a nonlinear integral equation with a nonintegrable perturbation, J. Integral Eqs., 1 (1979), 291-307.
10. E. C. Titchmarsh, Introduction to the theory of Fourier integrals, Oxford University Press, Oxford, 1937.

SOL/db

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2152v	2. GOVT ACCESSION NO. AD-A096640	3. RECIPIENT'S CATALOG NUMBER 640
4. TITLE (and Subtitle) 6) ASYMPTOTIC PROPERTIES OF VOLTERRA EQUATIONS WITH NONINTEGRABLE KERNELS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) 10) Stig-Olof/Londen		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) 13) DAAG29-80-C-0041v
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 9) Technical summary Repts		12. REPORT DATE Dec 1980
		13. NUMBER OF PAGES 13
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Volterra equations, nonlinear integral equations, asymptotic behavior, frequency domain methods		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We study the asymptotic behavior of the solutions of the scalar Volterra integrodifferential equation (E) $x'(t) + (a * g(x))(t) = f(t), \quad t \geq 0, \quad x(0) = x_0,$ where $a, f: \mathbb{R}^+ \rightarrow \mathbb{R}$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ are given functions, $*$ denotes convolu- tion and $x: \mathbb{R}^+ \rightarrow \mathbb{R}$ is the solution. We are in particular interested in (continued)		

ABSTRACT (continued)

the largely unsolved case when $a \notin L^1(\mathbb{R}^+)$ and f vanishes at infinity but does not belong to any $L^p(\mathbb{R}^+)$ space for $p < \infty$. The report examines both the linear ($g(x) \equiv x$) and the nonlinear ($g(x) \neq x$) version of (E).