

AD-A096 665

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/6 12/1

ON THE CHOICE OF THE EXTERIOR KNOTS IN THE B-SPLINE BASIS FOR A--ETC(U)

DEC 80 J KOZAK

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2148

NL

1 OF 1

40 0 0000H

END
DATE
FILMED
4-81
DTIC

LEVEL II (2)

MRC Technical Summary Report # 2148 ✓

ON THE CHOICE OF THE EXTERIOR KNOTS
IN THE B-SPLINE BASIS
FOR A SPLINE SPACE

J. Kozak

AD A 096665

1472

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

DTIC
ELECTE
MAR 23 1981

December 1980

Received May 29, 1980

FILE COPY

Approved for public release
Distribution unlimited

Sponsored by
U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

Council for International
Exchange of Scholars
Suite 300, Eleven DuPont Circle
Washington, D. C. 20036

81 3 19 082

ON THE CHOICE OF THE EXTERIOR KNOTS IN THE B-SPLINE BASIS
FOR A SPLINE SPACE

J. KOZAK

Technical Summary Report #2148
December 1980

Accession For	
NTIS GRA&I	X
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Special and/or
	Special
A	

ABSTRACT

The B-spline representation of a spline on some interval $[a,b]$ requires the introduction of additional knots outside $]a,b[$ that have nothing in common with the spline space itself. In this note, it is shown that locating all these additional knots at the endpoints a and b of the interval minimizes the matrix norm $\|\cdot\|_{\infty}$ of the matrix $(\lambda_i N_{j,k})^{-1}$ where λ_i are given linear interpolation conditions, and so is preferable when the B-spline representation of a spline interpolant is to be constructed. Such a choice usually also simplifies the algorithms. In particular, one is able to compute stably the B-spline coefficients of a complete spline interpolant by Gauss elimination without pivoting though the corresponding matrix fails to be totally positive or diagonally dominant.

AMS(MOS) Subject Classification - 65D10, 41A15

Key Words: B-spline basis, exterior knots, spline interpolation

Work Unit No. 3 - Numerical Analysis and Computer Science

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041 and Fulbright Grant No. 79-045-A.

SIGNIFICANCE AND EXPLANATION

↓
 The B-spline representation of a spline on some interval $[a,b]$ requires the introduction of additional knots outside $]a,b[$ that have nothing in common with the spline space itself. Nevertheless, their choice influences the numerical accuracy when the B-spline representation

$$\sum \alpha_i N_{i,k,t}$$

of a spline approximant is to be constructed. In this note we prove that locating all the additional knots at the endpoints a and b of the interval minimizes the norm

$$\|(\lambda_i N_{j,k,t})^{-1}\|_{\infty}$$

and is therefore preferable. Here $\underline{\lambda} := (\lambda_i)$ is an arbitrary sequence of linear functionals with support in I except for the assumption that the interpolation problem is correct, i.e. $(\lambda_i N_{j,k,t})$ nonsingular. The above choice usually also simplifies the algorithms. In particular, one is able to compute in a stable way the B-spline coefficients of a complete spline interpolant by Gauss elimination without pivoting though the corresponding matrix fails to be totally positive or diagonally dominant.

The proof relies on the observation that

$$(N_{i,k,\underline{t}}) = (N_{i,k,\underline{t}}) Q_{\underline{t}, \underline{t}}$$

with \underline{t} being two knot sequences that coincide in $[a,b]$. The matrix

$Q_{\underline{t}, \underline{t}}$ for the particular choice of additional knots \underline{t} described above turns to be totally positive and its columns sum up to 1, i.e. $Q_{\underline{t}, \underline{t}}$ is composed of (discrete) B-splines.

p-2

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

ON THE CHOICE OF THE EXTERIOR KNOTS IN THE B-SPLINE BASIS
FOR A SPLINE SPACE

J. KOZAK

1. The Result.

In a practical computation one is rarely able to make statements about the inverse of a given matrix, particularly if the linear system to be solved depends on several free parameters. This note is intended to demonstrate that in an important linear problem, that of spline interpolation, the properties of a chosen basis allow us to draw an interesting conclusion.

To start, let $I := [a, b] \subset \mathbb{R}$ be a given real interval, partitioned by the sequence

$$a =: t_k < t_{k+1} < \dots < t_n < t_{n+1} := b, \quad (1.1)$$

with $t_i < t_{i+k}$, all i and some integer $1 < k < n$. For the purpose of using B-splines, the sequence is extended by

$$t_1 < t_2 < \dots < t_{k-1} < t_k, \quad t_{n+1} < t_{n+2} < \dots < t_{n+k}. \quad (1.2)$$

To simplify the distinction between both parts of $\underline{t} := (t_i)_{i=1}^{n+k}$ put

$\text{int}(\underline{t}) := (t_i)_{i=k}^{n+1}$, $\text{ext}(\underline{t}) := (t_i)_{i=1; i=n+2}^{k-1; n+k}$. The collection of polynomial splines of order k on I with knot sequence \underline{t} is defined by

$$S_{k, \underline{t}}(I) := \{f \mid f|_{[t_i, t_{i+1}]} \text{ is a polynomial of degree } < k, \\ \text{jump}_{t_i} f^{(r)} = 0, r < k - \text{card}\{j \mid t_j = t_i\}, \text{ all } i\}.$$

Any $f \in S_{k, \underline{t}}(I)$ admits a unique B-spline representation

$$f = \sum_{i=1}^n \alpha_i N_{i, k, \underline{t}} \quad (1.3)$$

that has proved very successful in practical computations. Here, $(N_{i, k, \underline{t}})$ is the B-spline partition of the unity, i.e.

$$N_{i, k, \underline{t}}(x) := (t_{i+k} - t_i) [t_i, t_{i+1}, \dots, t_{i+k}] (\cdot - x)_+^{k-1}.$$

Consider now the following interpolation problem: for given interpolation conditions

$\underline{\lambda} := (\lambda_i)_{i=1}^n$ and prescribed numbers $\underline{r} := (r_i)_{i=1}^n \in \mathbb{R}^n$ find $f \in S_{k,\underline{t}}(I)$ such that

$$\lambda_i f = r_i, \quad \text{all } i. \quad (1.4)$$

There $\underline{\lambda}$ is an arbitrary sequence of linear functionals with support in I except for the assumption that the interpolation problem is correct, i.e.

$$f \in S_{k,\underline{t}}(I) \text{ and } \lambda_i f = 0, \text{ all } i, \text{ implies } f = 0.$$

The particular representation (1.3) leads to the solution

$$\underline{\alpha} := (\alpha_i)_{i=1}^n = A_{\underline{t}}^{-1} \underline{r} \quad (1.5)$$

with $A_{\underline{t}} := (\lambda_i N_{j,k,\underline{t}})_{i,j=1}^n$. We note that the solution f does not depend on $\text{ext}(\underline{t})$, but $A_{\underline{t}}$ and consequently $\underline{\alpha}$ do. This fact can influence the numerical accuracy of the computed spline.

Cox [8] considered various ways of choosing the additional knots. As he concluded from numerical evidence, in the case of general-purpose algorithms the coincident choice at end points is preferable. In fact, he observed in [7] that the spectral condition number of the matrix obtained in particular spline least-squares problems is considerably smaller for

$$\text{ext}(\underline{t}) = (\underbrace{t_k, t_k, \dots, t_k}_{k-1}, \underbrace{t_{n+1}, t_{n+1}, \dots, t_{n+1}}_{k-1}) \quad (1.6)$$

compared with an equidistant and an average choice. We note also that in the book by de Boor [5] the choice (1.6) is the rule.

Theorem 1.1. Let the knot sequences $\underline{\tau}, \underline{t}$ satisfy

$$\text{int}(\underline{\tau}) = \text{int}(\underline{t}),$$

$$\tau_i < t_i, \quad i = 1, 2, \dots, k-1,$$

$$\tau_i > t_i, \quad i = n+2, n+3, \dots, n+k,$$

and let $n > 2(k-1)$. If the interpolation problem (1.4) is correct, then

$$\|A_{\underline{\tau}}^{-1}\|_{\infty} > \|A_{\underline{t}}^{-1}\|_{\infty}. \quad (1.7)$$

The inequality (1.7) displays the numerical advantage of using the $\text{ext}(\underline{t})$ as defined in (1.6). Also, following [2], consider the interpolation map

$$P: \mathbb{R}^n \rightarrow S_{k, \underline{t}}(\mathbb{I}) : \underline{r} \rightarrow P\underline{r} := (\lambda_i P\underline{r} = r_i, \text{ all } i).$$

Then

$$\|P\| := \sup_{\|\underline{r}\|_\infty < 1} (\sup_{x \in \mathbb{I}} |P\underline{r}(x)|) = \sup_{\|\underline{r}\|_\infty < 1} (\sup_{x \in \mathbb{I}} |\sum_{i=1}^n \alpha_i N_{i, k, \underline{t}}(x)|) < \|A_{\underline{t}}^{-1}\|_\infty \quad (1.8)$$

and (1.6) gives the best bound of the form (1.8). It is also clear from [2] that $A_{\underline{t}}$ must become singular, as $\tau_1 \rightarrow -\infty$ or $\tau_{n+k} \rightarrow \infty$. We shall explicitly observe

$$\|A_{\underline{t}}^{-1}\|_\infty > \frac{1}{\|A_{\underline{t}}\|_\infty} \max \left(\prod_{r=2}^{k-1} \frac{t_{k+1} - \tau_r}{t_{k+1} - \tau}, \prod_{r=2}^{k-1} \frac{\tau_{n+r} - t_n}{t_{n+r} - t_n} \right). \quad (1.9)$$

Our final observation concerns complete even order spline interpolation. Let $k = 2m$ and

$$\underline{\lambda} := (\delta_{t_k}^{(1)}, \delta_{t_k}^{(1)}, \dots, \delta_{t_k}^{(m-1)}, \delta_{t_{k+1}}, \dots, \delta_{t_n}, \delta_{t_{n+1}}^{(m-1)}, \dots, \delta_{t_{n+1}}^{(m-1)}) \quad (1.10)$$

with $\delta_{t_k}^{(r)} f := f^{(r)}(t_k)$. It is proved in [4] that the Gauss elimination without pivoting can be applied safely if the matrix is totally positive. Unfortunately for $\underline{\lambda}$ given in (1.10) the matrix is not totally positive. But (1.6) gives us $A_{\underline{t}}$ of the form

$$\begin{matrix}
 m & \left\{ \begin{array}{l}
 x \\
 x \ x \\
 x \ x \ x \\
 x \ x \ x \ x \\
 x \ x \ x \ x \ x \\
 \cdot \\
 \cdot \\
 \cdot \\
 x \ x \ x \ x \ x \\
 x \ x \ x \ x \ x \\
 0 \quad x \ x \ x \\
 \quad x \ x \\
 \quad \quad x
 \end{array} \right. \\
 & \left. \right\} m
 \end{matrix}$$

and we need to factor only the submatrix

$$\underline{A}_t \begin{pmatrix} m+1, m+2, \dots, n-m \\ m+1, m+2, \dots, n-m \end{pmatrix},$$

and this submatrix is totally positive.

2. Its Proof.

Consider two knot sequences \underline{t} , $\underline{\tau}$ that satisfy (1.1), (1.2). Assume that $\text{int}(\cdot)$ is strictly increasing. The general case will follow from continuity properties of divided differences. Of course there exists a matrix $Q_{\underline{t} \underline{\tau}}$ such that

$$A_{\underline{\tau}} = A_{\underline{t}} Q_{\underline{t} \underline{\tau}}, \quad (2.1)$$

but it is not so obvious that we can find $Q_{\underline{t} \underline{\tau}}$ explicitly when $\text{int}(\underline{\tau}) = \text{int}(\underline{t})$. Recall Marsden's identity [9]

$$(y-x)^{k-1} = \sum_{i=1}^n \varphi_{i,k,\underline{t}}(y) N_{i,k,\underline{t}}(x), \quad x, y \in I. \quad (2.2)$$

Here

$$\varphi_{i,k,\underline{t}} := \prod_{\ell=1}^{k-1} (\cdot - t_{\ell+i}). \quad (2.3)$$

As observed in [1], (2.2) implies

$$(y-x)_+^{k-1} = \sum_{i=1}^n \prod_{\ell=1}^{k-1} (y - t_{\ell+i})_+ N_{i,k,\underline{t}}(x), \quad y \in \text{int}(\underline{t}), x \in I. \quad (2.4)$$

But (2.4) holds for an $y \leq t_k = a$ too, since then both sides vanish identically. Hence

$$[y_0, y_1, \dots, y_k] (\cdot - x)_+^{k-1} = \sum_{i=1}^n [y_0, y_1, \dots, y_k] \prod_{\ell=1}^{k-1} (\cdot - t_{\ell+i})_+ N_{i,k,\underline{t}}(x) \quad \text{on } I \quad (2.5)$$

under the assumption

$$y := (y_i)_{i=0}^k \in \text{int}(\underline{t}) \cup]-\infty, t_k[.$$

Similarly, with $\underline{y} \in \text{int}(\underline{t}) \cup]t_{n+1}, \infty[$,

$$\begin{aligned} [y_0, y_1, \dots, y_k] (\cdot - x)_+^{k-1} &= [y_0, y_1, \dots, y_k] (x - \cdot)_+^{k-1} = \\ &= \sum_{i=1}^n [y_0, y_1, \dots, y_k] \prod_{\ell=1}^{k-1} (t_{\ell+i} - \cdot)_+ N_{i,k,\underline{t}}(x) \quad \text{on } I. \end{aligned} \quad (2.6)$$

One finds that (2.5), (2.6) are the identity [3, (5.10)], adjusted to the finite interval

I .

Lemma 2.1. Let $n > 2(k-1)$, $\text{int}(\underline{\tau}) = \text{int}(\underline{t})$. Then

$$(N_{i,k,\underline{\tau}})_{i=1}^n = (N_{i,k,\underline{t}})_{i=1}^n Q_{\underline{t} \underline{\tau}} \quad (2.7)$$

with

$$Q_{\underline{t}, \underline{I}} := (q_{ij, \underline{t}, \underline{I}})_{i,j=1}^n$$

and

$$q_{ij, \underline{t}, \underline{I}} := \begin{cases} (\tau_{j+k} - \tau_j) \prod_{\ell=1}^j \varphi_{i, k, \underline{t}}(\tau_{\ell+k}) / \varphi_{j-1, k+2, \underline{I}}(\tau_{\ell+k}), & 1 \leq i \leq j \leq k-1, \\ (\tau_{j+k} - \tau_j) \prod_{\ell=j}^i \varphi_{i, k, \underline{t}}(\tau_{\ell}) / \varphi_{j-1, k+2, \underline{I}}(\tau_{\ell}), & n-k+2 \leq j \leq i \leq n, \\ \delta_{i-j} & \text{otherwise.} \end{cases}$$

Proof. Assume for a moment that $\text{int}(\underline{t})$ is strictly increasing. Choose $1 \leq j \leq k-1$.

Since $n+1 > 2(k-1) + 1 = 2k-1$ we can use (2.5) for any such j to obtain

$$N_{j, k, \underline{I}} = \sum_{i=1}^n (\tau_{j+k} - \tau_j) [\tau_j, \tau_{j+1}, \dots, \tau_{j+k}] \prod_{\ell=1}^{k-1} (\cdot - t_{\ell+1})_+ N_{i, k, \underline{t}} \text{ on } I,$$

and further

$$\begin{aligned} & (\tau_{j+k} - \tau_j) [\tau_j, \tau_{j+1}, \dots, \tau_{j+k}] \prod_{\ell=1}^{k-1} (\cdot - t_{\ell+1})_+ = \\ & = (\tau_{j+k} - \tau_j) \sum_{r=j}^{j+k} \frac{\prod_{\ell=1}^{k-1} (\tau_r - t_{\ell+1})_+}{\prod_{\substack{\ell=j \\ \ell \neq r}}^{j+k} (\tau_r - \tau_{\ell})} \\ & = 0 + (\tau_{j+k} - \tau_j) \sum_{r=i+k}^{j+k} \frac{\prod_{\ell=1}^{k-1} (\tau_r - t_{\ell+1})}{\prod_{\substack{\ell=j \\ \ell \neq r}}^{j+k} (\tau_r - \tau_{\ell})} = q_{ij, \underline{t}, \underline{I}}, \text{ all } i. \end{aligned}$$

The proof for $n-k+2 \leq j \leq n$ follows in the same way from (2.6), and (2.7) obviously holds

for the remaining range of j . The proof is completed for a strictly increasing

$\text{int}(\underline{t})$, but $\text{int}(\underline{I}) = \text{int}(\underline{t})$ simplifies $q_{ij, \underline{t}, \underline{I}}$, say for the range

$1 \leq i \leq j \leq k-1$, to

$$q_{ij, \underline{t}, \underline{I}} = (\tau_{j+k} - \tau_j) [\tau_{i+k}, \tau_{i+k+1}, \dots, \tau_{j+k}] \left(\prod_{\ell=i+1}^{k-1} (\cdot - t_{\ell}) / \prod_{\ell=j}^{k-1} (\cdot - \tau_{\ell}) \right), \quad (2.8)$$

and the general case follows. ■

2. Its Proof.

Consider two knot sequences \underline{t} , $\underline{\tau}$ that satisfy (1.1), (1.2). Assume that $\text{int}(\cdot)$ is strictly increasing. The general case will follow from continuity properties of divided differences. Of course there exists a matrix $Q_{\underline{t} \underline{\tau}}$ such that

$$A_{\underline{\tau}} = A_{\underline{t}} Q_{\underline{t} \underline{\tau}}, \quad (2.1)$$

but it is not so obvious that we can find $Q_{\underline{t} \underline{\tau}}$ explicitly when $\text{int}(\underline{\tau}) = \text{int}(\underline{t})$. Recall Marsden's identity [9]

$$(y-x)^{k-1} = \sum_{i=1}^n \varphi_{i,k,\underline{t}}(y) N_{i,k,\underline{t}}(x), \quad x, y \in I. \quad (2.2)$$

Here:

$$\varphi_{i,k,\underline{t}} := \prod_{\ell=1}^{k-1} (\cdot - t_{\ell+1}). \quad (2.3)$$

As observed in [1], (2.2) implies

$$(y-x)_+^{k-1} = \sum_{i=1}^n \prod_{\ell=1}^{k-1} (y - t_{\ell+1})_+ N_{i,k,\underline{t}}(x), \quad y \in \text{int}(\underline{t}), x \in I. \quad (2.4)$$

But (2.4) holds for an $y < t_k = a$ too, since then both sides vanish identically. Hence

$$[y_0, y_1, \dots, y_k] (\cdot - x)_+^{k-1} = \sum_{i=1}^n [y_0, y_1, \dots, y_k] \prod_{\ell=1}^{k-1} (\cdot - t_{\ell+1})_+ N_{i,k,\underline{t}}(x) \text{ on } I \quad (2.5)$$

under the assumption

$$y := (y_i)_{i=0}^k \in \text{int}(\underline{t}) \cup]-\infty, t_k[.$$

Similarly, with $\underline{y} \in \text{int}(\underline{t}) \cup]t_{n+1}, \infty[$,

$$\begin{aligned} [y_0, y_1, \dots, y_k] (\cdot - x)_+^{k-1} &= [y_0, y_1, \dots, y_k] (x - \cdot)_+^{k-1} = \\ &= \sum_{i=1}^n [y_0, y_1, \dots, y_k] \prod_{\ell=1}^{k-1} (t_{\ell+1} - \cdot)_+ N_{i,k,\underline{t}}(x) \text{ on } I. \end{aligned} \quad (2.6)$$

One finds that (2.5), (2.6) are the identity [3, (5.10)], adjusted to the finite interval

I .

Lemma 2.1. Let $n > 2(k-1)$, $\text{int}(\underline{\tau}) = \text{int}(\underline{t})$. Then

$$(N_{i,k,\underline{\tau}})_{i=1}^n = (N_{i,k,\underline{t}})_{i=1}^n Q_{\underline{t} \underline{\tau}} \quad (2.7)$$

with

In order to investigate $Q_{\underline{t}, \underline{\tau}}$ further we compute, for $1 < i < j < k-1$,

$$\begin{aligned} q_{ij, \underline{t}, \underline{\tau}} &= (\tau_{j+k} - \tau_j) \sum_{\ell=1}^j \varphi_{i, k, \underline{t}}^{(\tau_{\ell+k})} / \varphi_{j-1, k+2, \underline{\tau}}^{(\tau_{\ell+k})} \\ &= \sum_{\ell=1}^j (\tau_{j+k} + \tau_{\ell} - \tau_{\ell} + \tau_j) \varphi_{i, k, \underline{t}}^{(\tau_{\ell+k})} / \varphi_{j-1, k+2, \underline{\tau}}^{(\tau_{\ell+k})} \\ &= - \sum_{\ell=1}^{j-1} \varphi_{i, k, \underline{t}}^{(\tau_{\ell+k})} / \varphi_{j-1, k+1, \underline{\tau}}^{(\tau_{\ell+k})} + \sum_{\ell=1}^j \varphi_{i, k, \underline{t}}^{(\tau_{\ell+k})} / \varphi_{j, k+1, \underline{\tau}}^{(\tau_{\ell+k})}, \end{aligned}$$

and consequently

$$\sum_{j=1}^n q_{ij, \underline{t}, \underline{\tau}} = \sum_{j=1}^{k-1} q_{ij, \underline{t}, \underline{\tau}} = \sum_{\ell=1}^{k-1} \varphi_{i, k, \underline{t}}^{(\tau_{\ell+k})} / \varphi_{k-1, k+1, \underline{\tau}}^{(\tau_{\ell+k})} =: w.$$

Since additionally $\text{int}(\underline{\tau}) = \text{int}(\underline{t})$, then

$$\begin{aligned} w &= \sum_{\ell=1+k}^{2k-1} \frac{\prod_{r=1}^{k-1} (\tau_{\ell} - t_{r+1})}{\prod_{\substack{r=k \\ r \neq \ell}} (\tau_{\ell} - \tau_r)} \\ &= [\tau_k, \tau_{k+1}, \dots, \tau_{2k-1}] \prod_{r=1}^{k-1} (\tau_{\ell} - t_{r+1}) - \sum_{\ell=k}^{i+k-1} \frac{\prod_{r=1}^{k-1} (\tau_{\ell} - t_{r+1})}{\prod_{\substack{r=k \\ r \neq \ell}} (\tau_{\ell} - \tau_r)} \\ &= 1 - 0 = 1. \end{aligned}$$

The following lemma summarizes this observation.

Lemma 2.2. Let $n > 2(k-1)$, and $\text{int}(\underline{\tau}) = \text{int}(\underline{t})$. Then

$$\sum_{j=1}^n q_{ij, \underline{t}, \underline{\tau}} = \sum_{j=1}^n q_{ij, \underline{\tau}, \underline{t}} = 1, \text{ all } i.$$

From Lemma 2.1 we conclude

$$A_{\underline{\tau}} = A_{\underline{t}} Q_{\underline{t}, \underline{\tau}},$$

and to complete now the proof of the theorem it is enough to show that, for the

particular \underline{t} and $\underline{\tau}$ as specified there,

$$Q_{\underline{t}, \underline{\tau}} > 0 := (q_{ij, \underline{t}, \underline{\tau}} > 0, \text{ all } i, j)$$

since then by Lemma 2.2 and invertible $A_{\underline{t}}$

$$\|A_{\underline{\tau}}^{-1}\|_{\infty} = \|Q_{\underline{t}, \underline{\tau}}^{-1} A_{\underline{t}}^{-1}\|_{\infty} > \frac{\|A_{\underline{t}}^{-1}\|_{\infty}}{\|Q_{\underline{t}, \underline{\tau}}\|_{\infty}} = \|A_{\underline{t}}^{-1}\|_{\infty}.$$

Also (1.9) is confirmed just by changing $A_{\underline{t}}$ and $Q_{\underline{t}, \underline{\tau}}$ in the previous argument. It was pointed out to me by de Boor [6] that $Q_{\underline{t}, \underline{\tau}}$ is known to be totally positive if \underline{t} is a refinement of $\underline{\tau}$ on I , i.e. $i.t(\underline{\tau}) = \text{int}(\underline{t})$, $\tau_k < t_1$, $\tau_{n+2} > t_{n+k}$. The following lemma indicates that this fact holds in the more general situation of the Theorem 1.1, and the columns of $Q_{\underline{t}, \underline{\tau}}$ deserve to be called discrete B-splines as in [3].

Lemma 2.3. Under assumptions of the Theorem 1.1

$$q_{ij, \underline{t}, \underline{\tau}} > 0, \quad (-)^{j-i} q_{ij, \underline{\tau}, \underline{t}} > 0, \quad \text{all } i, j. \quad (2.9)$$

Proof. Choose again $1 \leq i < j \leq k-1$. The assumption on $\underline{\tau}, \underline{t}$ reads

$$\tau_1 < t_i < t_k < \tau_{i+k}, \quad 1 \leq i \leq k-1. \quad (2.10)$$

From (2.8) we observe that (2.9) holds for $j = i$. Let $j > i+1$. The elements

$q_{ij, \underline{t}, \underline{\tau}}$ are linear in any t_{ℓ} , $i+1 \leq \ell \leq k-1$. Thus from (2.10)

$$\text{sgn}(q_{ij, \underline{t}, \underline{\tau}}) = \begin{cases} \text{sgn}(q_{ij, \underline{t}, \underline{\tau}})|_{t_{\ell} = \tau_{\ell}} \\ \text{or} \\ \text{sgn}(q_{ij, \underline{t}, \underline{\tau}})|_{t_{\ell} = t_k} \end{cases}$$

or after repeating the argument $k-i-1$ times

$$\left. \begin{aligned} \operatorname{sgn}(q_{ij}, \underline{t}_i) &= \operatorname{sgn}(q_{ij}, \underline{t}_i) \\ t_{i+1} &= \begin{cases} \tau_{i+1} \\ \text{or} \\ t_k \end{cases}, \dots, t_{k-1} = \begin{cases} \tau_{k-1} \\ \text{or} \\ t_k \end{cases} \end{aligned} \right\} = \operatorname{sgn}((\tau_{j+k} - \tau_j) [\tau_{i+k}, \tau_{i+k+1}, \dots, \tau_{j+k}] ((\cdot - t_k)^m / \prod_{\ell=j}^{m+1} (\cdot - \tau_{v_\ell}))) \quad (2.11)$$

for some m , $j-i-1 \leq m \leq k-i-1$, and some $j \leq v_j < v_{j+1} < \dots < v_{i+m} \leq k-1$. For $m = j-i-1$ the argument of $\operatorname{sgn}(\cdot)$ in (2.11) vanishes, and it is enough to consider $m \geq j-i$. Then

$$\begin{aligned} & [\tau_{i+k}, \tau_{i+k+1}, \dots, \tau_{j+k}] ((\cdot - t_k)^m / \prod_{\ell=j}^{m+1} (\cdot - \tau_{v_\ell})) \\ &= [\tau_{i+k}, \tau_{i+k+1}, \dots, \tau_{j+k}] z^{[\tau_{v_j}, \tau_{v_{j+1}}, \dots, \tau_{v_{m+1}}]} w^{\frac{(z-t_k)^m}{z-w}} \\ &= [\tau_{i+k}, \tau_{i+k+1}, \dots, \tau_{j+k}] z^{\frac{(z-t_k)^m}{(z-w)^{m+1+i-j}}} \\ &= [\tau_{i+k}, \tau_{i+k+1}, \dots, \tau_{j+k}] z \sum_{r=0}^m \binom{m}{r} (z-w)^{r-m-i+j-1} (\hat{w}-t_k)^{m-r} \\ &= \frac{1}{(j-i)!} \sum_{r=0}^{m+1-j} \binom{m}{r} (-)^{j-1} \frac{(m-r)!}{(m+1-j-r)!} (\hat{z}-\hat{w})^{r-1-m} (\hat{w}-t_k)^{m-r} + 0 \\ &= (-)^{j-1} \binom{m}{j-i} (\hat{z}-t_k)^{m+1-j} (\hat{w}-t_k)^{j-1} (\hat{z}-\hat{w})^{-m-1} \end{aligned}$$

for some \hat{w}, \hat{z} with $\tau_j \leq \tau_{v_j} < \hat{w} < \tau_{v_{m+1}} \leq \tau_{k-1}$, $\tau_{i+k} < \hat{z} < \tau_{j+k}$. Thus the first inequality in (2.9) is confirmed. The other follows similarly. ■

Acknowledgement.

I thank Professor Carl de Boer for his careful reading and questioning of the manuscript. His suggestions helped me very much in developing the final draft of the paper.

References

1. C. de Boor, On calculating with B-splines, *J. Approximation Theory* 6 (1972), 50-62.
2. C. de Boor, On bounding spline interpolation, *J. Approximation Theory* 14 (1975), 191-203.
3. C. de Boor, Splines as linear combinations of L -splines. A survey, MRC Technical Summary Report No. 1667, 1976.
4. C. de Boor, A. Pinkus, Backward error analysis for totally positive linear systems, *Numer. Math.* 27 (1977), 485-490.
5. C. de Boor, *A Practical Guide to Splines*, Springer-Verlag, 1978.
6. C. de Boor, Private communication.
7. M. G. Cox, Numerical methods for the interpolation and approximation of data by spline functions, Ph. D. Thesis, 1975.
8. M. G. Cox, The incorporation of boundary conditions in spline approximation problems, *NPL NAC Report No. 80*, 1977.
9. M. Marsden, An identity for spline functions and its application to variation diminishing spline approximations, *J. Approximation Theory* 3 (1970), 7-49.

JK/db

14 MRC-TSR-21481

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2148 (6)	2. GOVT ACCESSION NO. AD-A096665	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON THE CHOICE OF THE EXTERIOR KNOTS IN THE B-SPLINE BASIS FOR A SPLINE SPACE.		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) J. Kozak		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041 79-045-A
11. CONTROLLING OFFICE NAME AND ADDRESS See Item 18 below		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 3 - Numerical Analysis and Computer Science
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE December 1980
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		13. NUMBER OF PAGES 10
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
18. SUPPLEMENTARY NOTES U. S. Army Research Office P. O. Box 12211 Research Triangle Park North Carolina 27709		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) B-spline basis, exterior knots, spline interpolation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The B-spline representation of a spline on some interval [a,b] requires the introduction of additional knots outside [a,b] that have nothing in common with the spline space itself. In this note, it is shown that locating all these additional knots at the endpoints a and b of the interval minimizes the matrix norm $\ \cdot \ _{\infty}$. The matrix $(\lambda_i N_{j,k})^{-1}$ where λ_i are given linear interpolation conditions, and so is preferable when the B-spline representation of a spline interpolant is to be constructed. Such a choice usually also simplifies the algorithms. In particular, one is able to compute stably the B-spline coefficients of a complete spline		

221000

20. Abstract (continued)

interpolant by Gauss elimination without pivoting though the corresponding matrix fails to be totally positive or diagonally dominant.

DATE
FILMED
-8