

AD-A096 676

STATE UNIV OF NEW YORK AT BUFFALO DEPT OF CHEMISTRY
KINETICS AND MECHANISM FOR THE ELIMINATION OF HYDROGEN BETWEEN --ETC(U)
MAR 81 O T BEACHLEY
N00014-78-C-0562
NL

UNCLASSIFIED

1 OF 1
AD A
096676



END
DATE
FILMED
4 -81
DTIC

AD A 096676

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 7	2. GOVT ACCESSION NO. AD-A096676	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Kinetics and Mechanism for the Elimination of Hydrogen between Dimethylaluminium Hydride and Benzylamine		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) O. T. Beachley, Jr.		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214		8. CONTRACT OR GRANT NUMBER(s) N-00014-78-C-0562
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-053-686
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) LEVEL		12. REPORT DATE March 19, 1981
		13. NUMBER OF PAGES 15
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for Publication in Inorganic Chemistry		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Organoaluminium(III) Compounds Inorganic polymers Aluminium-nitrogen compounds Kinetics and Mechanism		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The rate of elimination of hydrogen from dimethylaluminium hydride and benzylamine has been measured at -63°C in toluene solution. All experimental data are consistent with a second-order rate law which is complicated by a series of equilibria. Initial rate data were correl-		




DD FORM 1473
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

81 3 23 024

DUC FILE COPY

ated with a rate law and fitted to an appropriate mechanism. The following steps of the mechanism determine the rate of elimination of hydrogen during the early stages of the reaction.



Additional equilibria which involve the aluminium-nitrogen product from the elimination reaction are proposed in order to fully describe the chemistry of the system. The large number of equilibria which complicate the kinetics of the reaction are related to the high basicity of benzylamine.

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By _____	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	

OFFICE OF NAVAL RESEARCH

(15) Contract N-00014-78-C-0562

Task No. NR 053-686

(9) TECHNICAL REPORT NO. 7

(6) Kinetics and Mechanism for the Elimination of
Hydrogen between Dimethylaluminum Hydride
and Benzylamine

by

(10) G. T./Beachley, Jr/

17 TR-7

Prepared for Publication

in

Inorganic Chemistry

[State University of New York at Buffalo
Department of Chemistry
Buffalo, New York 14214

(11) 19, Mar 1981

(12) 21

Reproduction in whole or in part is permitted for
any purpose of the United States Government

*This document has been approved for public release
and sale; its distribution is unlimited

400354




[Contribution from the Department of Chemistry,
State University of New York at Buffalo,
Buffalo, New York 14214].

Kinetics and Mechanism for the Elimination of
Hydrogen between Dimethylaluminium Hydride
and Benzylamine

by
O. T. Beachley, Jr.

Abstract

The rate of elimination of hydrogen from dimethylaluminium hydride and benzylamine has been measured at -63°C in toluene solution. All experimental data are consistent with a second-order rate law which is complicated by a series of equilibria. Initial rate data were correlated with a rate law and fitted to an appropriate mechanism. The following steps of the mechanism determine the rate of elimination of hydrogen during the early stages of the reaction.





Additional equilibria which involve the aluminium-nitrogen product from the elimination reaction are proposed in order to fully describe the chemistry of the system. The large number of equilibria which complicate the kinetics of the reaction are related to the high basicity of benzylamine.

Introduction

The elimination reaction which occurs between an organometallic compound and a protic acid is of fundamental significance and finds many important applications. Despite the importance of this class of reaction to aluminum chemistry, only one kinetics study¹ of a reaction has lead to a proposed mechanism. However, additional kinetics studies with a variety of acid-base pairs are necessary to elucidate the general nature of the elimination step in the mechanism as well as the factors which influence the rate of the elimination reaction and the degree of polymerization of the organometallic product.

The kinetics data for the elimination reaction which occurs between dimethylaluminum hydride and N-methylaniline¹ in toluene at -63°C was consistent with a second order rate law which was complicated by an equilibrium. The following steps of the mechanism determined the rate of elimination of hydrogen. The adduct was assumed to be rapidly formed upon mixing the alane and amine. All results were consistent with the



conclusion that adduct formation was a dead-end path for elimination. Since the elimination reaction is not a reaction of a preformed adduct, both K_d and k alter the rate of formation of hydrogen. The second pertinent conclusion derived from this kinetics study related to the observed formation of only a dimeric product, $[(\text{CH}_3)_2\text{AlN}(\text{CH}_3)(\text{C}_6\text{H}_5)]_2$.

and the predominance of the cis isomer (80%), the kinetically preferred isomer. A $2\pi_s + 2\pi_a$ cycloaddition reaction which minimized interactions between bulky phenyl groups was proposed and, therefore, precluded the formation of higher aluminium-nitrogen polymers.

In this paper we report the kinetics of and propose a mechanism for the elimination reaction which occurs between dimethylaluminium hydride and benzylamine, $N(CH_2C_6H_5)_2$ in toluene solution at $-63^\circ C$. Benzylamine is a significantly stronger base with less bulky substituents than the previously studied N-methylaniline. The goals of our experiments were to determine the nature the aluminium-nitrogen product and the mechanism of the elimination reaction, and to compare the kinetic data for the two amines.

Experimental Section

All compounds were manipulated in a vacuum line or a purified argon atmosphere. The benzylamine was dried over molecular sieves and distilled just prior to use. All other reagents or solvents were prepared or purified as previously described.¹

Reaction of $(CH_3)_2AlH$ with $N(CH_2C_6H_5)_2$ - The stoichiometry of the elimination reaction between $(CH_3)_2AlH$ and $N(CH_2C_6H_5)_2$ was examined and the product was fully characterized. When 0.1890 g (3.26 mmol) of $(CH_3)_2AlH$ was combined with 0.3508 g (3.27 mmol) of $N(CH_2C_6H_5)_2$ in toluene solution, 3.24 mmol of H_2 (measured with a Toepler pump and gas buret assembly) was formed. No methane was observed as a product. Additional experiments with excess benzylamine confirmed the stoichiometry.

The aluminium-nitrogen product, $[(CH_3)_2AlN(CH_2C_6H_5)H]_2$, was fully characterized. The colorless, crystalline solid has a melting point of 65-68°C. Decomposition as evidenced by bubbling started at about 140°C. Cryoscopic molecular weight measurements in benzene solution using an apparatus similar to that described by Shriver² demonstrated the compound to exist as a dimer. The following experimental data were observed: $[(CH_3)_2AlN(CH_2C_6H_5)]_2$ - formula weight - 327: [Calc. molality, (observed molecular weight)] 0.0968(327); 0.0604(318); 0.0439(317). Our 1H NMR measurements of $[(CH_3)_2AlN(CH_2C_6H_5)]_2$ dissolved in toluene, cyclopentane and CH_2Cl_2 suggest that the compound exists as a mixture of cis and trans isomers but the ratio of isomers has essentially no dependence on the polarity of the solvent (Table I). The infrared spectrum of $[(CH_3)_2AlN(CH_2C_6H_5)]_2$ was recorded in the range 4000-250 cm^{-1} by means of a Perkin-Elmer Model 457 spectrometer as a Nujol mull using CsI plates. The following are the spectral data (Bands due to the Nujol have been omitted): 3262(m); 1264(m); 1192(vs); 1167(sh); 1082(sh); 1044(sh); 1017(vs); 970(m); 895(w); 842(sh); 802(s); 737(sh); 685(vs); 632(sh); 562(m); 522(w); 492(w); 452(m).

Kinetics Experiments: The kinetics of the reaction between dimethylaluminium hydride and benzylamine in toluene at -63°C was monitored by following the rate of evolution of hydrogen manometrically. The experimental procedure for adding the amine to the alane solution and following the reaction was identical to that previously described.¹ As in the earlier study all experimental variables which might alter pressure measurements were maintained as constant as possible.

Table I

^1H NMR Spectral Data^a $[(\text{CH}_3)_2\text{AlNH}(\text{CH}_2\text{C}_6\text{H}_5)]_2$


<u>Assignment</u>	<u>Solvent</u>		
	<u>$\text{C}_6\text{H}_5\text{CH}_3$</u>	<u>C_5H_{10}</u>	<u>CH_2Cl_2</u>
AlCH_3 cis	10.30	10.64	10.61
	10.33	10.72	10.66
	10.36	10.80	10.71
N-H	8.78	b	8.64
CH_2	6.03	6.11	5.97
		6.13	6.01
	6.12	6.19	6.07
		6.21	
C_6H_5	2.73	2.82	2.43
	2.78		

^aAll data are given as τ (ppm) and are referenced to tetramethylsilane (10.00 ppm). The spectra were recorded at 90 MHz and ambient temperature with a Varian Model EM-390 spectrometer.

^bThe N-H line was not observed due to interference from cyclopentane.

Results and Discussion

The general elimination reaction which occurs between dimethylaluminium hydride and benzylamine can be summarized by the following equation. The kinetics of this reaction were investigated by following



the rates of evolution of hydrogen at -63°C in toluene solution. A variety of experimental conditions were attempted. When the amine was in excess of the alane, a clear solution was formed and hydrogen was evolved smoothly. However, when the initial concentration of the alane and amine were equal, a white precipitate was observed in the reaction vessel but no hydrogen was formed. Experiments using an excess of the alane were prohibited because a different final product would be formed.³ Thus, all kinetics data had to be derived from experiments which employed a greater than 10-fold excess of amine.

The kinetics data are summarized in Table II. Only plots of hydrogen pressure vs. time for the early part of the reaction give data which can be correlated with a rate law and fitted to an appropriate mechanism. First, second and fractional order kinetic plots had distinct curvatures. The slope of the linear portion of pressure vs. time plots gives an observed rate which is proportional to the initial dimethylaluminium hydride concentration for a given amine concentration. Thus, the alane dependence is given by the following expression. A plot of $1/k_0$ vs.

$$\frac{\Delta P_{\text{H}_2}}{\Delta t} = k_0 [(\text{CH}_3)_2\text{AlH}]_T \quad (\text{Constant } [\text{N}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}_2]_T) \quad (4)$$

Table II

Kinetic Data for $(\text{CH}_3)_2\text{AlH}-\text{N}(\text{CH}_2\text{C}_6\text{H}_5)_2$ Reaction

$[(\text{CH}_3)_2\text{AlH}]_T,^a$ M	$[\text{N}(\text{CH}_2\text{C}_6\text{H}_5)_2]_T,$ M	10^3 Initial Rate mm/sec	$10^5 k_o^b$ sec ⁻¹	$10^4 kK_2^c$ sec ⁻¹
0.123	1.94	5.23	5.92	1.95
0.0891	1.68	3.83	5.99	1.82
0.0833	1.23	4.38	7.31	1.82
0.0819	1.07	4.52	7.67	1.76
0.0819	2.00	3.13	5.32	1.82
0.0776	1.44	3.42	6.12	1.68
0.0675	0.924	4.27	8.79	1.86
0.0603	1.39	2.68	6.19	1.66
0.0546	0.720	4.07	10.36	1.94
0.0575	2.03	2.08	5.03	1.74








a - Concentrations of $(\text{CH}_3)_2\text{AlH}$ are based on the number of moles of the monomeric unit.

b - Calculated by dividing the initial rate, converted to units of M/sec, by $[(\text{CH}_3)_2\text{AlH}]_T$. (See text).

c - Calculated from the rate law by using the calculated value of K_1 (1.21 M^{-1}), initial rate, $[(\text{CH}_3)_2\text{AlH}]_T$ and $[\text{N}(\text{CH}_2\text{C}_6\text{H}_5)_2]_T$. (See text).

$[N(CH_2C_6H_5)_2H_2]_T$ is linear and demonstrates the inverse dependence of the rate of hydrogen evolution on the amine concentration. The terms $[(CH_3)_2AlH]_T$ and $[N(CH_2C_6H_5)_2H_2]_T$ express the total dimethylaluminum hydride and benzylamine concentrations before elimination.

Equations 5-12 can be used to explain all of the kinetic data and experimental observations, and are totally consistent with the chemistry of the system. The only assumption which must be made for this mechanism is that the initial formation of adduct is extremely rapid.^{1,4,5}



$$\frac{dP_{H_2}}{dt} = \frac{-d[(CH_3)_2AlH]_T}{dt} = \frac{k[(CH_3)_2AlH]_T[N(CH_2C_6H_5)_2H_2]_T}{\frac{K_1[N(CH_2C_6H_5)_2H_2]_T^2}{K_2} + \frac{[N(CH_2C_6H_5)_2H_2]_T}{K_2} + 1} \quad (13)$$

The rate law is appropriate during the early stages of a reaction which utilizes a large excess of amine. Equilibria for K_3 and K_4 (eq. 10 and 11) are included in the mechanism in order to describe the overall chemistry of the system but have been eliminated from the rate law in order to fit the observed initial rate data. If K_2 is assumed to be small, the rate law can be simplified even further to give the following expression which is consistent with the data.

$$\frac{dP_{H_2}}{dt} = \frac{kK_2[(CH_3)_2AlH]_T}{K_1[N(CH_2C_6H_5)H_2]_T + 1} \quad (14)$$

Calculated Kinetic Constants

$$kK_2 = 1.80 \times 10^{-4} \text{ sec}^{-1}$$

$$K_1 = 1.21 \text{ M}^{-1}$$

The kinetic constants kK_2 and K_1 are calculated from the slope and intercept of the $1/k_0$ vs $[N(CH_2C_6H_5)H_2]_T$ plot. Using this value of K_1 , kK_2 can be recalculated from the initial concentrations and rate data for the individual experiments (Table II). The results of these calculations support the proposed mechanism and rate law. It is regrettable that it is not possible to calculate values of k and K_2 from the available data.

The proposed mechanism for the elimination reaction between dimethylaluminum hydride and benzylamine is analagous to that proposed for the N-methylaniline reaction¹ except that equilibria (K_1 , K_3 and K_4) have been added. The steps of the mechanism which are required for H_2 formation (eq. 8 and 9) are identical to those proposed for the

$\text{H}(\text{CH}_3)_2\text{Al}-\text{N}(\text{CH}_3)(\text{C}_6\text{H}_5)\text{H}$ reaction.¹ The equilibrium K_1 involving the bisamine adduct, $\text{H}(\text{CH}_3)_2\text{Al}[\text{N}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}_2]_2$, is necessary to explain the inverse dependence of the rate of hydrogen evolution on the amine concentration under the pseudo first order conditions (excess amine). Equilibria K_3 and K_4 , which occur after H_2 has been eliminated, pertain to overall chemistry of the system. The complications of five equilibria in the mechanism are responsible for the reaction not following simple first order kinetics under pseudo first order conditions. The occurrence of these equilibria for the benzylamine reaction are consistent with its significantly high basicity due to the electronic and steric effects. For comparison purposes, equilibrium constants⁶ which describe the basicity of benzylamine and N-methylaniline toward the proton in aqueous solution differ by 10^5 .


The observed decreasing initial rate of hydrogen evolution with increasing benzylamine concentration at constant alane concentration is a consequence of equilibrium K_1 - eq. 7. The excess amine reduces the concentration of the adduct, $\text{H}(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}_2$, the compound which must dissociate to form the species required for the elimination reaction, the monomeric alane and amine. Bisamine adducts which involve five-coordinate aluminium are known⁷⁻¹² but a strong base and minimum steric interactions between the acid and base are necessary. Two compounds,^{13,14} $\text{H}_3\text{Al}[\text{N}(\text{CH}_3)_3]_2$ and $\text{H}_3\text{Al}[(\text{CH}_3)_2\text{NC}_2\text{H}_4\text{N}(\text{CH}_3)_2]$, have been characterized by X-ray structural studies. However, in the case of dimethylaluminium hydride-amine compounds, no bisamine adducts have been previously observed. Apparently, $\text{N}(\text{CH}_3)_3$ ¹⁵ and $\text{N}(\text{C}_6\text{H}_5)(\text{CH}_3)\text{H}$ ¹ have too many steric interactions or are too weak a base toward $(\text{CH}_3)_2\text{AlH}$ for a

bisamine adduct to be formed even at -46°C or -63°C , respectively.

The step in the mechanism which leads to the formation of hydrogen (eq. 9) is considered to be a second order reaction between a monomeric alane and the amine. It was not possible to obtain direct kinetic data for this second order process by following the formation of hydrogen when the alane and amine concentrations were equal.¹ However, other experimental observations support our hypothesis. When the alane and benzylamine are in equal concentrations, an insoluble solid is formed upon mixing at -63°C but no hydrogen is eliminated. The insoluble solid is most likely the adduct, $\text{H}(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)_2$, a very polar compound. The insolubility of this adduct and the equilibria which precede the actual elimination reaction (eq. 7 and 8; K_1 and K_2) must reduce the concentrations of the reactive monomeric alane and amine to effectively hinder the formation of hydrogen. If the elimination reaction was a first order reaction of the preformed adduct, hydrogen formation should have been observed even if the adduct were insoluble. Second order reactions of the adduct are inconsistent with the kinetic data.

The equilibria represented by K_3 , K_4 and K_5 (eq. 10, 11 and 12) have been added to the overall mechanism of the total reaction in order to fully describe the chemistry of the system. These equilibria occur after hydrogen has been formed and involve the aluminium-nitrogen products of the reaction. Thus, they are not required to explain our kinetic observations for the initial part of the reaction. However, these complicating equilibria help to explain why the full reaction does not exhibit simple first order kinetics under pseudo first order condi-

tions. The fact that these equilibria occur in the benzylamine-alane elimination reaction but not in the corresponding N-methylaniline reaction¹ must be related to the higher basicity of benzylamine. The reaction of $(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}$ with $\text{N}(\text{CH}_2\text{C}_6\text{H}_5)_2$ (eq. 10, K_3) is clearly supported by ^1H NMR data (see following discussion). The subsequent reaction of $\text{H}_2(\text{C}_6\text{H}_5\text{CH}_2)\text{Al}(\text{CH}_3)_2\text{N}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}$ with $(\text{CH}_3)_2\text{AlH}$ would be a function of the increased basicity of the benzylamino group after elimination. This suggestion that the benzylamino-group can effectively compete with benzylamine for $(\text{CH}_3)_2\text{AlH}$ is based on related observations in aluminium-nitrogen chemistry.^{16,17} The analogous effect is considered to be responsible for the dimerization of $(\text{CH}_3)_2\text{AlN}(\text{CH}_3)\text{C}_2\text{H}_4\text{N}(\text{CH}_3)_2$ with free $-\text{N}(\text{CH}_3)_2$ groups, rather than the formation of the chelated monomer.¹⁷ Additional complications in the overall mechanism can be envisioned if a second elimination reaction occurred as shown in the following equation (15). A subsequent proton transfer reaction and



then dissociation of this product would give $(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}$, the typical reaction product.

The comparisons of the calculated kinetic constants for the analogous benzylamine- and N-methylaniline¹-dimethylaluminium hydride elimination reactions are of interest to determine how the amine effects the rate of elimination. Qualitatively, benzylamine reacts faster than N-methylaniline. Similarly, these observations are reflected in the calculated values of kK_2 ($1.80 \times 10^{-4} \text{ sec}^{-1} \text{ N}(\text{CH}_2\text{C}_6\text{H}_5)_2$; 5.46×10^{-5}

$\text{sec}^{-1} \text{N}(\text{C}_6\text{H}_5)(\text{CH}_3)\text{H}$. Benzylamine is a much stronger base than N-methylaniline. Thus, K_2 should be smaller for the benzylamine, probably less than $9.09 \times 10^{-3} \text{ M}$, the observed value of K_2 for the $\text{H}(\text{CH}_3)_2\text{AlN}(\text{C}_6\text{H}_5)(\text{CH}_3)\text{H}$ dissociation equilibrium.¹ In order for kK_2 to be larger for the benzylamine reaction, the rate constant for its elimination reaction (k_2) must be greater than $6.01 \times 10^{-3} \text{ M}^{-1} \text{ sec}^{-1}$, the value calculated for the N-methylaniline reaction.¹ There are many reasons why dimethylaluminium hydride-benzylamine would eliminate hydrogen faster than N-methylaniline. There are twice as many reactive N-H bonds with less steric hindrance for the formation of the transition state for the proposed four-centered S_{Ei} process.¹ Our current limited data does not warrant more detailed comparisons at this time.

The final product from the elimination reaction in the absence of excess amine is an aluminium-nitrogen dimer $[(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}]_2$, which exists as a 1:1 mixture of cis/trans geometrical isomers. This isomer ratio is not significantly affected by solvent polarity. The ^1H NMR data indicate 49.5% cis isomer in cyclopentane, 47.9% cis in toluene and 51.8% cis in methylene chloride. If a $2\pi_{\text{s}} + 2\pi_{\text{a}}$ concerted cycloaddition reaction of two $(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}$ monomer units is the mechanism for dimerization,¹ only small differences can exist between the relative energies of the transition states for the formation of the two isomers. The benzyl group would be expected to have small steric effects in the proposed transition state. However, if the monomer, $(\text{CH}_3)_2\text{AlN}(\text{CH}_2\text{C}_6\text{H}_5)\text{H}$, reacts with $\text{N}(\text{CH}_2\text{C}_6\text{H}_5)_2$ as proposed in eq. 10 (K_3), subsequent sequential reactions can also be envisioned to produce the dimer. The current kinetic study does not permit us to distinguish

between these two paths but the concerted cycloaddition reaction seems more plausible, especially when the alane and amine are reacted in equal amounts. When excess benzylamine is added to a cyclopentane solution of the dimer, the unique aluminium-methyl lines of the cis/trans isomers in the ^1H NMR spectrum are lost. Only one Al-CH_3 line is observed, which is consistent with fast exchange. As previously mentioned, the equilibria represented by equations 10 and 12 (K_3 and K_5) are consistent with these observations.

The reaction between an alane and amine cannot be considered simple or straightforward. Many factors must influence the various steps of the reaction mechanism. As the basicity and steric effects of the amine change, many equilibria involving reactants and products are apparently introduced into the mechanism. More kinetics studies will be required to elucidate the relative importance of the different equilibria and the factors influencing the magnitude of the second-order rate constant for elimination.

Acknowledgment. This work was supported in part by the Office of Naval Research. We are indebted to Professor Gordon M. Harris for many helpful discussions on the interpretation of the kinetic data.

References

1. Beachley, O. T., Jr.; Tessier-Youngs, C. Inorg. Chem. 1979, 18, 3188.
2. Shriver, D. F. "The Manipulation of Air Sensitive Compounds"; McGraw-Hill: 1969; p. 95.
3. Bernstein, J. D. Ph.D. Thesis. SUNY-Buffalo, Buffalo, N. Y., 1975.
4. Eisch, J. J.; Rhee, S.-G. J. Am. Chem. Soc. 1974, 96, 7276.
5. Eisch, J. J.; Rhee, S.-G. J. Am. Chem. Soc. 1975, 97, 4673.
6. Handbook of Chemistry and Physics, CRC Press; 1974; p. D-147.
7. Wiberg, E.; Graf, H.; Schmidt, M.; Uson, R. Z. Naturforsch 1952, 7b, 578.
8. Fetter, N. R.; Bartocha, B.; Brinckman, F. E.; Moore, D. W. Canad. J. Chem. 1963, 41, 1359.
9. Davidson, J. M.; Wartik, T. J. Am. Chem. Soc. 1960, 82, 5506.
10. Fetter, N. R.; Moore, D. W. Canad. J. Chem. 1964, 42, 885.
11. Young, A. R.; Ehrlich, R. Inorg. Chem. 1965, 4, 1358.
12. Ashby, E. C. J. Am. Chem. Soc. 1964, 86, 882.
13. Heitsch, C. W.; Nordman, C. E.; Parry, R. W. Inorg. Chem. 1963, 2, 508.
14. Palenik, G. J. Acta Cryst. 1964, 17, 1573.
15. Beachley, O. T. Jr.; Bernstein, J. D. Inorg. Chem. 1973, 12, 183.
16. Beachley, O. T. Jr.; Racette, K. C. Inorg. Chem. 1975, 14, 2534.
17. Beachley, O. T. Jr.; Racette, K. C. Inorg. Chem. 1976, 15, 2110.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D. C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D. C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D. C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D. C. 20360	1	Naval Ship Research & Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Divison, Code 6100 Naval Research Laboratory Washington, D. C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

No.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco, Calif. 96503 1

Mr. James Kelly
DTNSRDC Code 2804
Annapolis, Maryland 21402 1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. N. Grimes Department of Chemistry University of Virginia Charlottesville, Virginia 22901	1	Professor H. Abrahamson Department of Chemistry University of Oklahoma Norman, Oklahoma 73019	1
Dr. M. F. Hawthorne Department of Chemistry University of California Los Angeles, California 90024	1	Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. D. B. Brown Department of Chemistry University of Vermont Burlington, Vermont 05401	1	Dr. B. Foxman Department of Chemistry Brandeis University Waltham, Massachusetts 02154	1
Dr. W. B. Fox Chemistry Division Naval Research Laboratory Code 6130 Washington, D. C. 20375	1	Dr. T. Marks Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. J. Adcock Department of Chemistry University of Tennessee Knoxville, Tennessee 37916	1	Dr. G. Geoffrey Department of Chemistry Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. A. Cowley Department of Chemistry University of Texas Austin, Texas 78712	1	Dr. J. Zuckerman Department of Chemistry University of Oklahoma Norman, Oklahoma	1
Dr. W. Hatfield Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. D. Seyferth Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Professor Ralph Rudolph Department of Chemistry University of Michigan Ann Arbor, Michigan 48109	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Professor Richard Eisenberg Department of Chemistry University of Rochester Rochester, New York 14627	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1

