


VIRGINIA UNIV CHARLOTTESVILLE DEPT OF ENGINEERING SC--ETC F/6 12/2
MULTIPLE OBJECTIVE EVALUATION AND CHOICEMAKING UNDER RISK WITH --ETC(U)
FEB 81 C C WHITE, A P SAGE N00014-80-C-0542
UVA/SE-81-2 NL

UNCLASSIFIED

NL

1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 2644, 2645, 2646, 2647, 2648, 2649, 2650, 2651, 2652, 2653, 2654, 2655, 2656, 2657, 2658, 2659, 2660, 2661, 2662, 2663, 2664, 2665, 2666, 2667, 2668, 2669, 2670, 2671, 2672, 2673, 2674, 2675, 26

END
DATE
OF MED.
5-8
PTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 097418

SYSTEMS ENGINEERING RESEARCH REPORT 81-2

MULTIPLE OBJECTIVE EVALUATION AND CHOICEMAKING
UNDER RISK WITH PARTIAL PREFERENCE INFORMATION

Chelsea C. White
Andrew P. Sage


Supported by:

Engineering Psychology Programs
Office of Naval Research
ONR Contract Number N0014-80-C-0542
Work Unit Number NR-197-065

Approved for Public Release
Distribution Unlimited
Reproduction in Whole or in Part is
Permitted for Any Use of the U.S. Government

February 1981

DTIC
SELECT
S APR 7 1981



SCHOOL OF ENGINEERING AND
APPLIED SCIENCE

DEPARTMENT OF ENGINEERING SCIENCE AND SYSTEMS

UNIVERSITY OF VIRGINIA
CHARLOTTESVILLE, VIRGINIA 22901

81 3 26 038

+ DTIC FILE COPY

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER UVA/SE-81-2	2. GOVT ACCESSION NO. AD-A097418	3. RECIPIENT'S CATALOG NUMBER 19
4. TITLE (and Subtitle) MULTIPLE OBJECTIVE EVALUATION AND CHOICEMAKING UNDER RISK WITH PARTIAL PREFERENCE INFORMATION		5. DATE OF REPORT (or Period Covered) Research Report
7. AUTHOR(s) Chelsea C. White, III Andrew P. Sage		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Engineering Science and Systems University of Virginia Charlottesville, Virginia 22901		8. CONTRACT OR GRANT NUMBER(s) NA14-80-C-0542
11. CONTROLLING OFFICE NAME AND ADDRESS Engr. Psychology Programs Department of the Navy, Office of Naval Research Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-197-065
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE February 1981
		13. NUMBER OF PAGES 150
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES None		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multiple Objectives Decision Aiding Under Risk Partial Preference Information		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper, we consider a single stage, multiattribute decision making problem under risk. Each outcome is associated with a vector of value scores. The set of value score vectors is partially ordered in the usual manner. This partial order is used to induce partial orders on the set of alternatives based on: expected value score, first order stochastic dominance, and second order stochastic dominance. We show that properly aggregating value scores produces a more refined partial order on the set of value		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

401907

20.

score vectors, which then induces a more refined partial order on the alternative set. This result forms the basis of a decision aiding procedure which: (1) identifies the nondominated set of alternatives, a set guaranteed to contain the most preferred alternative under mild assumptions, (2) asks the decision maker to choose a most preferred alternative from the nondominated set, and (3) if this choice cannot be made, aggregates some, but not necessarily all, of the value scores to strengthen the partial order on the alternative set, thus reducing the size of the nondominated set and presumably enhancing alternative selection. The potential value of this procedure is that it is interactive, it can accommodate a variety of levels of online preference feedback from the decision maker, and it does not necessarily require that the value scores be completely aggregated.

MULTIPLE OBJECTIVE EVALUATION AND CHOICEMAKING
UNDER RISK WITH PARTIAL PREFERENCE INFORMATION*

Chelsea C. White, III

and

Andrew P. Sage

Department of Engineering Science and Systems
Thornton Hall
University of Virginia
Charlottesville, VA 22901
(804) 924-5395

ABSTRACT: In this paper, we consider a single stage, multi-attribute decision making problem under risk. Each outcome is associated with a vector of value scores. The set of value score vectors is partially ordered in the usual manner. This partial order is used to induce partial orders on the set of alternatives based on: expected value score, first order stochastic dominance, and second order stochastic dominance. We show that properly aggregating value scores produces a more refined partial order on the set of value score vectors, which then induces a more refined partial order on the alternative set. This result forms the basis of a decision aiding procedure which: (1) identifies the nondominated set of alternatives, a set guaranteed to contain the most preferred alternative under mild assumptions, (2) asks the decision maker to choose a most preferred alternative from the nondominated set, and (3) if this choice cannot be made, aggregates some, but not necessarily all, of the value scores to strengthen the partial order on the alternative set, thus reducing the size of the nondominated set and presumably enhancing alternative selection. The potential value of this procedure is that it is interactive, it can accommodate a variety of levels of on-line preference feedback from the decision maker, and it does not necessarily require that the value scores be completely aggregated.

*This research has been supported by ONR Contract N0014-80-C-0542.

1. Introduction

The issue addressed in this paper involves the problem of selecting the k most preferred alternatives from a finite set of alternatives, where the outcome which results from any selected alternative is determined probabilistically and where the problem is characterized by multiple, noncommensurate, and conflicting objectives. Specifically, let $\Pi = \{\pi_1, \dots, \pi_p\}$ be the finite set of all alternatives available to the decision-maker, where $k < p$. The set of all possible outcomes of each of the alternatives is $\Omega = \{\omega_1, \dots, \omega_M\}$. Assume that N lowest level objectives and Q attributes or objectives measures have been identified. Let $x_q(\omega_m)$ be the q^{th} attribute score if the m^{th} outcome has occurred, and let $x(\omega_m) = \{x_1(\omega_m), \dots, x_Q(\omega_m)\}$. Assume for each objective n , $1 \leq n \leq N$, a value function v_n has been assessed that is isotone (monotonically nondecreasing) in preference; that is, outcome m is preferred to outcome ℓ with respect to the n^{th} objective iff (if and only if) $v_n[x(\omega_m)] > v_n[x(\omega_\ell)]$. We remark it is commonly considered desirable that each value score be associated with a single attribute which is distinct from the attributes associated with the other objectives, (Keeney and Raiffa, 1976; p. 223); that is, $Q = N$ and $v_n[x(\omega_m)] = v_n[x_n(\omega_m)]$. We will not require this as an assumption. Let $v^m = \{v_1[x(\omega_m)], \dots, v_N[x(\omega_m)]\} \in \mathbb{R}^N$ and $V = \{v^m, m=1, \dots, M\}$. The finite set V is the set of all value score vectors and for our purposes is essentially equivalent to the

outcome set Ω . Since more is better with respect to value scores, the relation on V considered throughout this paper is: $v' R v$ iff $v'_n \geq v_n$ for all $n=1, \dots, N$. Each alternative is equivalent to a probability mass function over Ω and hence equivalently over V ; i.e., $\pi_p(m) = \pi_p(v^m)$ is the probability that the p^{th} alternative will result in value score vector v^m . All (subjective or objective) probabilities $\pi_p(m)$ are assumed to have been determined.

We assume that there exists a scalar-valued function $U: V \rightarrow \mathbb{R}$ such that

$$u(x) = U[v(x)]$$

where u is a multiattribute utility function and where both u and U are assumed to be unassessed. Furthermore, we assume that an objectives hierarchy has been specified and that value scores can be aggregated according to this objectives hierarchy. For example, let v_{N-1} and v_N be value scores associated with two lowest level objectives which are both associated with a single higher level objective. (The two lowest level objectives might be "increase comfort" and "decrease noise", and the higher level objective might be "improve aesthetics.") Then we assume that there exist an aggregation function $f: \mathbb{R}^2 \rightarrow \mathbb{R}^1$ and a utility function $\bar{U}$ such that

$$\begin{aligned} U(v_1, \dots, v_N) \\ = \bar{U}[v_1, \dots, v_{N-1}, f(v_{N-1}, v_N)]. \end{aligned}$$

These assumptions allow us to "move up" the objectives hierarchy and ultimately assess a scalar-valued value function $\bar{v}$ such that




FIGURE 1: Decision Tree Representation of the Static Choicemaking Problem Under Risk with Vector-Valued Outcomes.

$u(x) = u[\bar{v}(x)]$, if we so desire.

The multiattribute utility theory (MAUT) approach for solving the problem of selecting the k most preferred alternatives is to assess a utility function $u: V \rightarrow \mathbb{R}$ which has the property that value score vector v^m is preferred to value score vector v^l iff $u(v^m) > u(v^l)$. Alternatives can then be compared on the basis of expected utility; i.e., π' is preferred to π iff

$$\sum_{v \in V} u(v) \pi'(v) > \sum_{v \in V} u(v) \pi(v).$$

Since the expected utility generated by an alternative is a scalar, such a procedure linearly orders the alternatives and allows for the k most preferred alternatives to be easily identified.

Properly assessing the utility function, however, can be both time consuming and stressful for the decisionmaker. A choicemaking approach presented in (White and Sage, 1980), for the special case where each alternative is associated with a single outcome with probability one (i.e., the riskless case), has been developed to avoid, or reduce to the extent possible, the difficult and possibly unnecessary task of aggregating value scores. In this paper, we extend the approach presented in (White and Sage, 1980) to the case where outcomes of alternatives are described probabilistically in order to possibly reduce the time, effort, and stress often associated with MAUT utility assessment.

In order to put the results of this paper in perspective, it is useful to briefly describe the approach presented in (White and Sage, 1980). This approach is comprised of a general four step procedure for the riskless case:

1. Determine the set of alternatives that are nondominated with respect to the relation R .
2. Ask the DM (decision maker) to select the most preferred alternative from the nondominated set of alternatives.
3. If the DM can accomplish Step 2, remove the most preferred alternative from the alternative set. If the number of most preferred alternatives is less than k , go to Step 1; otherwise, stop.
4. If the DM cannot accomplish Step 2 due to the fact that the nondominated set is not small enough, then aggregate some but not necessarily all of the value scores. Typically, but not necessarily, several lower level objectives are aggregated into a single, higher level objective. Under proper assumptions, this aggregation procedure strengthens the relation R and hence will not increase the number of alternatives in the nondominated set. Go to Step 1.

The potential usefulness and behavioral relevance of the above approach are due to the facts that:

- a) The most preferred alternative will always be in the nondominated set as long as the utility function is restricted to be isotone on V .

- b) Alternative selection is generally easier if made from a subset of the alternative set than from the entire alternative set.
- c) Selection of the k most preferred alternatives may not require explicit aggregation of all the value scores, which is required by MAUT, and may, therefore, be accomplished with relatively less time, effort, and stress.

Of course, what constitutes a "small enough" nondominated set so that Step 2 can be accomplished is highly dependent on the DM, the contingency task structure, the scope or scale of the perceptions needed to aggregate value scores, the time allotted to the task, etc. We also remark that selection of the non-dominated alternative set should sometimes be accomplished along with a sensitivity study in order to insure that the DM considers all possible candidates for the most preferred alternatives.

The aggregation procedure mentioned in Step 4 is accomplished by determining an aggregation function $f: R^N \rightarrow R^L$ consistent with the objectives hierarchy and the DM's desires concerning aggregation of preferences. The aggregation function typically aggregates the value scores of two or more objectives into the value score of a single, higher level objective; i.e., $L=1$. See the automobile purchasing problem in (White and Sage, 1980) for an example.

For the case where each alternative is associated with a single outcome with probability one, the alternative set and the set of value score vectors can be viewed as equivalent and

hence the relation which describes preference on the value score vector set is inherited by the alternative set. Thus, for the certain case aggregating value scores to "strengthen" the relation on the value score vector set equivalently strengthens the relation on the alternative set. We will give a formal definition to a "strengthened" relation and discuss the implications of strengthening a relation beyond tending to reduce the associated nondominated set in the next section. When a one-to-one correspondence does not exist between the alternative set and the value score vector set, as is true for the problem considered here, the relations describing preference on these sets are different. It therefore does not necessarily follow that strengthening the relation on the value score vector set by properly aggregating value scores produces a stronger relation on the alternative set. If strengthening the value score vector set relation does strengthen the alternative set relation, then choice aiding for the risky case can proceed in a manner analogous to the procedure described in (White and Sage, 1980) and briefly outlined above.

In this paper we examine three relations on the alternative set which are induced by the relation on the value score vector set. The relations are associated with:

1. the expected value score vector
2. first order stochastic dominance
3. second order stochastic dominance

We show that for each of these three relations properly aggregating value scores strengthens the relation on the value score vector set which in turn induces a stronger relation on the alternative set. These three relations are considered in Sections 3, 4, and 5. Section 2 presents several preliminary results, and Section 6 summarizes the results and discusses their implications.

2. Preliminary Results and Comments

Assume ξ is a (preference-or-indifference) relation on an arbitrary set S , where $s' \xi s$ means s' is "weakly preferred" to s with respect to ξ . The strict preference relation induced by ξ is denoted by ξ_p ; i.e., $s' \xi_p s$ iff $s' \xi s$ and not $s \xi s'$. We do not require ξ to be complete; therefore, two elements of S can be related by any one of four possible relationships: $s' \xi s$ and not $s \xi s'$, $s \xi s'$ and not $s' \xi s$, $s' \xi s$ and $s \xi s'$, and not $s' \xi s$ and not $s \xi s'$ (i.e., s and s' are incomparable).

Let $\xi \subseteq \xi'$ be defined as: $s' \xi s$ implies $s' \xi' s$, and define $\xi_p \subseteq \xi'_p$ similarly. Thus, if s' is preferred to s with respect to ξ , then s' is also preferred to s with respect to ξ' , assuming $\xi \subseteq \xi'$. We say ξ' is stronger than (more properly, at least as strong as) ξ if $\xi \subseteq \xi'$ since ξ' is able to express preference between at least as many pairs of elements of S as can ξ . Several characterizations of the relationship between two relations ξ and ξ' on S , where $\xi \subseteq \xi'$, are given following preliminary definitions.

Let $\bar{S}(\xi)$ be the set of all nondominated elements in S with respect to ξ ; i.e., let

$$\bar{S}(\xi) = \{s \in S: \text{there does not exist an } s' \in S \text{ such that } s' \xi_p s\}$$

Define ξs as the set of all elements in S that are more preferred to $s \in S$ with respect to ξ ; i.e., $\xi s = \{s' \in S: s' \xi s\}$. Similarly, let $s \xi = \{s' \in S: s \xi s'\}$.

LEMMA 1: (a) Let $\xi \subseteq \xi'$. Then $\xi s \subseteq \xi' s$ and $s\xi \subseteq s\xi'$ for all $s \in S$. (b) Assume $\xi_p \subseteq \xi'_p$. Then, $\bar{S}(\xi'_p) \subseteq \bar{S}(\xi_p)$.

Proof: (a) The fact that $\xi s \subseteq \xi' s$ and $s\xi \subseteq s\xi'$ for all $s \in S$ follows directly from the definition of $\xi \subseteq \xi'$.

(b) To show that $\bar{S}(\xi'_p) \subseteq \bar{S}(\xi_p)$, we will show that $\bar{S}(\xi_p)^c \subseteq \bar{S}(\xi'_p)^c$, where c stands for complement. Let $s \in \bar{S}(\xi_p)^c$. Then, there is an $s' \in S$ such that $s' \xi_p s$ which due to $\xi_p \subseteq \xi'_p$ implies $s' \xi'_p s$. Thus, $s \in \bar{S}(\xi'_p)^c$, and the proof is complete. $\square$

The first part of Lemma 1 states that for each element in a set, strengthening the relation on the set allows more elements to be identified as being weakly more preferred to the given element. Analogously, for each element in the set, strengthening a relation allows more elements to be identified as being weakly less preferred to the given element. Equivalently, for each element in the set, strengthening a relation reduces the number of elements that are incomparable.

The second part of Lemma 1 shows that if the relations are preference relations, then the nondominated set for the stronger relation is a subset of the nondominated set for the weaker relation. It has been shown (see, for example, (White and Sage, 1980)) that the most preferred element of a set is in its non-dominated set if utility functions are assumed isotone. Thus, strengthening the preference relation on a set of alternatives may reduce the number of alternatives which are candidates for being the most preferred alternative. It is necessary, however, that the relation be a preference relation for the second part of Lemma 1 to hold, as

the following example demonstrates.

EXAMPLE 1: We now show that if either ξ and/or ξ' are not preference relations, then $\xi \subseteq \xi'$ may not imply that $\bar{S}(\xi') \subseteq \bar{S}(\xi)$. Let $S = (s^1, s^2) \in \mathbb{R}^2$, where $s^1 = (0,1)$ and $s^2 = (0,0)$.

Define:

- (i) $s' \xi s$ iff $s'_i \geq s_i$, $i=1,2$,
- (ii) $s' \xi' s$ iff $s'_1 \geq s_1$.

Note that $s' \xi s$ implies $s' \xi' s$, i.e. $\xi \subseteq \xi'$, and that neither relations are preference relations, e.g. it is possible that both $s' \xi s$ and $s \xi s'$. Note also that $s^1 = \bar{S}(\xi) \subseteq \bar{S}(\xi') = S$. □

We remark that the only way a nondominated set can increase after a relation has been strengthened is if:

1. There is an element s' in the nondominated set which is strictly more preferred to an element s on the basis of the weaker relation and
2. Strengthening the relation causes the new preference structure to be indifferent between s and s' .

Such a situation is unlikely to occur in most practical settings, and therefore its possibility should be of little operational concern. One exception to this statement is when an attribute(s) is dropped from consideration in order to strengthen the relation on the alternative set and there are ties in value scores associated with the other attributes, as demonstrated by Example 1.

For the choicemaking problem under risk being considered in this paper, the objective is to use a relation R on the set of alternatives Π to describe preference between alternatives in order to determine the nondominated alternatives, or equivalently, in

order to determine the alternatives that are candidates for being most preferred. If the nondominated set of alternatives is too large for choice selection, then we wish to strengthen R . Since R cannot be strengthened directly, we hope that aggregating value scores will cause R to become more discriminating. That is, if $R(R)$ is the relation on Π induced by the relation R on V , then we hope that aggregating value scores will strengthen R to R' , i.e. $R \subseteq R'$, which will in turn induce a stronger relation on the alternative set, i.e. $R(R) \subseteq R(R')$.

Throughout the remainder of the paper, the aggregation of value scores will be described functionally as $f: \mathbb{R}^N \rightarrow \mathbb{R}^L$, where usually but not necessarily $L < N$. Note that f induces a partial order on X : $x' R' x$ iff $f(x') \geq f(x)$. Throughout the remainder of the paper f will be assumed to be at least isotone, which is easily shown to imply $R \subseteq R'$.

3. Expected Value Score Dominance

We now consider the case where alternatives are compared on the basis of the expected value of their value score vectors. It is shown that aggregating value scores using a linear isotone aggregation function will strengthen this relation on the alternative set. We also show that a linear, strictly isotone aggregation will strengthen the associated preference relation, thus guaranteeing no alternatives will be added to the set of nondominated alternatives as the preference relation is strengthened.

Assume that:

- (i) $\pi' R_e \pi$ iff $\sum_{v \in V} v \pi'(v) \geq \sum_{v \in V} v \pi(v)$
- (ii) $\pi' P_e \pi$ iff $\pi'' R_e \pi$ and not $\pi R_e \pi'$.

Letting $f: \mathbb{R}^N \rightarrow \mathbb{R}^L$ be the aggregation function, we define:

- (i) $\pi' R'_e \pi$ iff $\sum_{v \in V} f(v) \pi'(v) \geq \sum_{v \in V} f(v) \pi(v)$
- (ii) $\pi' P'_e \pi$ iff $\pi' R'_e \pi$ and not $\pi R'_e \pi'$.

We refer to R_e as the expected value score relation on Π and P_e as its associated preference relation. R'_e and P'_e are the expected value score relation and preference relation, respectively, after a value score aggregation has been accomplished. Recall that a function is affine if it is linear plus a constant; i.e. $f(v) = Av + b$. Our main results for these relations are as follows.

LEMMA 2. (a) If f is affine and isotone, then $R_e \subseteq R'_e$.

(b) If f is affine and strictly isotone, then

$$\bar{\pi}(P'_e) \subseteq \bar{\pi}(P_e).$$

Proof: (a) Assume $\pi' R_e \pi$. By the isotonicity of f ,

$$f\left[\sum_{v \in V} v \pi'(v)\right] \geq f\left[\sum_{v \in V} v \pi(v)\right].$$

That f is affine then implies $\pi' R'_e \pi$.

(b) That $\pi' P_e \pi$ implies $\pi' P'_e \pi$ follows as above. The result then holds due to Lemma 1b. □

We remark that several decision aids for the single decision node under risk case assume that the aggregation function is linear, e.g. (Kelly, 1978). It is in general necessary to assume that the aggregation function is affine, as the following example demonstrates.

EXAMPLE 2. Let

$$v^1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad v^2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \text{and} \quad v^3 = \begin{bmatrix} 0.75 \\ 0.75 \end{bmatrix}$$

$$\pi'(v^1) = .2, \quad \pi'(v^2) = .2, \quad \pi'(v^3) = .6$$

$$\pi(v^1) = .3, \quad \pi(v^2) = .3, \quad \pi(v^3) = .4$$

We see that

$$\sum_{v \in V} v \pi'(v) = \begin{bmatrix} 0.65 \\ 0.65 \end{bmatrix}$$

$$\sum_{v \in V} v \pi(v) = \begin{bmatrix} 0.60 \\ 0.60 \end{bmatrix}$$

Thus, $\pi' R_e \pi$. Let $f(v_1, v_2) = \max\{v_1, v_2\}$, which is isotone.

Then,

$$\sum_{v \in V} f(v) \pi'(v) = 0.85$$

$$\sum_{v \in V} f(v) \pi(v) = 0.90$$

and hence $\pi R'_e \pi'$. Thus, $R \subseteq R'$ does not in general imply that $R_e \subseteq R'_e$ if f is isotone and nonlinear.

Consider also the case where the aggregation function is $f(v) = k_1 v_1 + k_2 v_2 + (1-k_1-k_2) v_1 v_2$, the multilinear case. A multilinear utility function is a necessary condition for mutual utility independence; see (Keeney and Raiffa, 1976) for details. Note that $f(v^1) = k_2$, $f(v^2) = k_1$, and $f(v^3) = 3(k_1 + k_2)/16 + 9/16$. It follows that

$$\sum_{v \in V} f(v) \pi'(v) = \frac{5}{16}(k_1 + k_2) + \frac{27}{80}$$

$$\sum_{v \in V} f(v) \pi(v) = \frac{3}{8}(k_1 + k_2) + \frac{9}{40}$$

Observe that if $(k_1 + k_2) \leq 9/5$, then $\pi' R'_e \pi$; however, when $(k_1 + k_2) \geq 9/5$, $\pi R'_e \pi'$. Thus, when the sum $k_1 + k_2$ becomes sufficiently large, $R \subseteq R'$ does not imply $R_e \subseteq R'_e$ but in fact implies that $R'_e \subseteq R_e$. □

Interestingly, if the number of outcomes equals 2, i.e. $M=2$, then the affine assumption in Lemma 2 can be deleted.

COROLLARY 1: Assume $M=2$. Then, if the aggregation function $f: R^N \rightarrow R^L$ is isotone, $R_e \subseteq R'_e$.

Proof: Note that

$$\sum_{v \in V} v \pi'(v) \geq \sum_{v \in V} v \pi(v)$$

is equivalent to

$$v^1 [\pi'(v^1) - \pi(v^1)] \geq v^2 [\pi'(v^1) - \pi(v^1)].$$

Assume $\pi'(v^1) - \pi(v^1) > 0$, which implies $v^1 \geq v^2$. Note that $v^1 \geq v^2$ implies $f(v^1) \geq f(v^2)$, which in turn implies

$$f(v^1) [\pi'(v^1) - \pi(v^1)] \geq f(v^2) [\pi'(v^1) - \pi(v^1)]$$

which is equivalent to the desired result. The $\pi'(v^1) - \pi(v^1) = 0$ case is trivial; the $\pi'(v^1) - \pi(v^1) < 0$ case proceeds as above. $\square$

The following example illustrates Lemma 2.

EXAMPLE 3. Assume there are five possible outcomes, four attributes initially under consideration, and six available alternatives, i.e. $M=5$, $N=4$, $P=6$. Table 1 presents assumed data, and Figure 2a displays the resulting domination digraph generated by the expected value scores listed in Table 2a. Figure 2a indicates that alternative 3 is dominated by alternatives 1, 2, and 4 and that the nondominated set of alternatives is $\{1, 2, 4, 5, 6\}$.

Assume $f(v) = Av$, where

$$A = \begin{bmatrix} .1 & .1 & .8 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Thus, the aggregation function is linear and the trade-off weights for the three attributes 1, 2, and 3 are .1, .1, and .8, respectively. Table 2b gives the expected value scores after the aggregation and Figure 2b displays the resulting digraph. Note that the aggregation has produced a nondominated set which is not larger (it is in fact smaller) than the nondominated set before the aggregation, which is in agreement with Lemma 2a. $\square$

		Outcome Number				
		1	2	3	4	5
Attribute Number	1	10	5	5	0	5
	2	10	0	0	0	0
	3	3	3	10	0	3
	4	5	5	5	0	10

(a) Value Scores for Each Outcome

		Outcome Number				
		1	2	3	4	5
Alternative Number	1	0.6	0.1	0.2	0.1	0.0
	2	0.7	0.0	0.1	0.2	0.0
	3	0.3	0.1	0.0	0.4	0.2
	4	0.3	0.0	0.1	0.1	0.5
	5	0.1	0.1	0.0	0.1	0.7
	6	0.0	0.1	0.1	0.0	0.8

(b) Probabilities for Example 3.

TABLE 1. Data for Example 3.


		Alternative Numbers					
		1	2	3	4	5	6
Attribute Number	1	7.5	7.5	4.5	6.0	5.0	5.0
	2	6.0	7.0	3.0	3.0	1.0	0.0
	3	4.1	3.1	1.8	3.4	2.7	3.7
	4	4.5	4.0	4.0	7.0	8.0	9.0

(a) Expected Value Scores Before the Aggregation


		Alternative Numbers					
		1	2	3	4	5	6
Attribute Number	1-2-3	4.63	3.93	2.19	3.62	2.76	3.46
	4	4.50	4.00	4.00	7.00	8.00	9.00

(b) Expected Value Scores After the Aggregation

TABLE 2. Expected Value Scores for Example 3.



(a)



(b)

FIGURE 2: Domination Diagrams Based on Expected Value Score Dominance for Example 3: (a) Before Attribute Aggregation, (b) After Attribute Aggregation.

We remark that the well-known mean-variance order (a definition is given, for example, in (Fishburn, and Vickson, 1978)) is a special case of the expected value score relation. To see this, let $Q=1$ and $N=2$, interpret $x(\omega_m)$ as the return resulting from outcome m , and define

$$v_1[x(\omega_m)] = x(\omega_m)$$

$$v_2[x(\omega_m)] = -(x(\omega_m) - \bar{x}_p)^2$$

if $\pi_p(m) \neq 0$, where $\bar{x}_p = \sum_m x(\omega_m) \pi_p(m)$. Alternatively, $v_1[x(\omega_m)] = \bar{x}_p$. In order for $v = \{v_1, v_2\}$ to be well-defined, we require that Ω and Π are modeled to have the following property: if m is such that $\pi_p(m) \neq 0$, then $\pi_q(m) = 0$ for all $q \in \Pi$, $q \neq p$. The relation R_e then becomes the mean-variance order.

4. First-Order Stochastic Dominance

In this section, we investigate multiattribute, first-order stochastic dominance as a means of defining a relation on the alternative set Π . We demonstrate that an isotone aggregation of value scores on the value score vector set V strengthens the relation on Π . An additional condition is required to prove that the preference relation on Π is strengthened.

Consider the following definitions:

- (i) $K = \{K \subseteq V: v \in K, v' \in V, \text{ and } v' \geq v \text{ imply } v' \in K\}$
- (ii) $\pi' R_1 \pi$ iff $\pi'(K) \geq \pi(K)$ for all $K \in \mathcal{K}$
- (iii) $\pi' P_1 \pi$ iff $\pi' R_1 \pi$ and not $\pi R_1 \pi'$.

Observe that (iii) is equivalent to:

- (iii)' $\pi' P_1 \pi$ iff $\pi' R_1 \pi$ and there is a $K \in \mathcal{K}$ such that $\pi'(K) \neq \pi(K)$.

We refer to R_1 as the first order stochastic dominance relation on Π and P_1 as its associated preference relation. Observe that $\mathcal{K}$ represents the collection of all the so-called increasing subsets of V with respect to the usual partial order R on $\mathbb{R}^N$. As has been stated in (Fishburn and Vickson, 1978), R_1 partially orders Π ; that is, R_1 is both transitive and antisymmetric.

Note that by extending the domain of the elements in Π to be the usual collection of Borel sets in $\mathbb{R}^N$, it is easy to show $\pi' R_1 \pi$ is equivalent to:

$$\int_S d\pi' \geq \int_S d\pi$$

for all increasing Borel sets S . As shown in (Lehmann, 1955) and (Fishburn and Vickson, 1978), such a statement is equivalent to the usual definition of first-order stochastic dominance:

$$\sum_{v \in V} u(v) \pi'(v) \geq \sum_{v \in V} u(v) \pi(v)$$

for all $u \in U_1$, where U_1 is the collection of all isotone scalar-valued functions on R^N . Thus, if $\pi' R_1 \pi$, then alternative π' can never produce a lower expected utility value than alternative π , as long as we restrict attention to isotone utility functions.

We remark that a particularly desirable aspect of the definition of the relations R_1 and P_1 is that K contains a finite number of elements (in fact, at most $\sum_{m=1}^M \binom{M}{m}$). Therefore, a finite algorithm can easily be constructed that can test for first-order stochastic dominance.

Let $f: R^N \rightarrow R^L$ be the aggregation function, and let $K' = \{K \subseteq V: v \in K, v' \in V, \text{ and } f(v') \geq f(v) \text{ imply } v' \in K\}$. Define R'_1 and P'_1 exactly as R_1 and P_1 were defined except substitute K' for K . We now present the main result of this section.

LEMMA 3. If f is isotone, then $R_1 \subseteq R'_1$.

Proof: A simple argument demonstrates that $K' \subseteq K$, which implies $R_1 \subseteq R'_1$ directly from the definition. □

Determining conditions which imply that $P_1 \subseteq P'_1$ is a more complicated task than determining conditions which imply that $R_1 \subseteq R'_1$ and requires further definitions. Let $K_1 = \{K \in K: K^C \in K\}$, $K_2 = K_1^C$, and define K'_i , $i=1, 2$, similarly with K' replacing K . Define $\tilde{K}(\pi', \pi) = \{K \in K_2: \pi'(K) \neq \pi(K)\}$; observe that if $\pi' P_1 \pi$, then $\tilde{K}(\pi', \pi) \neq \emptyset$.

LEMMA 4: Assume:

- (i) f is isotone
- (ii) $K_2' \cap \tilde{K}(\pi', \pi) \neq \emptyset$ for all $\pi', \pi \in \Pi$
such that $\pi' P_1 \pi$.

Then, $\bar{\Pi}(P_1') \subseteq \bar{\Pi}(P_1)$.

Proof: Lemma 3 implies $R_1 \subseteq R_1'$. It is easily shown that $K_2' \cap \tilde{K}(\pi', \pi) \neq \emptyset$ insures that $\pi' P_1 \pi$ implies $\pi' P_1' \pi$. Condition (ii) insures that this is true for all pairs in Π of interest and hence that $P_1 \subseteq P_1'$. The result then follows from Lemma 1b. □


We remark that condition (ii) in Lemma 4 is not very meaningful operationally since K_2' becomes known only after the value score aggregation. However, as we have discussed earlier, the desired characteristic, $\bar{\Pi}(P_1') \subseteq \bar{\Pi}(P_1)$, will probably rarely be affected by whether or not condition (ii) holds.

It is interesting to note that $\pi'(K) = \pi(K)$ for all $K \in K_1$ if $\pi' R_1 \pi$. This fact suggests that in trying to determine if $\pi' R_1 \pi$, a simple initial check would be to see if $\pi'(K) = \pi(K)$ for all $K \in K_1$. We observe that only half of the K_1 sets need to be checked since $\pi'(K) = \pi(K)$ iff $\pi'(K^C) = \pi(K^C)$.


We also remark that if all elements in the outcome set are nondominated, then $K = K_1$, and conversely. The discussion

preliminary to the statement of Lemma 4 can then be used to show that $\bar{\pi}(P_1) = \Pi$.

EXAMPLE 4. Consider the problem presented in Example 3. The data presented in Table 1a generate the digraph of elements in V in Figure 3a. The increasing sets in K associated with this digraph which are of interest are: $\{1\}$, $\{3\}$, $\{5\}$, $\{1,3\}$, $\{1,5\}$, $\{3,5\}$, $\{1,3,5\}$, and $\{1,2,3,5\}$. (We delete consideration of $\{1,2,3,4,5\} \in K$ and $\phi \in K$ since $\pi(\{1,2,3,4,5\}) = 1.0$ and $\pi(\phi) = 0.0$ for all $\pi \in \Pi$.) The probabilities associated with each of these sets for each alternative are displayed in Table 3a; Figure 3b presents the related digraph of elements in Π . As in Example 3, let f be linear and assume value scores 1, 2, and 3 are traded-off with weights .1, .1, and .8, respectively. The resulting digraph of elements in V is given in Figure 4a. This digraph has the following increasing sets of interest: $\{3\}$, $\{5\}$, $\{3, 5\}$, $\{1, 3\}$, $\{1, 3, 5\}$, and $\{1,2,3,5\}$. Notice that the total number of increasing sets has been reduced by the aggregation. The digraph of elements in Π associated with the data presented in Table 3b is displayed in Figure 4b, providing more preference information with respect to the alternatives than does the digraph in Figure 3b. □



(a)



(b)

FIGURE 3. Domination Digraphs Associated with First-Order Stochastic Dominance for Example 4 Before the Value Score Aggregation (a) The Value Score Digraph (b) The Alternative Digraph


Increasing Subset	Alternative					
	1	2	3	4	5	6
{1}	0.6	0.7	0.3	0.3	0.1	0.0
{3}	0.2	0.1	0.0	0.1	0.0	0.1
{5}	0.0	0.0	0.2	0.5	0.7	0.8
{1, 3}	0.8	0.8	0.3	0.4	0.1	0.1
{1, 5}	0.6	0.7	0.5	0.8	0.8	0.8
{3, 5}	0.2	0.1	0.2	0.6	0.7	0.9
{1, 3, 5}	0.8	0.8	0.5	0.9	0.8	0.9
{1, 2, 3, 5}	0.9	0.8	0.6	0.9	0.9	1.0

(a)


{3}	0.2	0.1	0.0	0.1	0.0	0.1
{5}	0.0	0.0	0.2	0.5	0.7	0.8
{3, 5}	0.2	0.1	0.2	0.5	0.7	0.8
{1, 3}	0.8	0.8	0.3	0.4	0.1	0.1
{1, 3, 5}	0.8	0.8	0.5	0.9	0.8	0.9
{1, 2, 3, 5}	0.9	0.8	0.6	0.9	0.9	0.0

(b)

TABLE 3: Probabilities of the Increasing Sets for Each Alternative, i.e. $P(v \in K \mid \pi)$ for each $K \in \mathcal{K}$ and $\pi \in \Pi$ (a) Before and (b) After the Value Score Aggregation in Example 4.



(a)



(b)

FIGURE 4: Domination Digraphs Associated with First Order Stochastic Dominance for Example 4 after the Value Score Aggregation
 (a) The Value Score Digraph (b) The Alternative Digraph.

It is of interest to give conditions that relate the two orders R_e and R_1 . Let the function $I_K: V \rightarrow \{0,1\}$ be such that $I_K(v) = 1$ if $v \in K$ and $I_K(v) = 0$ if $v \notin K$. Let $I = \{I_K\}_{K \in \mathcal{K}}$. Thus, $I(v)$ is a γ -vector if K contains γ subsets of V which identifies the increasing sets containing v . It is easy to demonstrate that $\pi' R_1^I \pi$ is equivalent to

$$\sum_V I(v) \pi'(v) \geq \sum_V I(v) \pi(v)$$

It is straightforward to show that the following results hold using the same procedure as was used in the proof of Lemma 2.

LEMMA 5. (a) Assume there is an isotone, affine function α such that $\alpha[I(v)] = v$ for all $v \in V$. Then $R_1 \subseteq R_e$.

(b) Assume there is an isotone, affine function β such that $\beta(v) = I(v)$ for all $v \in V$. Then $R_e \subseteq R_1$.

We now illustrate Lemma 5 with the following example.

EXAMPLE 5. Consider the data presented in Tables 1a and 3a for Examples 3 and 4, respectively. With reference to Table 3a, note that $I(v^1) = \text{col } \{1,0,0,1,1,0,1,1\}$, in that $1 \in \{1\}$, $1 \notin \{3\}$, $1 \notin \{5\}$, $1 \in \{1,3\}$, and so forth. The collection of column vectors $\{I(v^1), \dots, I(v^5)\}$ is then:

$$[I(v^1), \dots, I(v^5)] = \begin{bmatrix} 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 & 1 \end{bmatrix}$$

In order to illustrate Lemma 5a, we seek an isotone, affine function α such that

$$[v^1, \dots, v^5] = \alpha[I(v^1), \dots, I(v^5)]$$

where from Table 1a,

$$[v^1, \dots, v^5] = \begin{bmatrix} 10 & 5 & 5 & 0 & 5 \\ 10 & 0 & 0 & 0 & 0 \\ 3 & 3 & 10 & 0 & 3 \\ 5 & 5 & 5 & 0 & 10 \end{bmatrix}$$

Note that since $v^4 = \alpha I(v^4)$, α is required to be linear; i.e. α is a 4×8 matrix. We can now omit v^4 and $I(v^4)$ from further consideration. Standard algebraic procedures show that

$$\alpha = \begin{bmatrix} 5 & 0 & 0 & 0 & 0 & 0 & 0 & 5 \\ 10 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 7 & 0 & 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 5 & 0 & 0 & 0 & 0 & 5 \end{bmatrix}$$

Clearly, α is isotone and hence from Lemma 5a, $R_1 \subseteq R_e$, which is verified by Figures 2a and 3b. Since the aggregation function used was linear in both Examples 3 and 4, it is straightforward to show that $R_1' \subseteq R_e'$, using Lemma 5a and the same α as above.

Straightforward algebraic manipulation shows that there exists a unique 8×4 matrix β such that $\beta v = I(v)$ for all $v \in V$, where

$$\beta = \begin{bmatrix} 0 & 1/10 & 0 & 0 \\ -3/35 & 3/70 & 1/7 & 0 \\ -1/5 & 1/10 & 0 & 1/5 \\ -3/35 & 1/7 & 1/7 & 0 \\ -1/5 & 1/5 & 0 & 1/5 \\ -2/7 & 1/7 & 1/7 & 1/5 \\ -2/7 & 17/70 & 1/7 & 1/5 \\ 1/5 & -1/10 & 0 & 0 \end{bmatrix}$$

Note, however, that β is not isotone and thus it does not necessarily follow from Lemma 5b that $R_e \subseteq R_1$. In fact, we observe from Figures 2a and 3b that $R_e \subseteq R_1$ does not hold. □

5. Second-Order Stochastic Dominance

In this section, we examine multiattribute second-order stochastic dominance as a means of ordering the set of alternatives. Using necessary and sufficient conditions for multivariable second order stochastic dominance due to Fishburn and Vickson (1978), we show that a linear, isotone aggregation of value scores on the set V induces a strengthened relation on the alternative set Π . Here we will not define a strict preference relation on Π , noting that a stronger relation on Π will usually produce a nondominated set of alternatives after a value score aggregation which is no larger than the nondominated set of alternatives before the aggregation.

Define U_2 as the collection of all isotone, concave functions; i.e., $u \in U_1$ and $u[\lambda r + (1-\lambda)r'] \geq \lambda u(r) + (1-\lambda)u(r')$ for all $0 \leq \lambda \leq 1$ for any $r, r' \in R^N$. The concavity of a utility function is the functional representation of global risk aversion, describing the preference structure of a decision-maker who will never invest in an actuarially fair prospect.

Consider the following definition:

$$\pi' R_2 \pi \text{ iff } \sum_{v \in V} u(v) \pi'(v) \geq \sum_{v \in V} u(v) \pi(v)$$

for all $u \in U_2$.

We refer to R_2 as the second order stochastic dominance relation on Π .

Thus, if we know the decisionmaker is risk averse, then

$\pi' R_2 \pi$ implies that alternative π' is at least as good as alternative π (with respect to the expected utility criterion).

Clearly, it follows directly from the definitions that $R_1 \subseteq R_2$ and that R_2 is a partial order.

Fishburn and Vickson (1978) have presented a slightly less general version of the following necessary and sufficient conditions for $\pi'R_2\pi$: there exists a feasible solution to the set of a linear equalities and inequalities:

$$(i) \quad d_{ij} \geq 0 \text{ for all } i, j=1, \dots, M$$

$$(ii) \quad \sum_{j=1}^M d_{ij} = 1 \text{ for all } i = 1, \dots, M$$

$$\text{such that } \pi'(v^i) \neq 0$$

$$(iii) \quad \pi(v^j) = \sum_{i=1}^M \pi'(v^i) d_{ij} \text{ for all } j=1, \dots, M$$

$$(iv) \quad \sum_{i=1}^M d_{ij} v_n^j \leq v_n^i \text{ for all } i=1, \dots, M$$

such that $\pi'(v^i) \neq 0$ and all $n=1, \dots, N$,
 where v_n^j is the n^{th} scalar entry of $v^j \in V$;
 i.e. $v^j = \{v_1^j, \dots, v_N^j\}$.

Interpretations of the solution $\{d_{ij}\}$ to the above can be found in (Fishburn and Vickson, 1978).

Let $f: \mathbb{R}^N \rightarrow \mathbb{R}^L$ be the aggregation function, and define R_2' as R_2 was defined, except replace U_2 with U_2' , where

$$U_1' = \{u: u \text{ is isotone on } f(\mathbb{R}^N)\}$$

$$U_2' = \{u \in U_1': u \text{ is concave on } f(\mathbb{R}^N)\}.$$

We now present conditions on f which insure $R_2 \subseteq R_2'$.

LEMMA . Assume $f: \mathbb{R}^N \rightarrow \mathbb{R}^L$ satisfies the following conditions:

$$(1) \quad f_\ell(v^j) = \sum_{n=1}^N v_n^j b_{n\ell} \text{ where } b_{n\ell} \geq 0$$

for all $n = 1, \dots, N$ and $\ell = 1, \dots, L$

$$(2) \quad \text{if } v^i \neq v^j, \text{ then } f(v^i) \neq f(v^j).$$

Then, $R_2 \subseteq R_2'$.

Proof. We wish to show that if (iv) holds, then

$$\sum_{j=1}^M d_{ij} f_{\ell}(v^j) \leq f_{\ell}(v^i)$$

for all $i = 1, \dots, M$ such that $\pi'(v^i) \neq 0$ and all $\ell = 1, \dots, L$.

(2) implies that the number of v^i such that $\pi'(v^i) \neq 0$ does not change after the aggregation is performed. (1) can be used to easily show that (iv) implies the above inequality. $\square$

EXAMPLE 6. Again consider the problem presented in Example 3, using second-order stochastic dominance as the method of generating a partial order on Π . Figure 5 is the dominance digraph of elements in Π before the value score aggregation. Figure 2b is, coincidentally, the dominance digraph of elements in Π after the attribute aggregation. Observe that the assumption of risk aversion has caused the relationship $R_1 \subseteq R_2$ to be strict both before the aggregation (compare Figures 3b and 5) and after the aggregation (compare Figures 2b and 4b). $\square$

It seems reasonable to pose the following computationally attractive conjecture: assume, for each $n=1, \dots, N$, that π' stochastically dominates π in second order based on the outcome set $\{v_n^1, \dots, v_n^M\}$; that is, assume for each $n=1, \dots, N$, that (i) through (iii) hold and

$$(iv)' \quad \sum_{i=1}^M d_{ij} v_n^j \leq v_n^i$$

for all $i=1, \dots, M$ such that $\pi'(v^i) \neq 0$. Then π' stochastically




FIGURE 5. The Digraph for Example 6 Before the Value Score Aggregation

dominates π in second order based on the outcome set $\{v^1, \dots, v^M\}$; that is, (i) through (iv) hold.

The computational attractiveness of the above conjecture is due to the fact that there exist computationally simple checks for univariate second order stochastic dominance, c.f. (Fishburn and Vickson, 1978). Unfortunately, the above conjecture is false in general since the set of all $\{d_{ij}\}$ satisfying (iv)' for $n=k$ and (iv)' for $n=k'$, $k \neq k'$, may be null. It has been shown that the above statement and its converse hold, c.f. Theorem 2.11 (Fishburn and Vickson, 1978), under suitable independence assumptions. These independence assumptions, however, do not imply that the sufficient conditions for $R_2 \subseteq R_2'$ presented in Lemma 6 can be relaxed.

It is important to observe that the expected value score relation and the stochastic dominance relations differ fundamentally in their interpretation of a complete value score aggregation, i.e. for the case where the aggregation function $f: R^N \rightarrow R^1$ (the $L=1$ case). In this case, R_e' is a linear order and R_1' and R_2' may only be partial orders. R_e' essentially treats $f(v)$ as a utility function, i.e., $u(x) = f[v(x)]$, whereas R_1' and R_2' assume there exists a still unassessed function U which when composed with $f(v)$ produces the desired utility function, i.e., $u(x) = U[f[v(x)]]$. A discussion of the usefulness of assuming the existence of such a function U , especially for modeling risk, can be found in (Bodily, 1980). An example is now presented to illuminate the

fact that if there is a single attribute, presumably associated with the highest level objective and a complete value score aggregation, then the expected value score relation is linearly ordered but the stochastic dominance relations may not be.

EXAMPLE 7. Consider the problem presented in Example 3, and let the aggregation function $f(v) = Av$, where

$$A = [.05 \quad .05 \quad .40 \quad .50].$$


Thus, f linearly aggregates the value scores into a single, scalar value score with weights .05, .05, .40, and .50.

Equivalently, f linearly aggregates the value scores for attributes 1-2-3 and 4 with a weight of .5 each, c.f. Table 2b.


Figure 6 presents the relevant digraphs. Note that $R_1 \subseteq R_e$, in agreement with Example 5, and that $R_1 \subseteq R_2$.



(a)



(b)



(c)

FIGURE 6. Domination Digraphs for Example 7. (a) Expected Value Score, (b) First Order Stochastic Dominance, (c) Second Order Stochastic Dominance.

6. Conclusions

In this paper, we have examined a multiattribute, single decision node decision making problem under risk. The intent has been to investigate the extension of an approach to decision aiding under certainty presented in (White and Sage, 1980) to the case where outcomes are described probabilistically. We have seen that the risky case differs fundamentally from the certain case in that the relation on the alternative set is induced, at least in the three relations studied, by the relation on the value score vector set rather than being essentially equivalent to it. This fact has lead us to examine whether or not additional preference information supplied by the decision-maker in the form of value score aggregations would be passed on to the relation on the alternative set in the form of a stronger partial ordering. Under suitable conditions for the three induced partial orders examined, this "preference transmittal" has been shown to exist.

We observe that the value score partial order is significantly less computationally demanding than the partial orders generated by the stochastic dominance notions. Use of the value score relation requires determining and comparing the weighted sum of P N -vectors. The first-order stochastic dominance relation requires comparing P vectors having up to $\sum_{m=1}^M \binom{M}{m}$ scalar elements, where each scalar element is the sum of probabilities and where $\sum_{m=1}^M \binom{M}{m}$ represents the maximum number of elements in K . The second-

order stochastic dominance relation requires determining the feasibility of $P(P-1)$ linear programs, each having up to $M(M+N+2)$ decision variables ($2M$ of which are artificial variables) and up to $M(N+2)$ side constraints. Of course, these computational requirements may decrease if Lemma 5 can be used and/or if the fact that $R_1 \subseteq R_2$ is used. Clearly, the second-order stochastic dominance relation can be substantially more computationally demanding than either of the other two relations. The computational effort required to determine the dominance digraph for the first-order stochastic dominance relation and the value score relation is similar, provided that N and the number of increasing sets in K do not differ significantly and that K has been determined. Determining all of the increasing subsets (an algorithm for determining K is given in (White and Sage, 1980)), however, is significantly more computationally involved than the task of determining the domination digraph generated by the first-order stochastic dominance relation once K has been determined.

We remark that it is possible to model the mean variance order in the context of the expected value score relation without requiring Ω and Π to satisfy the property: $\pi_p(n) \neq 0$ implies $\pi_q(m) = 0$ for all $q \in \Pi$, $q \neq p$ for all m (see Section 3). To see this, let the value score vector depend on both the alternative and outcome; that is let $v_n = v_n[x(\omega_m), p]$. Assume $Q=1$ and $N=2$, and interpret $x(\omega_m)$ to be the return resulting from outcome m . We now define:

$$v_1[x(\omega_m), p] = x(\omega_m)$$

$$v_2[x(\omega_m), p] = -(x(\omega_m) - \bar{x}_p)^2.$$

Alternatively, define

$$v_1[x(\omega_m), p] = \bar{x}_p.$$

The resulting relation R_e is then the mean-variance order. It is straightforward to show that the results in Lemmas 2 and 3 easily generalize to the case where the value score vector can depend on both alternative and outcome. Extending the results in Sections 4 and 5 so that the value score vector can depend on the selected alternative is not so straightforward (e.g. note the obvious difficulties in defining R_1) and is a topic of future research.

REFERENCES:

Bodily, S. E., "Stability, Equality, Balance, and Multivariate Risk", Darden School Working Paper, Univ. of Virginia, Charlottesville, VA, August, 1980.

Fishburn, P. C., and Vickson, R. G., "Theoretical Foundations of Stochastic Dominance," Chapter 2 in Whitmore, G. A., and Findlay, M. C. (eds), Stochastic Dominance: An Approach to Decision Making Under Risk, Heath, Lexington, Ma, 1978.

Keeney, R. L., and Raiffa, H., Decisions with Multiple Objectives: Preferences and Value Tradeoffs, Wiley, NY, NY, 1976.

Kelly, C. W., "Decision Aids: Engineering Science and Clinical Art," Technical Report, Decisions and Designs, Inc., McLean, VA., 1978.

Lehmann, E. L., "Ordered Families of Distributions", Annals of Math. Statistics, 26, pp. 399-419, 1955.

White, C. C., and Sage, A. P., "A Multiple Objective Optimization-Based Approach to Choicemaking," IEEE Trans. on Systems, Man, and Cybernetics, SMC-10, pp. 315-326, 1980.

A.

DISTRIBUTION LIST

OSD

CDR Paul R. Chatelier
Office of the Deputy Under Secretary
of Defense
OUSDRE (E&LS)
Pentagon, Room 3D129
Washington, D.C. 20301

Department of the Navy

Director
Engineering Psychology Programs
Code 455
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217 (5 cys)

Director
Operations Research Programs
Code 434
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217

Director
Statistics and Probability Program
Code 436
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217

Director
Information Systems Program
Code 437
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217

Code 430B
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217

LCDR W. Moroney
Code 55MP
Naval Postgraduate School
Monterey, CA 93940

Department of the Navy

Commanding Officer
ONR Eastern/Central Regional Office
ATTN: Dr. J. Lester
Building 114, Section D
666 Summer Street
Boston, MA 02210

Commanding Officer
ONR Branch Office
ATTN: Dr. C. Davis
536 South Clark Street
Chicago, IL 60605

Commanding Officer
ONR Western Regional Office
ATTN: Dr. E. Gloye
1030 East Green Street
Pasadena, CA 91106

Office of Naval Research
Scientific Liaison Group
American Embassy, Room A-407
APO San Francisco, CA 96503

Director
Naval Research Laboratory
Technical Information Division
Code 2627
Washington, D.C. 20375 (6 cys)

Dr. Bruce Wald
Communications Sciences Division
Code 7500
Naval Research Laboratory
Washington, D.C. 20375

Dr. Robert G. Smith
Office of the Chief of Naval
Operations, OP987H
Personnel Logistics Plans
Washington, D.C. 20350

Naval Training Equipment Center
ATTN: Technical Library
Orlando, FL 32813

Department of the Navy

Human Factors Department
Code N215
Naval Training Equipment Center
Orlando, FL 32813

Dr. Alfred F. Smode
Training Analysis and Evaluation
Group
Naval Training Equipment Center
Code N-00T
Orlando, FL 32813

Dr. George Moeller
Human Factors Engineering Branch
Submarine Medical Research Lab
Naval Submarine Base
Groton, CT 06340

Dr. James McGrath, Code 302
Navy Personnel Research and
Development Center
San Diego, CA 92152

Navy Personnel Research and
Development Center
Planning and Appraisal
Code 04
San Diego, CA 92152

Navy Personnel Research and
Development Center
Management Systems, Code 303
San Diego, CA 92152

Navy Personnel Research and
Development Center
Performance Measurement and
Enhancement
Code 309
San Diego, CA 92152

CDR P. M. Curran
Code 604
Human Factors Engineering Division
Naval Air Development Center
Warminster, PA 18974

Dean of the Academic Departments
U.S. Naval Academy
Annapolis, MD 21402

Department of the Navy

Dr. Gary Poock
Operations Research Department
Naval Postgraduate School
Monterey, CA 93940

Dean of Research Administration
Naval Postgraduate School
Monterey, CA 93940

Mr. Warren Lewis
Human Engineering Branch
Code 8231
Naval Ocean Systems Center
San Diego, CA 92152

Dr. A. L. Slafkosky
Scientific Advisor
Commandant of the Marine Corps
Code RD-1
Washington, D.C. 20380

Department of the Army

Mr. J. Barber
HQs, Department of the Army
DAPE-MBR
Washington, D.C. 20310

Dr. Joseph Zeidner
Technical Director
U.S. Army research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Director, Organizations and
Systems Research Laboratory
U.S. Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Department of the Air Force

U.S. Air Force Office of Scientific
Research
Life Sciences Directorate, NL
Bolling Air Force Base
Washington, D.C. 20332

Department of the Air Force

Dr. Donald A. Topmiller
Chief, Systems Engineering Branch
Human Engineering Division
USAF AMRL/HES
Wright-Patterson AFB, OH 45433

Air University Library
Maxwell Air Force Base, AL 36112

Dr. Erl Alluuisi
Chief Scientist
AFHRL/CCN
Brooks AFB, TX 78235

Foreign Addresses

North East London Polytechnic
The Charles Myers Library
Livingstone Road
Stratford
London E15 2LJ
ENGLAND

Professor Dr. Carl Graf Hoyos
Institute for Psychology
Technical University
8000 Munich
Arcisstr 21
FEDERAL REPUBLIC OF GERMANY

Dr. Kenneth Gardner
Applied Psychology Unit
Admiralty Marine Technology
Establishment
Teddington, Middlesex TW11 OLN
ENGLAND

Director, Human Factors Wing
Defence & Civil Institute of
Environmental Medicine
Post Office Box 2000
Downsview, Ontario M3M 3B9
CANADA

Dr. A. D. Baddeley
Director, Applied Psychology Unit
Medical Research Council
15 Chaucer Road
Cambridge, CB2 2EF, ENGLAND

Other Government Agencies

Defense Technical Information Center
Cameron Station, Bldg. 5
Alexandria, VA 22314 (12 cys)

Dr. Craig Fields
Director, Cybernetics Technology
Office
Defense Advanced Research Projects
Agency
1400 Wilson Blvd
Arlington, VA 22209

Dr. Judith Daly
Cybernetics Technology Office
Defense Advanced Research Projects
Agency
1400 Wilson Blvd
Arlington, VA 22209

Other Organizations

Dr. Gary McClelland
Institute of Behavioral Sciences
University of Colorado
Boulder, CO 80309

Dr. Miley Merkhofer
Stanford Research Institute
Decision Analysis Group
Menlo Park, CA 94025

Dr. Jesse Orlansky
Institute for Defense Analyses
400 Army-Navy Drive
Arlington, VA 22202

Professor Judea Pearl
Engineering Systems Department
University of California-Los Angeles
405 Hilgard Avenue
Los Angeles, CA 90024

Professor Howard Raiffa
Graduate School of Business
Administration
Harvard University
Soldiers Field Road
Boston, MA 02163

Other Organizations

Dr. Arthur I. Siegel
Applied Psychological Services, Inc.
404 East Lancaster Street
Wayne, PA 19087

Dr. Paul Slovic
Decision Research
1201 Oak Street
Eugene, OR 97401

Dr. Amos Tversky
Department of Psychology
Stanford University
Stanford, CA 94305

Dr. Robert T. Hennessy
NAS - National Research Council
JH #819
2101 Constitution Avenue, N.W.
Washington, D.C. 20418

Dr. M. G. Samet
Perceptrics, Inc.
6271 Variel Avenue
Woodland Hills, CA 91364

Dr. Meredith P. Crawford
American Psychological Association
Office of Educational Affairs
1200 17th Street, N.W.
Washington, D.C. 20036

Dr. Ward Edwards
Director, Social Science Research
Institute
University of Southern California
Los Angeles, CA 90007

Dr. Charles Gettys
Department of Psychology
University of Oklahoma
455 West Lindsey
Norman, OK 73069

Dr. Kenneth Hammond
Institute of Behavioral Science
University of Colorado
Room 201
Boulder, CO 80309

Other Organizations

Dr. William Howell
Department of Psychology
Rice University
Houston, TX 77001

Journal Supplement Abstract Service
American Psychological Association
1200 17th Street, N.W.
Washington, D.C. 20036 (3 cys)

Dr. John Payne
Duke University
Graduate School of Business
Administration
Durham, NC 27706

Dr. Baruch Fischhoff
Decision Research
1201 Oak Street
Eugene, OR 97401

Dr. Leonard Adelman
Decisions and Designs, Inc.
8400 Westpark Drive, Suite 600
P. O. Box 907
McLean, VA 22101

Dr. Lola Lopes
Department of Psychology
University of Wisconsin
Madison, WI 53706

UNIVERSITY OF VIRGINIA

School of Engineering and Applied Science

The University of Virginia's School of Engineering and Applied Science has an undergraduate enrollment of approximately 1,400 students with a graduate enrollment of approximately 600. There are 125 faculty members, a majority of whom conduct research in addition to teaching.

Research is an integral part of the educational program and interests parallel academic specialties. These range from the classical engineering departments of Chemical, Civil, Electrical, and Mechanical and Aerospace to departments of Biomedical Engineering, Engineering Science and Systems, Materials Science, Nuclear Engineering and Engineering Physics, and Applied Mathematics and Computer Science. In addition to these departments, there are interdepartmental groups in the areas of Automatic Controls and Applied Mechanics. All departments offer the doctorate; the Biomedical and Materials Science Departments grant only graduate degrees.

The School of Engineering and Applied Science is an integral part of the University (approximately 1,530 full-time faculty with a total enrollment of about 16,000 full-time students), which also has professional schools of Architecture, Law, Medicine, Commerce, Business Administration, and Education. In addition, the College of Arts and Sciences houses departments of Mathematics, Physics, Chemistry and others relevant to the engineering research program. This University community provides opportunities for interdisciplinary work in pursuit of the basic goals of education, research, and public service.

DATE
FILMED
-8