


LEVEL *III*

(12)



AD A O 817 32

See 14/73

Distances of Random Variables and Point Processes

by

D.J. Daley

**Department of Statistics (I.A.S.)
Australian National University
Canberra 2600 Australia**

Institute of Statistics Mimeo Series #1331

March 1981

**DEPARTMENT OF STATISTICS
Chapel Hill, North Carolina**

DISTRIBUTION STATEMENT A
Approved for public release.

OTIC FILE COPY

81 4 10 034

12

DISTANCES OF RANDOM VARIABLES AND POINT PROCESSES

by

D.J. Daley
Department of Statistics (I.A.S.)
Australian National University
Canberra 2600 Australia

DTIC
SELECTED
APR 10 1981
C

ABSTRACT

The paper is basically a discussion and survey of some literature concerning the concept of a distance function for random variables and point processes. Work of Kalzanov on the closeness of renewal processes to Poisson processes is outlined, and there are comments on various possible comparison methods for point processes as given by Whitt.

Keywords: Point processes, renewal processes, distance functions, metrics, comparability of processes.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

Research done whilst a visitor in the Department of Statistics, University of North Carolina, Chapel Hill, and supported by the Office of Naval Research under Contract N00014-75-C-0809.

1. INTRODUCTION

When are two point processes close to one another? Indeed, when are two stochastic processes close to one another? These notes are basically intended as a review of some of the literature concerning the proximity of one point process to another, and necessarily therefore also includes notions of one point process being in some sense larger or thicker than another.

Since part of the difficulty in defining the distance between point processes may be seen in defining the distance between two random variables (as distinct from processes, which are indexed families of random variables), we start with some ideas related to the distance of two random variables, then review some work of Kalžanov measuring the closeness of a stationary renewal process to a Poisson process, proceed to ideas of comparison of point processes, and finally come to a (brief) section covering some ideas on distance functions for point processes.

2. DISTANCE BETWEEN RANDOM VARIABLES

The distance between two points in d -dimensional Euclidean space R^d is a familiar enough idea, so that the distance between realizations of two R^d -valued random variables (r.v.s) X and Y , defined on some common probability space (Ω, \mathcal{F}, P) say, is sensibly given by

$$|X(\omega) - Y(\omega)| \quad (2.1)$$

Accession For
NTIS GFA& ✓
DTIC TAB
Unannounced
Justification
By _____
Distribution/ _____
Availability Codes
Dist April 1968
A

for each ω in Ω , with $|\cdot|$ here denoting the usual distance function in R^d . As soon as we move away from just realizations of two r.v.s and seek to measure the closeness or otherwise of the r.v.s per se, by which we mean the measurable functions X, Y mapping Ω into R^d , we are forced either to perform some operation on the random function $|X(\omega) - Y(\omega)|$, or else to have recourse to some other notion not necessarily related to the distance of the realizations of the two functions involved.

Clearly the function

$$d_E(X, Y) = \int_{\Omega} |X(\omega) - Y(\omega)| P(d\omega) \quad (2.2)$$

is an obvious candidate for measuring the distance between the r.v.s X and Y . Provided the r.v.s X and Y have been given, and that $d_E(X, Y) < \infty$, the definition (2.2) is not easily disputed. However, it is not uncommon to be given not two r.v.s X and Y but their distributions in R^d : for the present it will be enough to consider the case of the distribution functions (d.f.s) F and G of R^1 -valued r.v.s, so that for example $F(t) = P\{X \leq t\}$. One metric on the space of d.f.s with finite first moment is the function

$$d_{df}(F, G) = \int_{-\infty}^{\infty} |F(t) - G(t)| dt. \quad (2.3)$$

Manipulation shows that $d_{df}(F, G) = d_E(X^*, Y^*)$ for the r.v.s X^* and Y^* , with d.f.s F and G respectively, defined by

$$X^*(\omega) = F^{-1}(U(\omega)), \quad Y^*(\omega) = G^{-1}(U(\omega)) \quad (2.4)$$

where F^{-1} and G^{-1} are the functional inverses of the d.f.s F and G , and $U(\omega)$ is a r.v. on (Ω, F, P) with the uniform

distribution, so that $P\{U(\omega) \leq t\} = t$ for $0 \leq t \leq 1$. Thus, (2.3) may be expressed as

$$d_{df}(F,G) = d_E(X^*,Y^*) = \int_0^1 |F^{-1}(u) - G^{-1}(u)| du. \quad (2.5)$$

When the d.f.s F and G are partially ordered in the distributional sense, $F \leq_d G$, meaning that $F(t) \geq G(t)$ (all t in \mathbb{R}), then $X^* \leq_{a.s.} Y^*$, and hence

$$d_{df}(F,G) = |EX - EY| = EY - EX.$$

If F and G cannot be ordered in this way, or if the pair (X,Y) of r.v.s is not equivalent in distribution to (X^*,Y^*) , then $d_E(X,Y)$ also includes some measure of the dispersion of X and Y . For example, when F and G are normal d.f.s with zero mean and variances 1 and $\sigma^2 < 1$, $d_E(X^*,Y^*) = (1-\sigma)(2/\pi)^{1/2}$. At the opposite extreme from $(X^*,Y^*) = (F^{-1}(U(\omega)), G^{-1}(U(\omega)))$ is the pair $(F^{-1}(U(\omega)), G^{-1}(1-U(\omega)))$, for which it may be shown that

$$\begin{aligned} \int_0^1 |F^{-1}(u) - G^{-1}(1-u)| du &= 2E|Z-\xi| \\ &= E|X-\xi| + E|Y-\xi| \end{aligned} \quad (2.6)$$

where the r.v. Z has $\frac{1}{2}(F(t) + G(t))$ as its d.f. and ξ is a median of Z . Intermediate between (2.6) and (2.5), when X and Y are independently distributed r.v.s with d.f.s F and G , $E|X-Y|$ again vanishes (like (2.6) but unlike (2.5)) if and only if $X = Y = \text{constant a.s.}$: (2.5) vanishes if and only if $X =_d Y$.

A major difficulty in extending the definition d_E or d_{df} to \mathbb{R}^d -valued r.v.s, relates to the need of an analogue of the construction of (X^*,Y^*) . Consider the following (cf. also Daley (1981)).

Problem 1. Define R^2 -valued r.v.s $\underline{X} = (X_1, X_2)$ and $\underline{Y} = (Y_1, Y_2)$ distributed uniformly over a circle of unit area with centre at the origin, and a square of unit area with centre at the origin, respectively. Amongst all r.v.s with these two specified distributions, is there a construction of r.v.s X^* , Y^* such that

$$d_E(X^*, Y^*) \leq d_E(X, Y) ?$$

A strategy that avoids the sample path problem, given two distributions, is the use of metrics on the distributions on the state space of the realizations induced by the r.v.s. Thus we have the variation metric

$$d_V(F, G) \equiv \frac{1}{2} \int_{R^d} |F(dt) - G(dt)| \quad (2.7)$$

for R^d -valued r.v.s with distributions F and G , and in particular, for integer-valued r.v.s X and Y ,

$$\begin{aligned} d_V(X, Y) &= \frac{1}{2} \sum_i |P\{X=i\} - P\{Y=i\}|, \\ &= P\{X^\dagger \neq Y^\dagger\} \end{aligned} \quad (2.8)$$

for appropriately defined r.v.s (X^\dagger, Y^\dagger) (and this construction is in general different from (X^*, Y^*)). Also, for real-valued r.v.s X and Y , we have the supremum metric

$$d_{\text{sup}}(X, Y) = \sup_t |P\{X \leq t\} - P\{Y \leq t\}|, \quad (2.9)$$

the Lévy metric and others.

3. DISTANCE BETWEEN POINT PROCESSES: INTRODUCTORY COMMENTS

Realizations of point processes on R can be described either as counting measures or by the distances of the nearest point to the right of the origin and by the lengths of the intervals between successive points enumerated respectively on each side of that point. For the present discussion we confine attention to point processes on $R_+ = (0, \infty)$, and assume without essential loss of generality that every realization has $N(R_+) \geq 1$ where N denotes the counting measure. We use N, N_1, N_2, \dots as generic names of point processes. The two descriptions referred to above are related via

$$S_0 = 0, \quad S_n = \inf\{t: N(0,t] \geq n\}, \quad (3.1)$$

and $X_n = S_n - S_{n-1}$ ($n = 1, 2, \dots$): X_1 is the forward recurrence time interval, and $\{X_n: n = 2, 3, \dots\}$ the successive intervals between points so long as $S_n < \infty$; otherwise we may set $X_n = \infty$.

Given two point processes N_i ($i = 1, 2$), with intervals $\{X_n\}$ ($i=1$) and $\{Y_n\}$ ($i=2$), the discussion of section 2 leads to consideration as "natural" distance functions for N_1 and N_2 either functions of $|N_1(t) - N_2(t)|$, or else of the sequence $\{X_n - Y_n\}$. The simplest constructions of point processes usually proceed via interval properties rather than counting properties (though as a counter-example of this conventional construction, recall Lewis and Shedler's (1979) discussion on realizations of inhomogeneous Poisson processes). Accordingly, the distance of N_1 and N_2 may be studied via functionals of $\{X_n^* - Y_n^*\}$ where the elements (X_n^*, Y_n^*) are constructed successively from the sequence

$\{U_n^*\}$ of independent identically distributed (i.i.d.) r.v.s uniformly distributed on $(0,1)$. Such a construction for renewal processes N_i is sensible, and also for point processes whose intervals are stochastically monotone Markov chains or even certain semi-Markov processes (see Sonderman (1980)). However, when the successive members of the sequence $\{X_n\}$ are not monotonically dependent on their predecessors, there may not necessarily exist a mapping linking $\{X_n^*\}$ and $\{Y_n^*\}$ that is unequivocally the optimum one.

4. THE MARGINAL COUNTING DISTRIBUTIONS

Let N_1 be a stationary renewal process for which $E N_1(0,1] = \lambda$ and the lifetime d.f. is F , so that $\{X_n : n = 2,3,\dots\}$ are i.i.d. r.v.s with d.f. F and

$$F_1(t) \equiv P\{X_1 \leq t\} = \lambda \int_0^t (1-F(u))du. \quad (4.1)$$

Kalžanov (1970, 1975a, 1975b) aimed to compare N_1 with the most "typical" (?) of point processes, namely, he took a Poisson process N_2 with rate parameter λ , so that

$$P\{N_2(0,t] = k\} \equiv v_k(t) = (\lambda t)^k e^{-\lambda t} / k!. \quad (4.2)$$

His comparison was based on the probabilities $\{v_k(\cdot)\}$ and

$$\begin{aligned} P_k(t) &= P\{N_1(0,t]=k\} = P\{S_k \leq t < S_{k+1}\} \\ &= \int_0^t (F^{(k-1)*}(t-u) - F^{k*}(t-u))dF_1(u) \quad (4.3a) \\ &= \lambda \int_0^t (F^{(k-1)*}(u) - 2F^{k*}(u) + F^{(k+1)*}(u))du, \end{aligned}$$

provided $k \geq 1$; for $k = 0$,

$$P_0(t) = 1 - F_1(t) . \quad (4.3b)$$

Kalžanov (1975a) claimed that

$$\delta_k \equiv \rho_1(P_k, v_k) \equiv \sup_t |P_k(t) - v_k(t)| \leq \begin{cases} \rho_1 & (k = 0) , \\ (3 + 4\ln(1/\rho_1))\rho_1 & (k = 1, 2, \dots) , \end{cases} \quad (4.4)$$

$$\text{where } \rho_1 \equiv \sup_t |F(t) - F_1(t)| \quad (4.5)$$

so that $\rho_1 = 0$ if and only if $F(t) = F_1(t) = 1 - e^{-\lambda t}$ and hence N_1 is a Poisson process. He does indeed establish (4.4) for $k = 0, 1, 2, 3$, and shows that $\sup_{\lambda t \leq \ln(1/\rho_1)} |P_k(t) - v_k(t)|$ is bounded as asserted, but his proof for $\lambda t > \ln(1/\rho_1)$ is erroneous, being based on the false assertion that for non-negative r.v.s X and Y and $a < b$,

$$\text{"Pr}\{a \leq X + Y < b\} \leq \text{Pr}\{a \leq X < b\} \text{" .} \quad (*)$$

Problem 2. Investigate whether the bounds at (4.4) hold for $k \geq 4$.

Can the coefficient of ρ_1 be tightened?

Problem 3. It was also claimed by Kalžanov (1975a) that

$$\delta_k \leq \begin{cases} \lambda\rho_2 & (k = 0) , \\ (5 + 2\ln(1/\lambda\rho_2))\lambda\rho_2 & (k = 1, 2, \dots) , \end{cases} \quad (4.6)$$

where $\lambda\rho_2 = \lambda \int_0^\infty |F(t) - F_1(t)| dt$. Investigate whether these bounds on δ_k hold for $k \geq 3$.

Since these publications may not be so readily accessible, an outline of the (correct) part of their derivation is given later in this section (see around (4.15)).

Kalžanov's motivation is clear enough; he is comparing N_1 with a Poisson process when the lifetime d.f. F and stationary forward recurrence time d.f. F_1 are close, and hence F is close to the exponential d.f., because as shown in Kalžanov (1970),

$$F_1(t) - (1 - e^{-\lambda t}) = \int_0^t (F_1(t-u) - F(t-u)) \lambda e^{-\lambda u} du, \quad (4.7)$$

so

$$\sup_t |F_1(t) - (1 - e^{-\lambda t})| \leq \rho_1 (1 - e^{-\lambda t}) \leq \rho_1, \quad (4.8)$$

$$\int_0^t |F_1(t) - (1 - e^{-\lambda t})| dt \leq \rho_2 (1 - e^{-\lambda t}) \leq \rho_2, \quad (4.9)$$

and
$$\sup_t |F(t) - (1 - e^{-\lambda t})| \leq \rho_1 + \sup_t |F_1(t) - (1 - e^{-\lambda t})| = 2\rho_1. \quad (4.10)$$

However, the bound (assumed true) is somewhat larger than is probably likely, because it exceeds 1 for $\rho_1 \geq .074778$, as in the table below:

ρ_1	$(3 + 4 \ln(1/\rho_1)) \rho_1$
.01	.214207
.02	.372962
.03	.510787
.04	.635020
.05	.749146
.06	.855219
.07	.954593
.074778	1.0 ₆

Suppose the lifetime d.f. F has a finite second moment. It may then be of more interest to compare the marginal probabilities or even the simpler probabilities

$$P_k^0(t) - F^{k*}(t) - F^{(k+1)*}(t) \quad (4.11)$$

of a renewal process $N_1^0(\cdot)$ with F as the d.f. of X_1 as well as X_2, X_3, \dots , with the corresponding probabilities $v_{k,\gamma}(t)$ or $v_{k,\gamma}^0(t)$ of a stationary renewal or renewal process respectively with gamma distributed intervals Y_2, Y_3, \dots matching X_2, X_3, \dots in their first two moments. Then for moderate (to large) values of k , using ϕ to denote the standard normal d.f., and $\sigma^2 = \text{var } X_n$ ($n \geq 2$),

$$P_k^0(t) \approx \phi((t-k\lambda^{-1})/(\sigma k^{1/2})) - \phi((t-(k+1)\lambda^{-1})/(\sigma(k+1)^{1/2})), \quad (4.12)$$

which is possible for all $t \geq 0$, and, of course, then

$P_k^0(t) \approx v_{k,\gamma}^0(t)$. ($P_k^0(t)$ may have discrete jumps, but the approximation will remain valid.) Indeed it is immediately obvious that, provided X_n has a finite third moment, the Berry - Esseen bound may be applied to bounding $|P_k^0(t) - v_{k,\gamma}^0(t)|$ uniformly in t , $\rightarrow 0$ as $k \rightarrow \infty$ like $O(k^{-1/2})$ - though large k is needed before this is necessarily smaller than one.

Recognizing from (4.3) that for $k \geq 1$,

$$P_k(t) = \int_0^t P_{k-1}^0(t-u) dF_1(u)$$

and that $P_k^0(t) = \int_0^\infty (F^{k*}(t) - F^{k*}(t-u)) dF(u)$, (4.13)

another possible approach to bounding $P_k(t)$ is to use the concentration function of F which behaves like $k^{-1/2}$ for increasing k (see section 2.2 of Hengartner and Theodorescu (1973)). Alternatively, smoothing inequalities on transforms may be useful (cf. Paulauskas (1971), Petrov (1975)).

For approximation purposes, observe that when $\phi(\cdot)$ denotes the standard normal probability density function,

$$\sup_t |\phi(t) - \sigma^{-1} \phi(t/\sigma)| = (1-\sigma')/\sigma' \quad (4.14)$$

where $\sigma' = \min(\sigma, \sigma^{-1})$.

Kalžanov's (1975a) partial derivation of (4.4) comes from (4.8) and (4.10) applied to

$$P_k(t) - v_k(t) = \int_0^t P_{k-1}(t-u)dF(u) - \int_0^t v_{k-1}(t-u)dG(u) \quad (4.15)$$

which holds for $k = 2, 3, \dots$, where we have written $G(u) = 1 - e^{-\lambda u}$, and, using convolution notation,

$$P_1(t) - v_1(t) = (F_1 - F * F_1)(t) - (G - G^{2*})(t).$$

(4.7) can be written as $F_1 - G = (F_1 - F) * G$. Then

$$\begin{aligned} P_1 - v_1 &= (F_1 - F) * G - F * F_1 + G * G \\ &= (G - F) * (F_1 + G), \end{aligned}$$

$$|P_1(t) - v_1(t)| \leq (F_1 + 2G + G^{2*})\rho_1.$$

Generally,

$$P_{k+1} - v_{k+1} = (P_k - v_k) * F + v_k * (F - G)$$

$$\begin{aligned} \text{so } |P_{k+1}(t) - v_{k+1}(t)| &\leq \sup_{u \leq t} |P_k(u) - v_k(u)| + (G^{k*} + G^{(k+1)*}) * (G^0 + G)\rho_1 \\ &\leq \rho_1 (F_1 + 3G + 4G^{2*} + \dots + 4G^{k*} + 3G^{(k+1)*} + G^{(k+2)*})(t). \end{aligned}$$

The coefficient of ρ_1 at (4.4) ≥ 12 for $\rho_1 \leq .074778$ so it holds for $k = 1, 2, 3$. For all k and finite t , the coefficient of ρ_1 in the bound on $P_{k+1}(t) - v_{k+1}(t)$ is at most $\rho_1(F_1(t) - G(t) + 4\lambda t) \leq \rho_1(\rho_1 + 4\lambda t)$.

Kalžanov (1975b) endeavours to bound the difference

$|P_k^{(1)}(t) - P_k^{(2)}(t)|$ of the probabilities $P_k^{(i)}(t)$ of stationary renewal processes N_i ($i=1, 2$) in terms of the metric $\sup_t |F^{(1)}(t) - F^{(2)}(t)|$ on the lifetime d.f.s $F^{(i)}(\cdot)$, assuming these to be absolutely continuous d.f.s with bounded densities and finite second moments. The argument

resembles that outlined above, and has the same appeal to the erroneous statement (*) .

Azlarov and Kalžanov (1976a) refers to Kalžanov (1975a) in deriving bounds on the difference of the stationary waiting time d.f. for a GI/M/k from the same d.f. in an M/M/k queue with the same arrival and service rates. Bounds are given for each of the cases of an IFR and DFR inter-arrival d.f. in GI/M/k. Details of the proof are omitted, so it is not clear whether (*) has been used. In Azlarov and Kalžanov (1976b) the loss probabilities in pure loss GI/M/k and M/M/k systems are compared, but only the abstract in Mathematical Reviews is available.

While for example the sequence $\{\delta_k\}$ provides some information on all the one-dimensional marginal distributions $\{P_k\}$ as compared with those $\{v_k\}$ of a Poisson process - or, indeed, we could consider $\{\delta_k\}$ for any two point processes via

$$\left\{ \sup_t |\Pr\{N_1(0,t]=k\} - \Pr\{N_2(0,t]=k\}| \right\} ,$$

we may prefer, as an indicator of the "distance" between the processes N_1 , some function that discounts behaviour for either large t or large k or both, such as (for appropriate $\epsilon > 0$ and z in $0 < z < 1$)

$$\sum_{k=0}^{\infty} z^k \int_0^{\infty} e^{-\epsilon t} |\Pr\{N_1(0,t]=k\} - \Pr\{N_2(0,t]=k\}| dt .$$

Alternatively, interest in large k may be measured by for example $\sup_t k^{\frac{1}{2}} |P_k^{(1)}(t) - P_k^{(2)}(t)|$.

Trivially, since there exist point processes with distinct probability distributions whose finite-dimensional distributions agree up to some finite order r (cf. Szasz (1970) and Oakes (1974)), no single member of $\{\delta_k\}$, nor even the entire sequence, nor any function

based on finite dimensional distributions of bounded order, will be a metric on the space of point processes. Of course $\sup_t |F^{(1)}(t) - F^{(2)}(t)|$ is a metric on the space of renewal processes, as these are characterized by their lifetime d.f.s.

An entry in Mathematical Reviews refers to Bol'sakov and Rakošic (1978) as having studied in detail Poisson and close-to-Poisson point processes. The meaning of the phrase in the context can only be guessed at.

5. COMPARABILITY OF POINT PROCESSES

The idea of studying the distance of one object from another, is an endeavour by implication to compare the two objects. Comparisons in stochastic analysis usually refers to the partial ordering of the objects, and consideration of the variety of definitions proposed for the comparability of point processes may be helpful. Let \prec denote a partial ordering on the space X of (real-valued) r.v.s on (Ω, F, P) : Stoyan (1977) surveys some of the ideas and properties of some partial orderings on X , mostly expressed in terms of partial orderings on functions of the probability measures P_X induced on (Ω, F, P) via $X \in X$.

For point processes, work of Schmidt (1976) that Stoyan reviews is covered in Whitt (1981) whose survey is the source of the definitions below; Whitt's numerical numbering of definitions is replaced by a more suggestive notation. The definitions are interspersed with some comments; write $N(0, t] = N(t)$ for brevity. Stoyan (1977) has an appealing terminology: if $N_1 \prec N_2$ for some partial ordering \prec on point processes, say that N_1 is \prec -thinner than N_2 , or, equivalently, that N_2 is \prec -thicker than N_1 .

Define

$$F(x|H_t) = \Pr\{T_{N(t)+1} - t \leq x | N(s), 0 < s \leq t\}, \quad (5.1)$$

the forward recurrence time d.f. at t conditional on the entire history H_t of $N(\cdot)$ up to time t . Then say

$$N_1 \leq_h N_2 \quad \text{if} \quad F^{(1)}(x|H_t) \leq F^{(2)}(x|H_t) \quad (5.2)$$

holds for all $x > 0$ and all histories H_t . In other words, the conditional forward recurrence time is stochastically larger in N_1 than N_2 for all realizations on $(0, t)$ and for all t .

In the particular case that (say) N_1 is a renewal process with lifetime d.f. F , and N_2 is a possibly inhomogeneous Poisson process with $EN_2(0, t] = \Lambda(t)$, (5.2) implies that

$$\sup_{0 \leq u \leq t} (F(u+x) - F(u)) / (1 - F(u)) \leq 1 - e^{-(\Lambda(t+x) - \Lambda(t))}. \quad (5.3)$$

This condition is more easily written via the representation, valid for all u for which $F(u) < 1$, that $F(u) = 1 - e^{-\mu(u)}$ for some non-decreasing function $\mu(\cdot)$, $\mu(u) \rightarrow \infty$ ($u \rightarrow \infty$). Then (5.3) is equivalent to

$$\mu(u+x) - \mu(u) \leq \Lambda(t+x) - \Lambda(t) \quad (\text{all } 0 \leq u \leq t),$$

and it follows that $\Lambda(t+0) = \infty$ for $\inf\{t: \mu(t+0) > \mu(t-0)\}$ if such set is non-empty. This would lead to a Poisson process exploding at such t , as would be the case also if $\mu(x) \rightarrow \infty$ for $x \rightarrow x_0 < \infty$.

Eliminating these cases, we are left with the (interesting) case that $\Lambda(t) \leq Ct$ (all t) for some finite constant $C \geq \lim_{v \downarrow 0} (\mu(u+v) - \mu(u)) / v \equiv h(u)$ (all u) provided the limit exists. Indeed, it suffices to write $m(t) = \sup_{0 < u < t} h(u)$, and then we may take

$$\Lambda(t) = \int_0^t m(u) du .$$

This is the smallest function $\Lambda(\cdot)$ satisfying (5.3); Miller (1979) also showed that such a nonhomogeneous Poisson process can be constructed so that N_2 will bound the renewal process N_1 with lifetime d.f.

$$F(x) = 1 - \exp\left(-\int_0^x h(u) du\right)$$

in the sense that $N_1 \leq_h N_2$. He also shows that an inhomogeneous Poisson process N_0 with $EN_0(0,t] = \int_0^t (\inf_{0 \leq u \leq x} h(u)) dx$ satisfies

$$N_0 \leq_h N_1 .$$

Returning to general point processes N_1 and N_2 , suppose that their realizations satisfy

$$N_1(x, x+dx) \leq N_2(x, x+dx) \text{ a.s.} \tag{5.4}$$

for all $0 < x < \infty$, i.e., every jump in the counting function $N_1(x, \omega)$ is also a jump in $N_2(x, \omega)$, and its size in N_2 is at least as large as in N_1 . When (5.4) is satisfied, write

$$N_1 \subseteq N_2 \tag{5.5}$$

since the definition is precisely that the sets $\{S_n^{(1)}\}$ as defined by (3.1) satisfy such an inclusion relation. If N_1 is obtained from N_2 by any thinning operation, then clearly $N_1 \subseteq N_2$.

If $N_1 \leq_h N_2$, then (there exists a common probability space on which N_1 and N_2 may be defined such that)* $N_1 \subseteq N_2$, but the converse need not hold. However, if $N_1 \subseteq N_2$ and both N_i are Poisson processes, then the converse will hold, because the intensity measures Λ_i will satisfy $\Lambda_1(A) \leq \Lambda_2(A)$ for all bounded Borel sets A , and for

*This parenthesized remark is to be understood in defining $N_1 \subseteq N_2$, $N_1 \leq_{int} N_2$ and $N_1 \leq_p N_2$.

inhomogeneous Poisson processes, $F(x|H_t) = 1 - \exp(-\Lambda(t,t+x))$ independent of H_t . If both N_1 are doubly stochastic Poisson processes for which the sample realizations of the intensity functions satisfy $\Lambda_1(A,\omega) \leq \Lambda_2(A,\omega)$ a.s. for all Borel sets A , then while we have $N_1 \subseteq N_2$, we need not have $N_1 \subseteq_h N_2$ because the "history" $H_t \equiv \{N_1(s): 0 < s \leq t\}$ is not equivalent to $\Lambda_1(\cdot,\omega)$. This statement could be rephrased in measure-theoretic language, but an illustration is probably worth more in word-value: let $\Lambda_1(\cdot,\omega)$ have the densities $\lambda_1(\cdot,\omega)$ given by

$$\lambda_1(t) = \begin{cases} 0 & \text{for } t \leq 1, & 2 & \text{for } t > 1 & \text{with probability } .5, \\ 2 & \text{for } 1 < t \leq 2, & 1 & \text{otherwise with probability } .5, \end{cases}$$

$$\lambda_2(t) = \begin{cases} 2 & \text{for all } t & \text{with probability } .5, \\ 2 & \text{for } t \leq 2, & 1 & \text{for } t > 2 & \text{with probability } .5. \end{cases}$$

Then $\Pr\{N_1(2,x) = 0 | N_1(0,2) = 0\} = (e^{-2(x-2)} + e^{-1}e^{-(x-2)}) / (1 + e^{-1})$

$$\Pr\{N_2(2,x) = 0 | N_2(0,2) = 0\} = (e^{-2(x-2)} + e^{-(x-2)}) / 2,$$

so that $F^{(1)}(x | N_1(0,2) = 0) > F^{(2)}(x | N_2(0,2) = 0)$, contra (5.2) if we were to have $N_1 \subseteq_h N_2$.

Particularly as inputs to queueing systems, there is much interest in the intervals $\{X_n\}$ following (3.1). For N_1 defined on a common probability space, say that

$$N_1 \leq_{\text{int}} N_2 \text{ if } X_n^{(1)} \geq X_n^{(2)} \text{ (all } n = 1, 2, \dots) \text{ a.s.} \quad (5.6)$$

In other words, the points of N_1 are more spaced out than those of N_2 .

If N_1 are renewal processes with lifetime d.f.s F, G for which $F(x) \leq G(x)$ (all x) (in the language of Stoyan, the d.f. F is larger than G), then there are realizations $\{X_n^{(1)}\}$ for which (5.6) is

necessarily satisfied. It need not be the case that if $N_1 \subseteq N_2$, then $N_1 \leq_{\text{int}} N_2$, nor conversely, although if N_i are renewal processes satisfying $N_1 \subseteq N_2$, then there will be renewal processes N'_i distributed like N_i for which $N'_1 \leq_{\text{int}} N'_2$. In terms of models for point processes, the ordering \leq_{int} arises naturally only when the process is conveniently specified by its intervals (cf. also the comments in section 3).

It will be evident that all three orderings presented so far have the property of implying that $N_1 \prec N_2$ yields

$$N_1(0, t] \leq N_2(0, t] \quad (\text{all } t) \quad \text{a.s.}, \quad (5.7)$$

and when (5.7) holds, we shall say that

$$N_1 \leq_p N_2. \quad (5.8)$$

To justify the notation as suggesting that some underlying probabilistic inequality may suffice, recall the equivalence of events

$$\{N_1(0, x_j] \leq n_j \quad (j = 1, \dots, k)\} = \{S_{n_j}^{(1)} \geq x_j \quad (j = 1, \dots, k)\},$$

and that the multivariate analogue of the inequality $X \leq_d Y$ between r.v.s X and Y is that

$$Ef(X_1, \dots, X_k) \leq Ef(Y_1, \dots, Y_k) \quad (5.9)$$

for all functions f increasing in each argument such that the expectations exist. If for all such f we have

$$Ef(\{S_n^{(1)}\}) \geq Ef(\{S_n^{(2)}\}) \quad (5.10)$$

then by an extension of a result of Strassen (see Chapter 17 of Marshall and Olkin (1979), or, for more detail, Kamae, Krengel and O'Brien (1977)) there is a probability space on which

$$S_n^{(1)} \geq S_n^{(2)} \quad (\text{all } n) \quad \text{a.s.}, \quad (5.11)$$

and hence (5.7) holds also.

Since (5.11) for $n = 1$ implies that $X_1^{(1)} \geq X_1^{(2)}$ a.s., it follows immediately that for renewal processes, $N_1 \leq_P N_2$ implies $N_1 \leq_{\text{int}} N_2$, but this result need not be true in general. For example, N_1 may be a renewal process with lifetime d.f. F , and N_2 an alternating renewal process with lifetime d.f.s G_1 and G_2 for which for all x , $G_1(x) \geq F(x) \geq G_2(x)$ and

$$\int_0^x G_1(x-u) dG_2(u) \geq F^{2*}(x); \quad \text{this is an easy construction}$$

(take $F(x) = G_2(x+a)$, $G_1(x) = G_2(x+a+b)$ for $0 < b < a$).

The last inequality Whitt lists is just the marginal distribution property that

$$N_1(t) \leq_d N_2(t) \quad (\text{all } t). \quad (5.12)$$

The quasi-Poisson processes of Szasz (1970) or Oakes (1974) afford simple examples of processes satisfying $N_1 \leq_d N_2$ but none of the other partial orderings. (As Whitt notes, (5.12) is not in fact a partial ordering on the space of point processes, because, as for example with the quasi-Poisson processes, we can have $N_1 \leq_d N_2$ and $N_2 \leq_d N_1$ without N_1 having a common (joint) distribution.) Note that if (5.12) holds, then, by the equivalence as below (5.8), $S_n^{(1)} \leq_d S_n^{(2)}$ (all n).

Schmidt (1976) formulated the definition (5.12), unnecessarily restricted to stationary point processes, and also formulated a distributional version of \leq_{int} for stationary point processes via the inequality

$$E_0 f(\{X_n^{(1)}\}) \geq E_0 f(\{X_n^{(2)}\}) \quad (5.13)$$

with f as at (5.13) and E_0 denoting expectation with respect to the Palm probabilities, for which $\{X_n^{(i)}\}$ are then stationary sequences. By the extension of the Strassen result, there then exist point processes N_i^0 for which $N_1^0 \leq_{\text{int}} N_2^0$, these processes having the Palm distribution of the stationary processes N_i .

Note that if N_i are renewal processes for which $N_1 \leq_d N_2$, then since

$$\Pr\{N_1(t) = 0\} \geq \Pr\{N_2(t) = 0\},$$

we must also have $N_1 \leq_{\text{int}} N_2$. But if the N_i are stationary renewal processes, neither \leq_{int} nor \leq_d need imply the other, as Schmidt showed.

In subsequent work, Whitt (1981b) has taken a more pragmatic view and compared some point processes numerically by studying how different processes may have different effects on a queueing system when fed as inputs into the system. More work of this nature has been done by Whitt and his colleagues.

6. DISTANCES BETWEEN POINT PROCESSES

For the time being we pass over the comparisons \leq_h and \leq_c for suggesting distances between point processes. Coming to \leq_{int} , and recalling (2.3), we could use

$$d_{\text{int},1}(N_1, N_2) \equiv \sup_{n \geq 1} n^{-1} \int_0^\infty |\Pr\{S_n^{(1)} > t\} - \Pr\{S_n^{(2)} > t\}| dt \quad (6.1)$$

where the factor n^{-1} has been introduced to average out the effect of $S_n^{(1)} - S_n^{(2)}$ being asymptotically like $n(EX_1^{(1)} - EX_1^{(2)})$ if the sequences $\{X_n^{(i)}\}$ are stationary. However, if $N_1 \leq_d N_2$ and $N_2 \leq_d N_1$, as at (5.11) and below, we should have $d_{\text{int},1}(N_1, N_2) = 0$, for (6.1) depends

only on the marginal distributions of $N^{(1)}(t)$, not the joint distribution over a set of values of t .

$$d_E(N_1, N_2) = \sup_{t>0} |EN_1(t) - EN_2(t)|/(t+1) \quad (6.2)$$

has a similar drawback. However

$$d_{E^*}(N_1, N_2) = \sup_{t>0} E|N_1(t) - N_2(t)|/(t+1) \quad (6.3)$$

does not have this drawback: instead we are confronted with the problem of requiring N_1 and N_2 to be defined on the same probability space, and, as noted in section 2, the appropriate construction is far from evident as soon as we dispense with renewal processes. As an alternative to (6.3), and one that is better suited to processes constructed by the interval properties, we have

$$d_{\text{int}}(N_1, N_2) = \sup_{n \geq 1} n^{-1} E|S_n^{(1)} - S_n^{(2)}|. \quad (6.4)$$

For renewal processes, defined via sequences $\{(X_n^{(1)*}, X_n^{(2)*})\}$ as in section 3, $d_{\text{int}} = d_{\text{int},1}$, although the proof is not immediately obvious.

Since renewal processes N_i are characterized by their renewal functions $U_i(\cdot)$, we could use, as an alternative to (6.2) (but only for renewal processes) the function

$$\sup_{t \geq 0} |U_1(t) - U_2(t)|/(t+1). \quad (6.5)$$

Matthes, Kerstan and Mecke (1978) (e.g. in their Sections 1.9 and 1.12) use the variation distance for point processes, and T.C. Brown (private correspondence) has written of using compensators to obtain total variation results for the distance of point processes

from Poisson processes. I have been told of an unpublished review by R.L. Farrell that may have relevance to this section. Mark Brown (1981 and correspondence) has compared renewal processes with decreasing failure rate (DFR): in the notation of section 2, he has shown that for distributions with the weaker property of increasing mean residual life,

$$\max(\rho_1(F, F_1), \rho_1(F, G), \rho_1(F_1, G)) \leq 1 - 2(EX)^2/EX^2; \quad (6.6)$$

for DFR d.f.s it is easily checked from a sketch of the tails of the d.f.s involved and the results of section 2, that the maximum on the left-hand side of (6.6) is in fact $\rho_1(F, F_1)$. He has also studied metrics in terms of the hazard functions being close.

We have not attempted to assess the effect of common operations on point processes (thinning, superposition, translation, cluster-formation, etc.) on the distance functions given above.

ACKNOWLEDGEMENT I am greatly indebted to Ross Leadbetter for his interest.

REFERENCES

- AZLAROV, T.A. & KALŽANOV, K. (1976a) Estimation of the stability of a certain system with waiting time. (In Russian). Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk 1976, no. 5, 68-69. MR 57 # 4354.
- & ——— (1976b) An estimate of the stability of a queueing system. (In Russian). Taskent. Gos. Univ. Naučn. Trudy Vyp. 490 Voprosy Matematiki (1976), 21-26. MR 58 # 3092.
- BOL'SAKOV, I.A. & RAKOŠIC, V.S. (1978) Applied Theory of Random Flows. (In Russian). Izdat. "Sovetskoe Radio", Moscow. MR 80g: 60055

- BROWN, M. (1981) Approximating DFR distributions by exponential distributions, with applications to first passage times.
Ann. Probab. 9
- DALEY, D.J. (1981) The closeness of uniform random points in a square and a circle. Tech. Report, Mimeo Series, Statistics Dept., University of North Carolina, Chapel Hill.
- HENGARTNER, W., & THEODORESCU, R. (1973) Concentration Functions. Academic Press, New York.
- KALŽANOV, K. (1970) The closeness of the Palm and Poisson flows. (In Russian). Izv. Akad. Nauk UzSSR Ser. Fiz-Mat. Nauk 14 (1970), no. 3, 69-71. MR 44 # 1073.
- (1975a) The stability of a Poisson flow. (In Russian). Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk 1975, no. 2, 75-77. MR 55 # 11431.
- (1975b) The nearness of two Palm flows. (In Russian). Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk 1975, no. 4, 25-30. MR 55 # 11432.
- KAMAE, T., KRENGEL, U. & O'BRIEN, G.L. (1977) Stochastic inequalities on partially ordered spaces. Ann. Probab. 5, 899-912.
- LEWIS, P.A.W. & SHEDLER, G.S. (1979) Simulation of nonhomogeneous Poisson processes by thinning. Naval Res. Logistics Quart. 26, 403-413.
- MARSHALL, A.W. & OLKIN, I. (1979) Inequalities: Theory of Majorization and Its Applications. Academic Press, New York.
- MATTHES, K., KERSTAN, J. & MECKE, J. (1978) Infinitely Divisible Point Processes. John Wiley & Sons, Chichester.
- OAKES, D. (1974) A generalization of Moran's quasi-Poisson process. Studia Sci. Math. Hungar. 9, 433-437.

- PAULAUSKAS, V. (1971) On a smoothing inequality (in Russian).
Litovsk. Mat. Sb. 11, 861-866 (= Selected Transl. Math. Statist. Probab. 14 (1977), 7-12).
- PETROV, V.V. (1975) Sums of independent Random Variables
Springer-Verlag, New York. (= Translation from original Russian edition, 1972).
- SCHMIDT, V. (1976) On the non-equivalence of two criteria of comparability of stationary point processes. Zast. Mat. 15, 33-37.
- SONDERMAN, D. (1980) Comparing semi-Markov processes. Math. Operat. Res. 5, 110-119.
- STOYAN, D. (1977) Qualitative Eigenschaften und Abschätzungen stochastischer Modelle. Akademie-Verlag, Berlin.
- SZASZ, D. (1970) Once more on the Poisson process. Studia Sci. Math. Hungar. 5, 441-444.
- WHITT, W. (1981a) Comparing counting processes and queues.
Adv. Appl. Probab. 13 (to appear).
- (1981b) Approximating a point process by a renewal process,
I: A general framework. Operat. Res. 29 (to appear).

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER	2. GOVT ACCESSION NO. AD-A097 622	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Distances of Random Variables and Point Processes.	5. TYPE OF REPORT & PERIOD COVERED TECHNICAL paper.	6. PERFORMING ORG. REPORT NUMBER Mimeo Series No. 1331
7. AUTHOR(s) D.J./Daley	8. CONTRACT OR GRANT NUMBER(s) 13 ONR Contract N00014-75-C-0809	10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS 1223
9. PERFORMING ORGANIZATION NAME AND ADDRESS 14 MIMED SER-1331	11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics and Probability Program (Code 436) Arlington, VA 22217	12. REPORT DATE March 1981
13. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	14. NUMBER OF PAGES 23	15. SECURITY CLASS (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release--Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) point processes, renewal processes, distance functions, metrics, comparability of processes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The paper is basically a discussion and survey of some literature concerning the concept of a distance function for random variables and point processes. Work of Kalžanov on the closeness of renewal processes to Poisson processes is outlined, and there are comments on various possible comparison methods for point processes as given by Whitt.		

420064
sk

END

DATE
FILMED

5 81

DTIC