

AD-A097 726

WASHINGTON UNIV SEATTLE DEPT OF CHEMISTRY F/G 7/3  
ENERGY TRANSFER AT HIGH TEMPERATURES IN THE CYCLOPROPANE VEM SY--ETC(U)  
MAR 81 T KASAI, D F KELLEY, B S RABINOVITCH N00014-75-C-0690

ML

UNCLASSIFIED

TR-19

1 of 1  
AS A  
097100



END

DATE  
FILMED

5-81

DTIC

12 LEVEL II

Energy Transfer at High Temperatures in the Cyclopropane VEM System. Temperature Dependence of  $\langle \Delta E \rangle$ .

16 T. Kasai, D. F. Kelley and B. S. Rabinovitch

Department of Chemistry BG-10  
University of Washington  
Seattle, WA 98195

AD A 097 726

9 Technical Report No. NR092-549 TRJ9

14 Contract NR00014-75-C-0690 NR 092-549

15 March 15, 1981

DTIC ELECTE  
S D  
APR 14 1981  
B

OFFICE OF NAVAL RESEARCH  
Department of the Navy  
Code 473  
800 N. Quincy  
Arlington, VA 22217

Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release; its distribution is unlimited.

DTIC FILE COPY

81 4 14 48

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1 REPORT NUMBER <del>NR 092-549</del> TR19	2 GOVT ACCESSION NO. AD-A097	3 RECIPIENT'S CATALOG NUMBER 726
4 TITLE (and Subtitle) Energy Transfer at High Temperatures in the Cyclopropane VEM System. Temperature Dependence of $\langle \Delta E \rangle$		5 TYPE OF REPORT & PERIOD COVERED Technical
		6 PERFORMING ORG REPORT NUMBER
7 AUTHOR(s) T. Kasai, D. F. Kelley and B. S. Rabinovitch		8 CONTRACT OR GRANT NUMBER(s) N00014-75-C-0690 NR 092-549
9 PERFORMING ORGANIZATION NAME AND ADDRESS Professor B. S. Rabinovitch Department of Chemistry BG-10 University of Washington Seattle, WA 98195		10 PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS
11 CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Code 743 Department of the Navy 800 N. Quincy Arlington, VA 22217		12 REPORT DATE March 15, 1981
		13 NUMBER OF PAGES 9
14 MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15 SECURITY CLASS (of this report)
		15a DECLASSIFICATION DOWNGRADING SCHEDULE
16 DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release; its distribution is unlimited.		
17 DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Prepared for publication in Chemical Physics		
18 SUPPLEMENTARY NOTES Prepared for publication in Chemical Physics		
19 KEY WORDS (Continue on reverse side if necessary and identify by block number) Cyclopropane Surfaces Energy Transfer Transients Gases Unimolecular Reaction High Temperature Variable Encounter Method Surfaces Vibrational Relaxation		
20 ABSTRACT (Continue on reverse side if necessary and identify by block number) The study of collisional energy transfer between cyclopropane molecules and a seasoned silica wall by the Variable Encounter Method has been extended upward in temperature to 1325 K. The value of $\langle \Delta E \rangle$ the average energy of a down-transition upon collision of the substrate with the wall continues to decrease monotonically. The data provide the most accurate and well-documented display of the temperature trend of $\langle \Delta E \rangle$ presently available.  (DELTA EPSILON) ↑		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102 LF 014 6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Energy Transfer at High Temperatures in the Cyclopropane VEM System.  
Temperature Dependence of  $\langle \Delta E \rangle$ .<sup>1</sup>

by T. Kasai, D. F. Kelley and B. S. Rabinovitch\*

Department of Chemistry BG-10, University of Washington  
Seattle, Washington 98195

Abstract

The study of collisional energy transfer between cyclopropane molecules and a seasoned silica wall by the Variable Encounter Method has been extended upward in temperature to 1325 K. The value of  $\langle \Delta E' \rangle$  the average energy of a down-transition upon collision of the substrate with the wall continues to decrease monotonically. The data provide the most accurate and well-documented display of the temperature trend of  $\langle \Delta E' \rangle$  presently available.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

Recently, collisional energy transfer data have been obtained in steady-state homogeneous thermal unimolecular polyatomic systems at temperatures of 773 K<sup>2a</sup>, 973 K<sup>2a</sup> and 1173 K<sup>2b</sup>, with cyclopropane-d<sub>2</sub> and cyclopropane substrates. These studies were made with various monatomic, diatomic and polyatomic bath gases and indicate that the relative collision efficiency,  $\beta_c$ , and the average amount of energy transferred on a downstep transition,  $\langle \Delta E' \rangle$ , both decrease with increase of temperature. Very weak colliders such as helium have been predicted on theoretical grounds<sup>3</sup> to follow a power law,  $\beta_c \propto T^{-n}$ , where  $n > 2$ . Some anomalies in shock tube work at temperatures around 1200 K - 1500 K have previously led to the suggestion that the trend in  $\beta_c$  may extend to significantly higher temperatures.<sup>4,5</sup> By contrast, a recent shock tube study<sup>6</sup> has led to the suggestion that  $\langle \Delta E' \rangle$  increases with rise of temperature for helium bath gas, although contradictory data exists.<sup>2</sup>

A decline of  $\langle \Delta E' \rangle$  with rise of temperature has also been observed<sup>in studies made</sup> by the recently developed Variable Encounter Method (VEM) for a variety of hydrocarbon substrate molecules,<sup>7-10</sup> over the range from above 400 K to above 1100 K. The collision partner in these studies was the seasoned wall surface, assumed to behave like a large polymeric molecule of effectively infinite mass. It is important to extend these approaches well into the high temperature region in order to link more firmly the results at lower temperatures with the shock tube regime. In the work reported here, we have extended the VEM study of the collisional relaxation of vibrational transients in the cyclopropane system up to 1325 K.

The details of the practise and theory of the experimental method are similar to previous studies.<sup>7-10</sup> Average numbers of collisions of substrate molecules per encounter with the reactor surface,  $m$ , were 5.6 and 7.6 in the present work. The purity of the reactant cyclopropane was 99.94%. Analysis of the reaction products was by gas chromatography with a squalane column and f.i.d.

Figure 1 shows experimental plots of the average reaction probability per collision,  $\bar{P}_c(m)$  versus temperature from 1020 K to 1325 K. The data were compared with stochastic calculations using several models for the energy transfer transition probabilities,  $P_{\Delta E}$ ; details are provided in ref. 7. The threshold energy  $E_0$  was taken as  $63.6 \text{ kcal mol}^{-1}$ . The best fit to the data is obtained with a gaussian model:  $P_{\Delta E} = A \exp\{-(\Delta E - \Delta E_{mp})^2 / 2\sigma^2\}$ ; where  $A$  is a normalization constant;  $\Delta E_{mp}$  is the most probable downstep energy transferred (independent of initial energy), and  $\sigma$  was set equal to  $0.7 \Delta E_{mp}$ . Calculated curves were fitted to the data (Fig. 1). Values of  $\langle \Delta E' \rangle$  ( $\text{cm}^{-1}$ ) that fit the data from both reactors at particular temperatures are shown, and the trend of decreasing  $\langle \Delta E' \rangle$  with increasing temperature is seen to continue throughout the temperature range studied. As shown in Fig. 2, these values are almost identical to, but a little less ( $\sim 30 \text{ cm}^{-1}$ ) than those reported for cyclopropane by Flowers et al.<sup>9</sup> in the region of temperature overlap.

As was mentioned elsewhere,<sup>10</sup> the most reliable data tend to be obtained at higher temperatures, where complexities due to surface catalytic effects are minimized. Paradoxically, the work in ref. 9 showed a slight and increasing catalytic side reaction of acetylene production, at and above 1000 K. Careful chemical analysis in the present study revealed no acetylene side reaction. Since both studies give rise to almost identical wall activation efficiencies, the assertion of Flowers, et al. that acetylene production represented a minor, and irrelevant side reaction is borne out.

Incubation times<sup>7</sup>  $\tau_{inc}$  have been defined previously and have been related to the mean first passage time,  $\bar{t}_{fp}$ , by the expression<sup>7</sup>

$$\tau_{inc} = \bar{t}_{fp} - t_{fp}^{ss}$$

where  $t_{fp}^{ss}$  is the steady state mean first passage time and  $\bar{t}_{fp}$  is the mean first passage time for the case of transients,

$$\bar{t}_{fp} = \int_0^{\infty} t P(t) dt ,$$

where  $P(t)$  is the probability that a molecule will cross the reaction threshold  $E_0$ , for the first time, at time  $t$ .  $\tau_{inc}$  also increases with temperature in this study as  $\langle \Delta E' \rangle$  declines, from 13.5 collisions at  $\sim 1040$  K to 18.6 collisions at 1300 K. Incubation times greater than unity signify, of course, that the conventional relative collisional efficiency,  $\beta_c$ , is less than unity, i.e., less than strong collider values.

In conclusion, the data of Figure 2 provide the most accurate illustration presently available of an example of a trend of  $\langle \Delta E' \rangle$  over a wide range of temperatures.

The decrease in  $\langle \Delta E' \rangle$  and  $\beta_c$  with rise of temperature can be explained as follows. A semi-empirical quasi-statistical accommodation model for collisional energy transfer, subject to conservation of angular momentum restrictions, was described some time ago by Lin and Rabinovitch.<sup>3</sup> Oref and Rabinovitch<sup>15</sup> gave a theoretical treatment based on this model. In either version, a decrease in  $\langle \Delta E \rangle$  with increase in temperature is predicted. The treatment of Lin and Rabinovitch took the form for the average of a down-transition,  $\langle \Delta E' \rangle$ ,

$$\langle \Delta E' \rangle = \frac{\sum_{\Delta E = E_{th}}^{E_{tot} - E_0} P(\Delta E) B(E_{th}) [\Delta E - E_{th}]}{\sum P(\Delta E) B(E_{th})} ;$$

where,  $E_{th}$  is the thermal energy of transitional modes (3 to 6, depending on the nature of the collision partner) that are formed in the collision complex;  $E_{tot}$  is the total internal energy of the substrate molecule and the transitional modes;  $E_0$  is the dissociation energy of the collision complex;  $P(\Delta E)$  is the probability of, and  $\Delta E$  is the energy in the transitional modes at the time of decomposition of the complex. The negative temperature behavior arises because

$E_{th}$  increases more strongly with  $T$  than does either the total energy of the collision complex or  $\Delta E$ . In addition, decrease of the residence time of molecules on the surface with increase of temperature further weakens the practical realization of the limiting assertion of quasi-statistical energy relaxation.

We may expect for weak colliders, especially He, that at sufficiently high temperatures, the collision-complex accommodation model<sup>3</sup> may give way to the Landau-Teller-type behavior for polyatomic molecules, as has been found previously (starting at much lower temperatures) for diatomics,<sup>11</sup> and as appears may be likely<sup>6</sup> for triatomics near, or above 2000 K. Whether such alteration of behavior will be seen at practical temperatures for the case of interaction of polyatomic substrates with fairly strong colliders is more problematical.

References

1. Work supported by the Office of Naval Research and the National Science Foundation
- 2.a) I. E. Klein, B. S. Rabinovitch, and K. H. Jung, J. Chem. Phys. 67, 3833 (1977); I. E. Klein and B. S. Rabinovitch, Chem. Phys. 35, 439 (1979); V. V. Krongauz, M. E. Berg, and B. S. Rabinovitch, Chem. Phys. 47, 9 (1980).
- b) E. Kamaratos, J. F. Burkhalter, D. G. Keil, and B. S. Rabinovitch, J. Chem. Phys. 83, 984 (1979).
3. Y. N. Lin and B. S. Rabinovitch, J. Phys. Chem. 74, 3151 (1970); D. C. Tardy and B. S. Rabinovitch, Chem. Rev. 77, 369 (1977).
4. B. S. Rabinovitch, D. G. Keil, J. F. Burkhalter, and G. B. Skinner, Proc. Xth Internat. Shock Tube Symposium, ed. G. Kamimoto, Kyoto, 1975, p. 579.
5. J. Troe, Proc. Xth Internat. Shock Tube Symposium, ed. G. Kamimoto, Kyoto, 1975, p. 29.
6. H. Endo, K. Glanzer, and J. Troe, J. Phys. Chem. 83, 2083 (1979).
7. D. F. Kelley, L. Zalotai, B. D. Barton, and B. S. Rabinovitch, J. Chem. Phys. 71, 538 (1979); D. F. Kelley, L. Zalotai, and B. S. Rabinovitch, Chem. Phys. 46, 379 (1980).
8. M. C. Flowers, F. C. Wolters, B. D. Barton, and B. S. Rabinovitch, Chem. Phys. 47, 189 (1980); M. C. Flowers, F. C. Wolters, D. F. Kelley, and B. S. Rabinovitch, Chem. Phys. Lett. 69, 543 (1980).
9. M. C. Flowers, F. C. Wolters, D. F. Kelley, and B. S. Rabinovitch, Chem. Phys., in press.
10. D. F. Kelley, T. Kasai, and B. S. Rabinovitch, J. Chem. Phys., in press.
11. J. D. Lambert, Vibrational and Rotational Relaxation in Gases, Clarendon Press, Oxford, 1977.
12. I. Oref and B. S. Rabinovitch, Chem. Phys. 26, 385 (1977)

Figure Captions

Figure 1 Plots of experimental values of  $\bar{P}_c(m)$  vs temperature for the two reactors:  $m = 7.6$ ,  $\bullet$  ;  $m = 5.6$ ,  $\blacksquare$  . Also shown are solid curves calculated using a gaussian transition probability model fitted to the  $m = 5.6$  data at each of three values of  $T$ . The  $\langle \Delta E' \rangle$  values ( $\text{cm}^{-1}$ ) were then used to construct the calculated curve for  $m = 7.6$ .

Figure 2 Calculated plots of  $\langle \Delta E' \rangle$  versus temperature for cyclopropane: data of Flowers et al. (ref. 9), - - - ; present data, — .






Fig. 6.40.57.

DATE  
FILMED  
-8