

AD-A098 905

NAVAL OCEAN RESEARCH AND DEVELOPMENT ACTIVITY NSTL S--ETC F/G 8/3
MEASURED FLOW IN ST. VINCENT AND GRENADA PASSAGES IN 1977.(U)

JUN 80 P A MAZEIKA, D A BURNS, T A KINDER


UNCLASSIFIED

NORDA-TN-62

NL

1 of 1
AD A
0098905

END
DATE
FILED
6-81
DTIC



MICROCOPY RESOLUTION TEST CHART
NBS 1963-A

LEVEL II

65 (12)

Naval Ocean Research
and Development Activity

NSTL Station, Mississippi 39529




NORDA Technical Note 62

Measured Flow in St. Vincent and Grenada Passages in 1977

5-1413

AD A098300



DIC FILE COPY

Paul A. Mazeika
Donald A. Burns
Thomas H. Kinder

DTIC
ELECTE
MAY 14 1981
S D E

Oceanography Division
Ocean Science and Technology Laboratory

JUNE 1980

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

81 5 14 057

ABSTRACT

Between January and November 1977 flow was measured with ten current meters in St. Vincent and Grenada Passages. Mean scalar speeds exceeded 23 cm/sec at all instruments and flow was predominantly westward. Subinertial variability calculated using wide band (0.0156 CPH) spectral estimates was large, amounting to 14%-77% of the individual record variances. All records showed changes in low frequency flow, such as abrupt changes in direction, stagnation (period of low flow), or 360° rotations in direction. Spectra showed peaks near 12-13 days in near-bottom records from St. Vincent passage, and peaks between 20 and 70 days in the other records.

Strong tidal signals were also found in the velocity records. Between 15% and 74% of the individual record variances were in diurnal, semidiurnal, or harmonic frequency bands. The semidiurnal frequency band contained from 11% - 67% of the record variances. Tidal harmonics were also clearly evident, especially at the two near-bottom instruments deployed sequentially in St. Vincent Passage, where the first five harmonics of the semidiurnal frequency accounted for 27% - 35% of the variance.

Each instrument recorded temperature, and most of the individual record variance was either subinertial (8% - 86%) or semidiurnal (4% - 82%). Together, these frequency bands accounted for 69% of the variance. Tidal harmonics accounted for 2% - 25% of the temperature variance.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Original
A	

ACKNOWLEDGMENTS

We wish to acknowledge the successful deployment and retrieval of current meter arrays by Louis A. Banchero. Captain John W. Arens and the crew of USNS BARTLETT provided at-sea support.

CONTENTS

	Page
List of Illustrations	iv
List of Tables	v
I. Introduction	1
II. Historical Flow Measurements in the Passages	1
III. Low Frequency Current Fluctuations in the Gulf of Mexico and Straits of Florida	3
IV. Presentation of Results	5
V. References	13

ILLUSTRATIONS

	Page
Figure 1. Caribbean Sea	14
Figure 2a. Temperature in St. Vincent Passage, 3 Feb 77	15
Figure 2b. Salinity in St. Vincent Passage, 3 Feb 77	16
Figure 3a. Temperature in Grenada Passage, 30 Jan 77	17
Figure 3b. Salinity in Grenada Passage, 30 Jan 77	18
Figure 3c. Temperature in Grenada Passage, 18 Feb 77	19
Figure 3d. Salinity in Grenada Passage, 18 Feb 77	20
Figure 4. Seasonal speed histograms, St. Vincent Passage	21
Figure 5. Seasonal speed histogram, Grenada Passage	22
Figure 6a. Progressive vector diagram at 105 m in St. Vincent Passage, array A	23
Figure 6b. Progressive vector diagram at 380 m in St. Vincent Passage, array A	24
Figure 6c. Progressive vector diagram at 880 m in St. Vincent Passage, array A	25
Figure 7a. Progressive vector diagram at 380 m in St. Vincent Passage, array B	26
Figure 7b. Progressive vector diagram at 820 m in St. Vincent Passage, array B	27
Figure 8a. Progressive vector diagram at 60 m in Grenada Passage	28
Figure 8b. Progressive vector diagram at 110 m in Grenada Passage	29
Figure 8c. Progressive vector diagram at 320 m in Grenada Passage	30
Figure 8d. Progressive vector diagram at 600 m in Grenada Passage	31
Figure 9. Changes in direction	32
Figure 10. Low frequency current and temperature, St. Vincent, array A	33
Figure 11. Low frequency current and temperature, St. Vincent, array B	34
Figure 12. Low frequency current and temperature, Grenada	35
Figure 13. Unfiltered spectra, St. Vincent array A, 105 m	36

	Page
Figure 14. Unfiltered spectra, St. Vincent, array A, 380 m	37
Figure 15. Unfiltered spectra, St. Vincent, array A, 880 m	38
Figure 16. Unfiltered spectra, St. Vincent, array B, 380 m	39
Figure 17. Unfiltered spectra, St. Vincent, array B, 820 m	40
Figure 18. Unfiltered spectra, Grenada, 60 m	41
Figure 19. Unfiltered spectra, Grenada, 110 m	42
Figure 20. Unfiltered spectra, Grenada, 320 m	43
Figure 21. Unfiltered spectra, Grenada, 600 m	44
Figure 22. Low frequency spectra of flow components, St. Vincent, array A	45
Figure 23. Low frequency spectra of flow components, St. Vincent, array B	46
Figure 24. Low frequency spectra of flow components, Grenada	47
Figure 25. Coherence of u components, St. Vincent, array A	48
Figure 26. Coherence of v components, St. Vincent, array A	49
Figure 27. Coherence of u and v components, St. Vincent, array B	50
Figure 28. Coherence of u components, Grenada	51
Figure 29. Coherence of v components, Grenada	52

TABLES

Table 1. Current meter array description	2
Table 2. Current meter observations in the Antilles Passages	4
Table 3. Percent frequency distribution of current speed	7
Table 4. Percent frequency distribution of temperature from current meters	8
Table 5. Major direction variations of flow	9
Table 6. Mean resultant flow and tidal components	10
Table 7. Variance distribution	11

I. INTRODUCTION

The circulations of the subtropical North Atlantic, the Caribbean Sea, and the Gulf of Mexico are highly interrelated. The net mass inflow into the Caribbean through various passages must balance the net outflow through the Yucatan Channel. The water balance which occurs in the Gulf of Mexico is relatively simple, since there is one channel for net inflow (Yucatan Channel) and one channel for net outflow (Straits of Florida). The Caribbean portion of this coupled system is more complex, since many passages have considerable variability of inflow and outflow (Fig. 1). The factors causing fluctuations of transport are little known and patterns of variability are only scantily explored. Simultaneous meteorological, hydrographic, and current meter data obtained continuously over several years throughout the Caribbean and all passages, are required for a comprehensive understanding of this system. The technical and financial difficulties in obtaining such synoptic data, however, are obvious. The results of the investigation presented in this study include only two of the southernmost Lesser Antilles passages, which may carry a high percentage of the net inflow from the Atlantic.

The current meter data were obtained during January–November 1977 in St. Vincent and Grenada Passages. Locations of the current meter arrays, depths of current meters, durations of records, and other information are shown in Figure 1 and Table 1.

All current meters were AMF Mod 610 Vector Averaging Current Meters, which recorded speed, direction, and temperature averaged over 15 minutes. Specifications of resolution, range, and accuracy of the current meters are described in the manufacturer's manual (AMF Electrical Products Development Division, 1973). The current meter arrays were taut line moorings similar to the mooring described and illustrated by Banchemo (1973). Bottom depth, before anchoring the arrays, was carefully explored by about ten crisscrossings of each channel, and vertical accuracy is estimated as 10–12 m.

The purpose of this investigation was to obtain flow characteristics during the entire period of the current meter's operational capability (approximately nine months). The results, besides amplifying knowledge of flow patterns and variability, are applicable also for rudimentary modeling and for guiding further investigations. Although the observations extended over two main seasons (winter and summer), the seasonal variability cannot be interpreted with only nine months of data. Nevertheless, some seasonal differences in flow regimes appeared. The main emphasis was to obtain long, continuous records for estimating variability at frequencies higher than seasonal. In addition, the current meter arrays were deployed at the beginning of a hydrographic survey in the passages and adjacent waters extending between the southern Lesser Antilles and Tobago–Barbados longitudes in January–February 1977. The results of this survey (Mazeika et al., 1980) show a complex mesoscale circulation just east of the southern Lesser Antilles.

II. HISTORICAL FLOW MEASUREMENTS IN THE PASSAGES

Previous current measurements taken in the Lesser Antilles and in some of the northern Caribbean passages have provided important information. They can be compared with the variability in other areas of this system and may eventually be used to design more complete investigations.

Stalcup et al. (1971) reported results obtained from four current meters moored in the Grenada Passage, and Burns and Carr (1975) analyzed 36 current meter

TABLE 1. CURRENT METER ARRAY DESCRIPTION (1)

PASSAGE	LAT (N)	LONG (N)	WATER DEPTH (m)	CH DEPTH (m)	DATA START	DATA END	DURATION OF DATA (DAYS)	INERTIAL PERIOD (HOURS)
St. Vincent								
A	13°25.4'	61°03.1'	915	105	30 Jan 77	24 May 77	114	
				380		24 May 77	114	51.69 (1.935 x 10 ⁻² cph)
B(2)	13°25'	61°03'	850	105(3)	26 May 77	11 July 77	46	
				380		9 Nov 77	167	51.7 (1.93 x 10 ⁻² cph)
Grenada								
	11°43.5'	61°55.5'	750	60(4)		25 Sept 77	238	
				820		7 Nov 77	165	
				110(4)	30 Jan 77	27 Apr 77	87	59.05 (1.693 x 10 ⁻² cph)
				320		6 Nov 77	280	
				600		6 Nov 77	280	

(1) The sampling rate was four samples/hour.

(2) Position of current meter arrays A and B were approximately the same. The direction of flow at the near bottom current meter of array B, however, was inconsistent with that of array A. The deepest current meter of array B was 50 m shallower and the bottom depth 75 m shallower than at A. Apparently the positions differed, although both arrays were in the main channel. The time lapse between end of A and start at B was about 26 hours.

(3) The current meter at 105 m of array B ran for only 46 days and the raw data plot indicated that the operation was not continuous.

(4) Comparison of current meter temperature records and concurrent hydrographic profiles suggested that the 60 m and 110 m Great Passage instruments were about 70 m deeper than their nominal depths.

records obtained between Venezuela and Hispaniola. Twenty-two of these records were from various passages, while the rest were obtained elsewhere within the Caribbean. Smith (1974) reported data taken from twelve current meters, with one array of four current meters located near Jungfern Passage. A summary of the basic information from these three sources encompassed 29 measurements (Table 2). These records ranged between 14 and 47 days in length, with a mean of 26.7 days. The observation depths ranged between 45 and 1509 m. The reasons for selecting depths were seldom explained and are often difficult to rationalize. The mean velocities estimated from progressive vector diagrams were between 1 and 2 knots (kn) (1 kn = 51.4 cm/sec) in the upper layers, but they were generally moderate or low in the layer between the salinity maximum and minimum cores. Two current meters were located near the salinity maximum (at 100 m in Jungfern and at 85 m in Mona Passages), but both measured rather weak flow. Mean velocities at three current meters below the salinity minimum were less than 8 cm/sec. A majority of current records indicated one or more significant, or even sharp, changes in the direction of flow. No apparent periodicity appears in this variation of direction. Some records indicated considerable persistence in direction, while others changed direction randomly over intervals of 2 to 37 days.

These records were also variable at tidal frequencies. Energy spectra of the historical current meter data in Grenada Passage showed the semidiurnal tides at all measured depths. In St. Vincent and St. Lucia Passages, semidiurnal tides dominated in the upper layers, while near the bottom semidiurnal mixed tides were present that had a significant diurnal component. Most other current records showed that mixed semidiurnal tides had a considerable diurnal component of energy, except for the Guadalupe/Antigua and Jungfern Passages where the tide is predominantly diurnal and has a considerable semidiurnal component.

Additionally the presence of an overtide at about a six-hour period was reported for almost all current meter data except that from Grenada and Anegada Passages. Significant energy density was shown in at least half of the records at about an eight-hour period. In Mona Passage a 50-hour period was noticeable. Pronounced current fluctuations at the inertial period were present in Guadalupe/Antigua, Montserrat/Antigua and Mona Passages. Low frequency 360° turns of the current were observed in several cases at various depths, with periods between 5 and 12 days.

III. LOW FREQUENCY CURRENT FLUCTUATIONS IN THE GULF OF MEXICO AND THE STRAITS OF FLORIDA

Fluctuations of a four- to five-day period were reported in Yucatan Strait by Hansen and Molinari (1979) from one-month-long current meter data taken at 1977 m (15 m above the bottom). Wunsch and Wimbush (1977) showed large nontidal current variability in Florida Strait at 5 and 14 day periods. They related this variability to corresponding oscillations in wind forcing. Düing et al. (1977) used two years' data of current and temperature measurements in the Florida current off Miami to show that low frequency energy maxima occurred at periods of 2 to 4, 4 to 5, and 8 to 25 days. Fluctuations, especially in the 10-13 day period band, were coherent with these maxima, and the wind stress curl led current by three days. Kielman and Düing (1974), analyzing a 50-day current meter record off Miami, showed persistent current fluctuations at periods of about 5-6 days; the downstream component was six times larger than the cross-stream component and led by about one-quarter period. They found that mesoscale wind forcing might have been important for current fluctuations, but that local and large-scale wind effects were not significant. Niiler (1976) showed fluctuations at about 10-20 day periods

TABLE 2. CURRENT METER OBSERVATIONS IN THE ANTILLES PASSAGES (1)

PASSAGE	LOCATION LAT. LONG.	MONTH, YEAR	OBSERV. LENGTH DAYS	OBSERV. DEPTH (m)	WATER DEPTH (m)	MEAN VELOCITY CM/SEC (2)	MEAN (3) DIRECTION	TYPE (4)
Grenada	11°36'N; 61°53'W	Mar-Apr 70	37	423	623	14	W	SM
	"	"	"	623	"	19	Nw	SM
	11°31'N; 61°53'W	Mar-Apr 70	"	225	477	10	NE 24; W 15	SM
	"	"	"	427	"	21	NE 26; W 11	SM
	11°08'N; 61°35'W	Aug 72	15	130	140	10	NE	SM
St. Vincent	11°51'N; 61°46'W	May 72	24	45	640	47	SW	SM
	13°30'N; 61°08'W	June 72	22	45	1600	90	NW	SM
	"	"	"	245	"	14	NW 20; NE 1	SM
	"	"	"	1590	"	3	SE 1; NW 21	SM
	13°38'N; 61°00'W	July 72	23	45	600	61	NW	SM
St. Lucia	"	"	"	590	"	9	Nw	SM
	14°19'N; 60°55'W	July 72	14	45	1030	49	NW	SM
	"	"	"	245	"	28	SSW	SM
Bermuda	"	"	22	1010	"	7	W 8; SW 14	SM
	15°02'N; 61°16'W	July 72	22	290	2060	11	SW 14; S 8	SM
Guadelupe- Antigua	16°36'N; 61°47'W	June 72	22	460	1045	5	SW	SM
Monserrat- Antigua	16°54'N; 62°03'W	June 72	18	260	760	6	N 8; WNW 10	SM
	"	"	22	745	"	5	WNW	SM
St. Martin St. Christopher	17°45'N; 63°04'W	June-July 72	22	245	735	6	NW 3; SW 8; NNW 8; SW 3	SM
Anegada	18°32'N; 64°02'W	June-July 72	22	280	2075	11	NE 17; NW 5	SM
Jungfern (5)	17°38'N; 65°12'W	Apr-May 74	47	100	1564	10	SE	SM
	"	"	"	450	"	8	NW 21; SE 26	SM
	"	"	"	800	"	4	N 10; NE 37	SM
	"	"	14	1534	"	6	NE	SM
Mona	18°11'N; 67°41'W	Sept-Oct 72	29	285	485	10	NE	SM
	18°19'N; 68°00'W	Jan 72	28	85	575	15	SW 10; SE 12	SM
	"	"	"	560	"	4	W 12; NW 11	SM
	18°31'N; 67°48'W	Oct 72	28	350	625	11	W 20; NNW 8	SM
"	"	"	615	"	4	NE	SM	

NOTES:

- (1) The first four current meters (Geodyne #850) listed in the Table were by WHOI; sampling rate 15 minutes. The rest of the current meters (Geodyne A101) were by NAVOCEANO; sampling rate 10 minutes.
- (2) Estimated mean velocity from progressive vector diagrams using 24 h mean data for WHOI and 12 h mean data for NAVOCEANO.
- (3) General direction of progressive vector diagrams. When flow was in more than one direction, the attached figure indicates the duration of each episode.
- (4) S = Semidiurnal,
SM = Semidiurnal mixed,
M = Diurnal mixed.
- (5) about 33 km to the west of the deepest part of Jungfern channel. The deepest current meter was below the 100 m level.

in the longshore currents on the western Florida Shelf and reported spectral properties similar to those of currents on the eastern Florida Shelf. Lee and Mayer (1977) found significant coherence between north-south currents at the shelf break (30 m isobath) off southwestern Florida and the east-west wind at a nine-day period, with the wind leading by about 90°. In the Florida current, meandering and fluctuations at periods of 5-7 days are persistent, as shown by Niiler (1975) in his extensive review of data, theory, and numerical models.

Similar forcing and interaction processes may operate in the eastern part of the Caribbean, which has not been as extensively studied as the Gulf of Mexico and the Straits of Florida. Meandering of trade winds at four- to five-day periods has been shown in the Caribbean area by Riehl (1954). Mofjeld and Wimbush (1977) showed high coherence between bottom pressure and local wind oscillations at a five-day period with a 4.5 hour phase lag in the eastern Caribbean.

Effects of local wind forcing on flow variability in the relatively narrow and shallow Lesser Antilles Passages may not be significant, but large-scale wind field fluctuations on both sides of the islands may be important. Other factors may also be considered in the passages and in the eastern Caribbean. It has been shown recently by Mazeika et al. (1980) and by Mazeika (1973) that intense mesoscale eddies, which may affect low frequency flow variability farther west, occur east of the Lesser Antilles. Thus, the temporal variability and occurrence of subinertial energy peaks in various frequency bands at different locations in the eastern Caribbean may result from a complex interaction of fluctuations in intensity and direction of wind forcing at various frequencies. Internal factors may also play a role.

IV. PRESENTATION OF RESULTS

Current meter data presented in this study are of much longer duration than any of the previous observations summarized in Table 1. The depths for current meters were selected to measure flow in the layer of strong baroclinicity which is near and above the salinity maximum, to measure flow in a thick intermediate layer between the salinity maximum and minimum cores, and to measure flow in the near-bottom layer. Vertical distributions of temperature and salinity, the location of current meters, and the mean flow for the day of hydrographic measurements in St. Vincent Passage are shown in Figures 2a and 2b. The temperature and salinity distributions are shown for Grenada Passage in Figure 3a and 3b one day after deployment of the current meter array and Figure 3c and 3d for three weeks later. Considerable change in the temperature and salinity distributions is noticeable between the two sections.

Most of the records contained few suspicious data, but there were some exceptions. In St. Vincent Passage array B was placed close to the position of array A, and began operation 26 hours after array A was recovered (Table 1). The deep current meters on these two arrays, however, showed mean flows in opposite directions and much different amplitudes of east-west semidiurnal tides (Table 6). We have no other reason to suspect these records: the abrupt differences may have been caused by local topography (both instruments were within 30 m of bottom). The 105 m record of array B was short (46 days) and noisy, so we have excluded it from most discussions. The Grenada records at 110 m and 320 m were noisier than the remainder of the records, but appear quite usable for our purposes. Additionally, an array was deployed in St. Lucia Passage, but was not recovered.

Tables 3 and 4 show the percent frequency distributions of speed and temperature, and their means. A high percentage of speeds (Table 3) in the range of 30-40 cm/sec occurred at all depths, but only the shallowest current meters showed high percentages in the range of 70-80 cm/sec. Percent frequency of temperature (Table 4) shows a decreasing range of temperature with depth, as usual. Discrepancies occurred in the temperature records for the Grenada Passage at 60 m and 110 m between current meter and CTD data. Temperature differences indicate about 70 m greater depth than nominal for the two current meters.

Seasonal frequency distributions of speeds are shown in Figure 4 for St. Vincent Passage and in Figure 5 for Grenada Passage. There are changes in the frequency distribution between winter and summer that may indicate some trends, but no conclusions on seasonal variability can be made at this point.

The most conspicuous events in direction of flow are summarized in Table 5 and can be compared with the progressive vector diagrams presented in Figures 6 through 8. Typical examples of change in direction, circular rotation, and stagnation (speed < 3 cm/sec) are shown in Figure 9 by segments of progressive vector diagrams with points of two-hour averaged speed and direction. Stagnation at 105 m (array A, St. Vincent Passage) on March 17-25 (Table 5) actually occurred as a slow cyclonic circling (about 5 circles) visible on a large-scale, two-hour average progressive vector diagram (not shown). Also the stagnation that occurred during July 2-9 at 600 m in Grenada Passage was actually several slow anticyclonic rotations. Significant low frequency changes in current direction occurred with these data even more often than with the previously described historical data. Persistence in the same direction ranged between 3 and 130 days. Frequencies of long-period rotations with these data are similar to the historical data described above, ranging from 3 to 18 days. Direction changes were frequently uncorrelated between various depths at the same array and appeared as random events. Such changes may suggest current reversals throughout the passage, but may also be meanders or translations of narrow currents. Obviously a single current meter array in a passage is not sufficient to detect the cause for direction change and the mechanism by which flow is reorganized. Extensive synoptic data may be required on both sides of the islands and in the passages to understand the nature of the forcing, the process of this change, and the effects on the hydrographic distribution.

Table 6 lists resultant (vector mean) speed, direction, and persistence for each record. Persistence is defined as the ratio of resultant speed to scalar mean speed, expressed in percent, and is a measure of the directionality of the flow. For example, a unidirectional flow would have a persistence of 100%. Scalar mean speeds ranged from 23.5 cm/sec to 51.5 cm/sec, with most in the range of 20 to 30 cm/sec. Resultant speeds varied from 8.5 cm/sec to 45.8 cm/sec, with most in the range from 14 cm/sec to 22 cm/sec. The persistences were high, varying from 36% to 91%. Only one record had a persistence below 49%. In addition, Table 6 shows major tidal flow constituents. A least-square fit of a periodic trial function, using the Gauss-Seidel technique, was applied to the u and v current components. The maximum number of points that can be handled by the program is 16,280 (170 days). The purpose was to obtain a first look at tidal signatures; therefore, no block averaging of data was attempted.

Daily vectors are shown in Figures 10 through 12. In the two near-bottom St. Vincent records a clear oscillation appears which has a peak-to-peak amplitude of about 20 to 30 cm/sec and a period of about 10 to 15 days. At the Grenada mooring, the 60 m and 600 m records had longer period fluctuations. These

TABLE 3. PERCENT FREQUENCY DISTRIBUTION OF CURRENT SPEED

PASSAGE	DEPTH (m)	SPEED CM/SEC										MEAN SPEED CM/SEC
		0	10	20	30	40	50	60	70	80	>80	
St. Vincent												
A	105	1.4	5.8	9.7	15.1	15.6	15.2	16.1	12.7	8.4	8.4	51.5
	380	16.6	31.6	21.5	17.0	9.4	3.3	0.5	0.1			23.3
	880	6.5	16.0	14.7	17.3	19.4	14.6	8.0	3.1	0.4		37.2
F	380	12.5	27.7	19.5	16.6	12.3	8.4	2.6	0.4			27.6
	820	13.0	30.8	27.9	17.7	7.4	2.5	0.6	0.1			23.6
Grenada	60	6.4	18.4	20.8	19.3	13.6	9.6	7.4	2.7	1.8		34.8*
	110	26.3	22.1	16.6	11.3	7.1	4.5	3.8	2.6	5.5		28.1
	320	15.0	25.6	23.6	17.2	10.3	6.3	1.7	0.3			25.8
	600	17.5	25.2	26.0	20.2	8.4	2.4	0.3				23.5

TABLE 4. PERCENT FREQUENCY DISTRIBUTION OF TEMPERATURE FROM CURRENT METERS

PASSAGE	DEPTH (m)	°C													MEAN			
		4	6	8	10	12	14	16	18	20	22	24	26	28				
St. Vincent																		
"	105						0.16	2.53	13.18	23.86	42.41	17.56	0.30					22.19
	380		2.86		60.22	36.46	0.46											11.70
	880	87.33	12.67															5.52
	380		0.92		43.93	53.58	1.57											12.10
	820	74.76	25.24															5.61
St. Vincent	60 (1)				0.01	0.53	4.06	36.65	45.31	13.29	0.15							18.34
	110 (1)				0.42	12.04	60.67	21.79	5.08									15.39
	320		1.46		55.94	42.46	0.14											11.88
	600		95.76	4.24														7.11

Temperature at 60 m was about 4° and at 110 m -3° lower than CTD temperatures at the hydrographic stations on both sides of the current meter array. The current meters were apparently about 70 m deeper than indicated.

TABLE 5. MAJOR DIRECTION VARIATIONS OF FLOW

PASSAGE	CURRENT METER DEPTH (m)	DIRECTION OF FLOW (1)	STAGNATION (2) (DATES)	ROTATION	
				DATES	DURATION (DAYS) CHARACTERISTICS
St. Vincent					
A	105	WNW 17; NW 38; NNW 25; W 5; NW 2	March 17-25	March 17-May 26	40 Cyclonic; 4 full rotations in sequence
	380	NNW 14; NW 32	March 6-11	March 1-9	8 Anticyclonic; full circle
	880	NNW 35; NW 48; E 5; NW 5			
B	380	NW 116; S 5; W 5	Sept 25-Oct 2 Nov 2-5	Oct 24-28	4 Anticyclonic; semicircle
	820	NE	June 13-16 Sept 24-30		
Grenada	60	W 24; WSW 130; SW 16; WSW 40		July 7-16 Aug 1-9	9 Anticyclonic; full circle 8 Cyclonic; semicircle
	110	N 20; NE 58		Feb 27-March 11 April 18-May 2	12 8 Cyclonic; semicircle Cyclonic; semicircle
	320	E 4; WNW 13; SW 10; NW 35; NNW 2 NW 17; NNW 47	Feb 16-19 Feb 22-26 March 1-7	March 13-17 April 18-May 2 July 7-25	4 14 18 Cyclonic; semicircle Anticyclonic; full circle Anticyclonic; full circle
	600	WSW	July 13-14 Aug 10-18 Oct 11-14	April 22-25 May 5-13	3 8 Cyclonic; full circle Cyclonic; full circle

(1) Near general directions estimated from progressive vector diagrams. Significant changes in direction are shown in sequence with adjacent figures indicating number of days in the corresponding direction.

(2) Stagnation is defined as an extended period (2 or more days) when the current speeds are very low or zero.

fluctuations were less regular and had fewer realizations than the shorter period oscillations in St. Vincent Passage. The peak-to-peak amplitude was about 60 cm/sec at 60 m depth and about 40 cm/sec at 600 m depth. The period of these fluctuations appeared to be from 50 to 90 days. Similar fluctuations may have existed in the St. Vincent records, but the length (167 days or less) was too short to define clearly fluctuations that were 50 to 90 days long.

Spectra were computed on all unfiltered records (except for array B, 105 m) using the BMDX92 package developed by the UCLA Health Science Computing Facility. For all records the mean and linear trends were removed before computing the spectra. A few large and isolated spikes were removed from the temperature records, and beginning and ending transients were removed from the temperature and velocity records; but the records were not otherwise manipulated. Figures 13-21 show the spectra of velocity (east-west and north-south components) and temperature, and Table 7 lists the variance distributions from these calculations. In all velocity records a considerable variance existed at the lowest frequency estimates (up to 77% of the variance, see Table 7), a prominent semidiurnal tidal peak (11% to 67% of the variance), and the energetic peaks at tidal harmonic frequencies (up to 35% of the variance), especially in St. Vincent Passage. Temperature spectra showed similar characteristics: considerable low frequency variance (up to 86%), prominent semidiurnal peaks (4% to 82%), and energetic tidal harmonics (up to 25%). With these rather wide-band estimates (0.0156 CPH), neither the diurnal tide nor inertial oscillations show as distinct peaks. Tidal frequencies were eliminated to investigate low frequency variability. Spectra were computed using the BMDX routines after low-pass filtering with a cutoff period of about 48 hours (Figs. 22, 23, and 24). Both near-bottom records from St. Vincent show peaks near a 12-13 day period (0.0033 CPH), corresponding to the oscillations seen in the daily vector diagrams (Figs. 10-12). Less pronounced peaks are shown in the records from shallower depths, corresponding to periods near 35 or 70 days (about 1.2×10^{-3} CPH or 6×10^{-4} CPH). The Grenada Passage records had a broad peak near 70 days (6×10^{-4} CPH) and a possible weak peak near 26 days (2×10^{-3} CPH). Figures 25 through 29 show coherences of u and v components between various depths at the three current meter arrays.

TABLE 7. VARIANCE DISTRIBUTION (1)

CURRENT METER			FLOW (2)					TEMPERATURE (2)				
DAYS			TOTAL VARIANCE (CM/SEC) ²	% Low Frequency	% Diurnal	% Semidiurnal	% Harmonics	TOTAL VARIANCE (°C) ²	% Low Frequency	% Diurnal	% Semidiurnal	% Harmonics
	St. Vincent											
A	105 m	114	959	64.1	1.3	23.0	4.3	4.07	63.7	0.5	78.9	1.7
	380 m	114	608	30.2	1.0	60.1	3.7	0.77	38.1	1.5	39.7	11.7
	880 m	112	1356	22.9	0.6	40.3	27.2	0.13	13.6	0.5	55.5	38.2
B												
	380 m	167	499	19.2	1.5	67.2	1.7	0.64	35.7	2.2	36.6	11.7
	820 m	165	482	14.7	0.9	32.5	35.1	0.29	8.2	0.4	82.1	1.3
Grenada												
	60 m	238	600	77.0	1.7	12.9	2.0	1.68	62.0	1.7	15.4	1.7
	110 m	87	1245	44.5	4.5	11.2	7.4	1.3	79.1	1.4	13.5	1.6
	320 m	280	751	60.0	4.1	12.3	5.3	0.65	78.1	1.7	8.2	1.7
	600 m	280	267	69.9	1.3	25.0	0.5	0.21	70.4	1.1	4.2	1.1

(1) As defined in unfiltered spectra.

(2) Low: Two lowest estimates.

Diurnal: Estimate centered on 0.031 cph
 Semidiurnal: Two estimates 0.078, 0.094 cph
 Harmonics: sum of 1st 0.16 , 0.17 cph
 2nd 0.23 , 0.25 cph
 3rd 0.31 , 0.33 cph
 4th 0.39 , 0.40 cph
 5th 0.48 , 0.50 cph
 Width of estimates: 0.0156 cph

V. REFERENCES

AMF Electrical Products Development Division (1973). Vector Averaging Current Meter, Model 610A. AMF Incorporated, Herndon, Virginia.

Banchero, L. A. (1973). Results of Current Observations Church Gabbro. Naval Oceanographic Office, Technical Note 7300-2-73.

Burns, D. A. and M. Carr (1975). Current Meter Data Report for the Eastern Part of the Caribbean Sea. Naval Oceanographic Office, Technical Note 6110-6-75.

Düing, W. D., C. N. K. Mooers, and T. N. Lee (1977). Low-frequency Variability in the Florida Current and Relations to Atmospheric Forcing from 1972 to 1974. J. Marine Res., 35, 129-161.

Hansen, D. V. and R. L. Molinari (1979). Deep Currents in the Yucatan Strait. J. Geophys. Res., 84, 359-362.

Kielman, J. and W. Düing (1974). Tidal and Sub-inertial Fluctuations in the Florida Current. J. Phys. Oceanog., 4, 227-236.

Lee, T. and D. A. Mayer (1977). Low Frequency Current Variability and Spin-off Eddies Along the Shelf Off Southeast Florida. J. Marine Res., 35, 193-220.

Mazeika, P. A., D. A. Burns, and T. H. Kinder (1980). Mesoscale Circulation East of the Southern Lesser Antilles. J. Geophys. Res., 85, 2743-2758.

Mazeika, P. A. (1973). Circulation and Water Masses East of Lesser Antilles. Deutsche Hydrog. Zeit. 26, 49-73.

Mofjeld, H. O. and M. Wimbush (1977). Bottom Pressure Observations in the Gulf of Mexico and Caribbean Sea. Deep Sea Res., 24, 987-1004.

Niiler, P. P. (1976). Observations of Low-frequency Currents on the West Florida Continental Shelf. Memoires Societe Royale des Sciences de Liege, 6^e Serie, Tome X, 331-358.

Niiler, P. P. (1975). Variability in Western Boundary Currents. In: Numerical Models of Ocean Circulation. National Academy of Sciences, Washington, D. C., p. 216-236.

Riehl, H. (1954). Tropical Meteorology. McGraw-Hill Book Co., Inc., New York p. 392.

Smith, D. R. (1974). Results of Current Observations Caribbean Operations. Naval Oceanographic Office, Technical Note 6110-1-74.

Stalcup, M. C., W. G. Metcalf, and M. Zemanovic (1971). Current Measurements in the Lesser Antilles. Woods Hole Oceanographic Institution, Ref. 71-51, Technical Report (unpublished manuscript).

Wunsch, C., and M. Wimbush (1977). Simultaneous Pressure, Velocity and Temperature Measurements in the Florida Straits. J. Marine Res., 35, 75-104.




Figure 2a. Vertical temperature ($^{\circ}\text{C}$) distribution in St. Vincent Passage. Circled crosses show current meter locations. The mean velocities and directions shown at each current meter were computed for the day of hydrographic data (3 February, 1977; 5 days after deployment of the current meter array). The main channel is on the left (south side). The trench on the right side is a canyon in the western end of the passage, while on the eastern end the bottom depth is less than 400 m.

STATION NUMBER	17	18	19	20
	5.64	5.63	5.64	5.82




Figure 2b. Vertical salinity (‰) distribution (less 30 ‰) in St. Vincent passage, 3 February 1977.




Figure 3a. Vertical temperature ($^{\circ}\text{C}$) distribution in Grenada passage, 30 January 1977. Circled crosses show current meter locations. The mean velocities and directions shown at each current meter were computed for the day of hydrographic data (one day after deployment of current meter array).




Figure 3b. Vertical salinity (‰) distribution (less 30‰) in Grenada Passage, 30 January 1977.




Figure 3c. Vertical temperature (OC) distribution in Grenada Passage, 18 February 1977.







Figure 3d. Vertical salinity (‰) distribution (less 30‰) in Grenada Passage, 18 February 1977. Circled crosses show current meter locations. The mean velocities and directions shown at each current meter were computed for the day of hydrographic data (20 days after deployment of current meter array).

WINTER (February, March, April)



SUMMER (July, August, September)





Figure 4. Seasonal (winter and summer) speed histogram in St. Vincent Passage.

WINTER (February, March, April)




Figure 5. Seasonal (winter and summer) speed histogram in Grenada Passage.

ST. VINCENT PASSAGE A 105M




Figure 6a. Progressive vector diagram at 105 m in St. Vincent passage (current meter array A, see Table 1). Each point is a 24 hour resultant vector. S indicates the start of a diagram; adjacent is the first continuation. The second continuation is a short segment in the upper right corner. Both continuations are northwestward.




Figure 6b. Progressive vector diagram at 380 m in St. Vincent Passage (array A).

ST. VINCENT PASSAGE A 880M




Figure 6c. Progressive vector diagram at 880 m in St. Vincent Passage (array A).

ST. VINCENT PASSAGE B 380M



Figure 7a. Progressive vector diagram at 380 m in St. Vincent Passage (array B; see Table 1). The upper segment is a continuation westward.

GRENADA PASSAGE 60M




Figure 8a. Progressive vector diagram at 60 m in Grenada Passage (Table 1).
Continuations are in sequence from the top to bottom, flowing westward.

GRENADA PASSAGE 110M




Figure 8b. Progressive vector diagram at 110 m in Grenada Passage.

GRENADA PASSAGE 320M




Figure 8c. Progressive vector diagram at 320 m in Grenada Passage. The continuations at the left side flow northward.

GRENADA PASSAGE 600M




Figure 8d. Progressive vector diagram at 600 m in Grenada Passage. Continuations are in sequence from top to bottom, flowing westward.




Figure 9. Segments of progressive vector diagrams illustrating changes in direction. Each point represents a 2 hour average.
upper: St. Vincent, 380, 28 Oct-3 Nov
middle: St. Vincent, 880 m, 16 Feb-28 March
lower: Grenada, 320 m, 30 Jan-21 March

ST. VINCENT PASSAGE A




Figure 10. Daily vectors of flow and temperature (continuous line) from low-pass filtered data, St. Vincent Passage, array A (see Table 1). The number to the left of the temperature scale is the record mean.




Figure 11. Daily vectors of flow and temperature (continuous line) from low-pass filtered data, St. Vincent Passage, array B (see Table 1). The number to the left of the temperature scale is the record mean.

GRENADA PASSAGE




Figure 12. Daily vectors of flow and temperature (continuous line) from low-pass filtered data, Grenada Passage. The number to the left of the temperature scale is the record mean.

13-25.4N 61-03.1W 105M




Figure 13. Variance spectra of flow components E (zonal) and N (meridional), and temperature (T^2/cph) at 105 m in St. Vincent Passage (array A, see Table 1). Data were not filtered.

13-25.4N 61-03.1W 380M




Figure 14. Variance spectra of flow components and temperature (T^2/cph), at 380 m in St. Vincent Passage (array A, see Table 1). Data were not filtered.

13-25.4N 61-03.1W 880M




Figure 15. Variance spectra of flow components and temperature (T^2/cph) at 880 m in St. Vincent Passage (array A, see Table 1). Data were not filtered.

13-19.0N 60-59.9W 380M




Figure 16. Variance spectra of flow components and temperature (T²/cph) at 380 m in St. Vincent Passage (array B, see Table 1). Data were not filtered.

13-19.0N 60-59.9W 820M




Figure 17. Variance spectra of flow components and temperature (T^2/cph) at 820 m in St. Vincent Passage (array B, see Table 1). Data were not filtered.

11-43.5N 61-55.5W 60M




Figure 18. Variance spectra of flow components and temperature (T^2/cph) at 60 m in Grenada Passage (see Table 1). Data were not filtered.

11-43.5N 61-55.5W 110M




Figure 19. Variance spectra of flow components and temperature (T^2/cph) at 110 m in Grenada Passage (see Table 1). Data were not filtered.

11-43.5N 61-55.5W 320M




Figure 20. Variance spectra of flow components and temperature (T^2/cph) at 320 m in Grenada Passage (see Table 1). Data were not filtered.

11-43.5N 61-55.5W 600M




Figure 21. Variance spectra of flow components and temperature (T^2/cph) at in Grenada Passage (see Table 1). Data were not filtered.




Figure 22. Low frequency variance spectra of E and N components at 105 m, 380 m, and 880 m in St. Vincent Passage (array A, see Table 1).

ST VINCENT PASSAGE P




Figure 23. Low frequency variance spectra of E and N components at 380 m and 810 m in St. Vincent Passage (array B, see Table 1).




Figure 24. Low frequency variance spectra of E and N components at 60 m, 320 m, and 600 m in Grenada Passage (see Table 1).




Figure 25. Coherence of E-W components of flow in St. Vincent Passage between depths of 105 m, 380 m, and 880 m (array A).




Figure 26. Coherence of N-S components of flow in St. Vincent passage between depths of 105 m, 380 m and 880 m (array A).


ST VINCENT PASSAGE B

UB20-0390

1.0

0.9

0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0.0



ST VINCENT PASSAGE B

UB20-0390

1.0

0.9

0.8
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0.0




Figure 27. Coherence of E-W and N-S components of flow in St. Vincent Passage between depths of 380 m and 820 m (array B).




Figure 28. Coherence of E-W component of flow in Grenada Passage between depths of 60 m 320 m and 600 m.




Figure 29. Coherence of N-S components of flow in Grenada Passage between depths of 60 m, 320 m and 600 m.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

showed peaks near 12-13 days in near-bottom records from St. Vincent passage, and peaks between 20 and 70 days in the other records.

Strong tidal signals were also found in the velocity records. Between 15% and 74% of the individual record variances were in diurnal, semidiurnal, or harmonic frequency bands. The semidiurnal frequency band contained from 11%-67% of the record variances. Tidal harmonics were also clearly evident, especially at the two near-bottom instruments deployed sequentially in St. Vincent Passage, where the first five harmonics of the semidiurnal frequency accounted for 27%-35% of the variance.

Each instrument recorded temperature, and most of the individual record variance was either subinertial (8%-86%) or semidiurnal (4%-82%). Together, these frequency bands accounted for 69% of the variance. Tidal harmonics accounted for 2%-25% of the temperature variance.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DATE
FILMED
-88