

AD-A100 565

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

ON THE BUCKLING OF LINEAR VISCOELASTIC RODS. (U)

MAY 81 M E GURTIN, V J MIZEL, D W REYNOLDS

DAAG29-80-C-0041

UNCLASSIFIED

MRC-TSR-2222

NL

1 of 1
AD-A100 565



END
DATE
FILMED
7-81
DTIC

— LEVEL II

(2)

AD A100565

MRC Technical Summary Report #2222

ON THE BUCKLING OF LINEAR VISCOELASTIC
RODS

Morton E. Gurtin, Victor J. Mizel and
David W. Reynolds

Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53706

DTIC
JUN 24 1981

May 1981

Received April 15, 1981

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P.O. Box 12211
Research Triangle Park
North Carolina 27709

81 C 23 073

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

ON THE BUCKLING OF LINEAR VISCOELASTIC RODS

Morton E. Gurtin¹, Victor J. Mizel¹, and David W. Reynolds²

Technical Summary Report #2222
May 1981

ABSTRACT

In this note we consider the linearized equation for the buckling of a viscoelastic rod from an undeformed virgin state. We show that this equation does not exhibit buckled solutions for axial end thrusts which - after application - are held constant. Though this result is apparently known, there appears to be no proof available in the literature.

We show further that if the load is allowed to vary with time, then, in contrast to elastica theory, there is an uncountably infinite number of buckled solutions.

AMS(MOS) Subject Classification: 73H05

Key Words: viscoelasticity, buckling

Work Unit No. 2 - Physical Mathematics

Accession For	
NTIS Grant	☒
DTIC	☐
AD	☐
AS	☐
CS	☐
DS	☐
ES	☐
FS	☐
GS	☐
HS	☐
IS	☐
JS	☐
KS	☐
LS	☐
MS	☐
NS	☐
OS	☐
PS	☐
QS	☐
RS	☐
TS	☐
US	☐
VS	☐
WS	☐
XS	☐
YS	☐
ZS	☐
Dist	Special
A	

¹Professor of Mathematics, Department of Mathematics, Carnegie-Mellon University, Pittsburgh, PA 15213.

²Research Assistant, Department of Mathematics, Carnegie-Mellon University, Pittsburgh, PA 15213.

SIGNIFICANCE AND EXPLANATION

The recent use of polymeric materials for structural purposes renders important the careful study of viscoelastic buckling, a phenomenon quite different from that associated with elastic materials, especially when the time scale of interest is large. In this paper we show that the linearized equation for the buckling of a viscoelastic rod from its virgin state does not exhibit buckled solutions for axial end thrusts which - after application - are held constant. We show further that if the load is allowed to vary with time, then, in contrast to elastica theory, there is an uncountably infinite number of buckled solutions.

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

ON THE BUCKLING OF LINEAR VISCOELASTIC RODS

Morton E. Gurtin¹, Victor J. Mizel¹, and David W. Reynolds²

1. Basic equations.

Consider a thin inextensible rod, pinned³ at the ends, and acted on by an axial end thrust $P(t)$ in such a way that its center-line bends in a plane. For each material point x and time t , let $\varphi(x,t)$ denote the angle between the horizontal axis and the tangent to the rod at x (Figure 1), and let $m(x,t)$ designate the bending moment at x . Here, for convenience, we label material points by their positions $x \in [0,L]$ in the undeformed, straight configuration the rod is assumed to be in prior to time zero.

We work within the quasi-static theory; thus balance of moments has the form⁴

$$m' + P \sin \varphi = 0, \quad (1)$$

where $m' = \partial m / \partial x$. We assume that the rod is linearly viscoelastic in the sense of the constitutive equation⁵

$$m(x,t) = \beta \varphi'(x,t) + \int_0^t \alpha(t-s) \varphi'(x,s) ds \quad (2)$$

giving the bending moment as a function of the past history of the curvature φ' . Here $\beta > 0$ is the instantaneous flexural

¹Professor of Mathematics, Department of Mathematics, Carnegie-Mellon University, Pittsburgh, PA 15213.

²Research Assistant, Department of Mathematics, Carnegie-Mellon University, Pittsburgh, PA 15213.

³In the interest of brevity, we consider only pinned ends. Our results go through, without change, for any of the standard boundary conditions. (In this connection, cf. [1].)

⁴Cf., e.g., [2], §262.

⁵Here we use the fact that $\varphi(x,t) = 0$ for $t < 0$.




Figure 1. The deformed rod

rigidity, while α , assumed continuous and <0 , measures internal dissipation; the unique solution $G(s)$, $s \geq 0$, of the initial-value problem

$$\dot{G}(s) = \alpha(s), \quad G(0) = \beta$$

is the (moment-curvature) relaxation function.

Equations (1) and (2) yield the integro-differential equation

$$\beta \varphi''(x,t) + \int_0^t \alpha(t-s) \varphi''(x,s) ds + P(t) \sin \varphi(x,t) = 0,$$

or equivalently, using the standard notation for convolutions,

$$\beta \varphi'' + \alpha * \varphi'' + P \sin \varphi = 0.$$

Since the ends of the rod are pinned, the relevant boundary conditions are

$$m(0,t) = m(L,t) = 0$$

for all t , or, by (2) and the standard uniqueness theorem for linear Volterra integral equations,

$$\varphi'(0,t) = \varphi'(L,t) = 0$$

for all t . Further, as the end of the rod corresponding to the material point $x = L$ can move only horizontally, we have the additional constraint

$$\int_0^L \sin \varphi(x,t) dx = 0.$$

In this note we confine our attention to the linear theory and hence replace $\sin \varphi$ by φ in the above equations. The resulting boundary-value problem then takes the form¹

$$\begin{aligned} \beta \varphi'' + \alpha * \varphi'' + P\varphi &= 0, \\ \varphi'(0,t) = \varphi'(L,t) &= 0, \end{aligned} \tag{3}$$

$$\int_0^L \varphi(x,t) dx = 0.$$

¹Cf. [3], [4] for a discussion of equations similar to (3)₁, but with $P = \text{constant}$. To our knowledge no previous paper has allowed $P(t)$ to depend on t .

2. Impossibility of buckled solutions when $P(t) = \text{constant}$.

In this section we show that the problem (3) has only the zero solution when

$$P(t) = P = \text{constant for } t \geq 0.$$

With this in mind, let $0 < P_1 < P_2 < \dots < P_n < \dots$ and $\psi_1(x), \psi_2(x), \dots, \psi_n(x), \dots$ denote the eigenvalues and eigenvectors of the corresponding elastic problem:

$$\begin{aligned} \beta \psi_n'' + P_n \psi_n &= 0, \\ \psi_n'(0) &= \psi_n'(L) = 0, \end{aligned} \tag{4}$$

$$\int_0^L \psi_n(x) dx = 0;$$

i.e.,

$$\psi_n(x) = \cos \frac{n\pi x}{L}, \quad P_n = \beta \frac{n^2 \pi^2}{L^2}, \quad n = 1, 2, \dots \tag{5}$$

Suppose that $\varphi(x, t)$ is a sufficiently smooth solution of (3) and define

$$\varphi_n(t) = \int_0^L \varphi(x, t) \psi_n(x) dx. \tag{6}$$

Then, since φ and ψ_n have vanishing spatial derivatives at the ends of the rod, two integrations by parts in conjunction with (4)₁ yield

$$\int_0^L \varphi'' \psi_n dx = \int_0^L \varphi \psi_n'' dx = - \frac{P_n}{\beta} \varphi_n.$$

Thus multiplying (3)₁ by ψ_n and integrating with respect to x over the interval $(0,L)$ results in the integral equation

$$(P-P_n)\varphi_n - \frac{P_n}{\beta} \alpha * \varphi_n = 0 \quad (7)$$

for $n = 1, 2, \dots$

If $P \neq P_n$, (7) is a Volterra equation of the second kind, which has the unique solution

$$\varphi_n(t) = 0 \quad (8)$$

for all t . On the other hand, if $P = P_n$,

$$\alpha * \varphi_n = 0,$$

and hence, by Titchmarsh's theorem¹, (8) holds in this case as well. By (6) and (8), φ is orthogonal to each function ψ_n of the form (5)₁. This is clearly possible only if $\varphi(x,t)$ is independent of x , and the desired conclusion, $\varphi \equiv 0$, follows from (3)₃.

¹Cf., e.g., [5], Theorem 152.

3. Existence of buckled solutions with $P(t) \neq \text{constant}$.

There are buckled solutions for certain nonconstant loads.
To see this consider solutions of the form

$$\varphi(x,t) = \gamma(t)\psi_n(x)$$

for $t \geq 0$. By $(4)_{2,3}$ this function satisfies the boundary conditions $(3)_2$ and the constraint condition $(3)_3$. Moreover, by $(4)_1$, φ will satisfy the differential equation $(3)_1$ provided γ and P satisfy the integral equation

$$[P(t) - P_n]\gamma(t) - \frac{P_n}{\beta} \int_0^t \alpha(t-s)\gamma(s)ds = 0. \quad (9)$$

This equation can have many solutions. A simple example is

$$\gamma(t) = \text{constant}, \quad P(t) = P_n Q(t),$$

with

$$Q(t) = \frac{G(t)}{G(0)}.$$

For this solution the angle $\varphi(x,t)$ jumps to its elastic shape at $t = 0$ and remains there for $t > 0$. The requisite axial thrust is initially the elastic buckling load P_n , but for $t > 0$, $P(t)$ decreases in proportion to the relaxation function $G(t)$.

Solutions continuous in time at $t = 0$ are also possible.
For example,

$$\gamma(t) = At \quad (A = \text{constant}), \quad P(t) = P_n \frac{1}{t} \int_0^t Q(s) ds,$$

forms a solution. (Note that, again, $P(0^+) = P_n$.)

More generally, (9) can be rewritten as

$$\frac{P(t)}{P_n} = \frac{\gamma(t) + \gamma * \dot{Q}(t)}{\gamma(t)}; \quad (10)$$

thus any function $\gamma > 0$ on $[0, \infty)$ yields a buckled solution with P given by (10). It is not difficult to show that $P(0^+)$ exists when $\gamma(0^+) = 0$, provided $\dot{\gamma}(0^+) > 0$. Thus any smooth function γ on $[0, \infty)$ with $\gamma > 0$ on $(0, \infty)$, $\gamma(0) = 0$, and $\dot{\gamma}(0^+) > 0$ also generates a buckled solution.¹

Thus, interestingly, in contrast to elastica theory there is an uncountably infinite number of buckled solutions (even modulo multiplicative constants).

Acknowledgment. This work was supported by the Army Research Office (MEG, DWR) and the National Science Foundation (VJM).

¹Conditions on $P(t)$ which insure the existence of nontrivial $\gamma(t)$ are given in [1]. This paper also investigates the asymptotic behavior of buckled solutions.

References

- [1] Reynolds, D. W., On the buckling of viscoelastic rods.
In preparation.
- [2] Love, A.E.H., A Treatise on the Mathematical Theory of Elasticity, Dover: New York, 1944, Fourth Edition.
- [3] Distéfano, J. N., Sulla stabilità in regime viscoelastico a comportamento lineare I. Atti Acad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur (8) 27, 356-361.
- [4] Distéfano, J. N., Creep buckling of slender columns. J. Strut. Div. Proc. of A.S.C.E. 91, 127-150.
- [5] Titchmarsh, E. C., Introduction to the Theory of Fourier Integrals, Oxford: Oxford, 1948, Second Edition.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2222	2. GOVT ACCESSION NO. AD-A100	3. RECIPIENT'S CATALOG NUMBER 565
4. TITLE (and Subtitle) ON THE BUCKLING OF LINEAR VISCOELASTIC RODS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Morton E. Gurtin, Victor J. Mizel and David W. Reynolds		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 2 - Physical Mathematics
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE May 1981
		13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) viscoelasticity, buckling		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this note we consider the linearized equation for the buckling of a viscoelastic rod from an undeformed virgin state. We show that this equation does <u>not</u> exhibit buckled solutions for axial end thrusts which - after application - are held constant. Though this result is apparently known, there appears to be no proof available in the literature. We show further that if the load is allowed to vary with time, then, in contrast to elastica theory, there is an uncountably infinite number of buckled solutions.		

**DAT
FILM**