

AD-A101 785

NAVAL HEALTH RESEARCH CENTER SAN DIEGO CA

F/G 6/15

EFFECTS OF TRIAZOLAM (0.5 MG) ON SLEEP, PERFORMANCE, MEMORY, AN--ETC(U)

MAR 81 C L SPINWEBER, L C JOHNSON

UNCLASSIFIED

NAVHLTHRSCHC-81-1

NL

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

END

DATE

FILED

4 81

DTIC

AD A101785

LEVEL II

EFFECTS OF TRIAZOLAM (0.5 mg)
ON SLEEP, PERFORMANCE, MEMORY, AND
AROUSAL THRESHOLD

C. L. SPINWEBER & L. C. JOHNSON

REPORT NO. 81-1

See 1473

DTIC
ELECTE
JUL 25 1981
S D



This document has been approved
for public release and sale; its
distribution is unlimited.

NAVAL HEALTH RESEARCH CENTER

P. O. BOX 1000
SAN DIEGO, CALIFORNIA 92161

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

FILE COPY

81 7 22 038

81 7 22 008

EFFECTS OF TRIAZOLAM (0.5 mg) ON SLEEP, PERFORMANCE, MEMORY,
AND AROUSAL THRESHOLD

Cheryl L. Spinweber and Laverne C. Johnson
Clinical Psychophysiology Department

Naval Health Research Center
P.O. Box 85122
San Diego, CA 92138

Report No. 81-1, supported by Naval Medical Research and Development Command, Bethesda, Maryland, Department of the Navy, under research Work Unit MRO41.01.003-0157. This research was partially supported by The Upjohn Company, Kalamazoo, Michigan, in a grant to the San Diego State University Foundation under Account No. 528008. The views presented in this paper are those of the authors. No endorsement by the Department of the Navy has been given or should be inferred.

The authors thank Michael H. Bonnet, Ph.D., for his contributions to the design of this protocol.

SUMMARY

Currently marketed sleeping pills are long-acting drugs which impair morning performance following bedtime administration, thus rendering them unsuitable for use in operational settings. Triazolam, a newer, short-acting benzodiazepine, was evaluated to determine its usefulness as a hypnotic and to measure its effects on performance, memory, and arousal threshold. The rapid metabolism and clearance of triazolam suggested that it might promote improved sleep without producing a drug-induced performance impairment the following morning.

Twenty male poor sleepers, mean age 21 ± 2.37 years, were studied. To qualify as a poor sleeper, subjects had to rate their sleep quality as "poor" or "very poor," report a usual sleep latency greater than 45 minutes, and indicate that the problem had persisted for at least 6 months. To meet EEG sleep criteria on the screening night, poor sleepers had to exhibit sleep latencies of 30 minutes or longer and have at least 5% of their total sleep time in Stages 3 + 4, slow wave sleep.

Following the screening night, subjects received placebos in a single-blind paradigm for 3 consecutive baseline nights. Following the placebo-baseline nights, 10 subjects received 0.5 mg triazolam for 6 nights while the other 10 continued to receive placebo in a double-blind paradigm. After the 6 treatment nights, all subjects received placebo on 2 withdrawal nights. All-night sleep EEGs were recorded according to the usual laboratory procedures.

Performance batteries were administered 20-40 minutes after the morning awakening. Tests included the Wilkinson 4-Choice Reaction Time Test, the Digit Symbol Substitution Test, the Williams Word Memory Test, and the Card Sorting Task. On night 10, the performance night, subjects were aroused from Stage 2 sleep, at 1.5, 3, and 5 hours post-drug ingestion, to perform the same tasks, in order to assess the acute effects of triazolam on performance. In the morning following night 10, subjects were presented with the Memory Checklist. This list contained the 90 words which were presented in the Williams Word Memory Task during nighttime test sessions plus 90 filler words. Subjects were instructed to pick out the words they remembered from the nighttime sessions.

Arousal threshold was obtained on 3 study nights. Tones were played over a loudspeaker and incremented in 5 dB steps until the sleeper was awakened by the tone.

Triazolam was found to be an effective hypnotic which significantly reduced sleep latency and increased total sleep time. Morning performance was unimpaired by bedtime administration of triazolam. However, at 1.5, 3, and 5 hours post-administration, ability to perform was significantly lowered. In addition, triazolam was found to produce anterograde amnesic effects--impairment of memory for stimuli presented during the acute phase of drug action. Triazolam also significantly elevated arousal threshold during the night.

Triazolam is suitable for use as a sleeping pill in settings in which personnel will be scheduled for 7.5 hours of sleep. However, since triazolam does impair ability to perform visuo-motor and cognitive tasks for up to 5 hours post-administration, impairs memory for events occurring during the night, and elevates arousal threshold, its usefulness in operational settings in which there might be a need to arouse personnel to perform duties is limited.

INTRODUCTION

In sleep laboratory studies, triazolam (Halcion), a triazolobenzodiazepine, has been demonstrated to be an effective hypnotic in doses ranging from 0.25–1.0 mg.¹⁻⁹ In addition, results of outpatient studies and other investigations which used self-report measures to evaluate hypnotic efficacy have shown that sleep is subjectively improved after triazolam administration.¹⁰⁻¹⁸

Pharmacokinetic measures indicate that both triazolam and its active metabolite, 7- α -hydroxy triazolam, have short half-lives, reported to be within the range of 2.1–10 hours.¹⁹⁻²¹ The relatively rapid metabolism and clearance of triazolam suggest that, at least at lower doses within the effective range, this benzodiazepine may promote improved sleep without producing a drug-induced performance impairment the following morning. Several studies have assessed morning performance following bedtime triazolam administration.^{2,5,7,10,17,22} The results of these studies differ, depending upon dose size, time post-ingestion when tests were administered, and the type of tasks used. Only one of these performance studies used insomniacs as subjects.⁵

Acute effects of triazolam (0.25 mg and 0.5 mg) on performance have been demonstrated at 3.5 hours post-administration in testing following arousal from sleep.² Nicholson and Stone⁷ have reported an impairment of visuo-motor performance up to 5 hours post-administration of a 0.25 mg dose during daytime testing in subjects who remained awake. We believe that ours is the first study of benzodiazepine hypnotics to employ multiple systematic arousals from sleep, designed to delineate temporal parameters of acute effects and provide behavioral correlates of benzodiazepine pharmacodynamics in the sleeping subject.

MATERIALS AND METHODS

Subjects

Twenty male poor sleepers, mean age 21 ± 2.37 years, were studied. Poor sleep was defined by both EEG and subjective criteria. Subjective criteria included responses to a questionnaire designed to elicit the individual's reports of his sleep quality. To qualify as a poor sleeper, subjects had to rate their sleep quality as "poor" or "very poor," indicate a usual sleep latency greater than 45 minutes (sleep onset insomnia), and report having this sleep onset problem for at least 6 months. The subject's subjective report was discussed further in a personal interview. To meet EEG sleep criteria on the screening night, poor sleepers had to exhibit sleep latencies (time from lights out to the onset of Stage 2 sleep) of 30 minutes or longer and have at least 5% of their total sleep time in slow wave sleep (SWS) (Stages 3 + 4). During screening nights, average sleep latency for the 20 subjects was 53.6 ± 34 minutes.

Subjects were screened for possible psychiatric conditions, sensitivity to benzodiazepines, alcohol or drug abuse, and recent illnesses. All subjects were in good health and denied current or recent use of any type of sleep medication or other drugs. There were no sleep complaints other than those associated with falling asleep.

All subjects were informed about the general nature of the experiment and willingly signed Informed Consent and Privacy Act statements. All subjects were asked to refrain from napping and taking drugs or alcohol during the course of the study. Breath analyzer and urine tests, used aperiodically, indicated no detectable use of alcohol or other drugs during the study.

Based on screening night findings, 20 possible poor sleepers were rejected because of sleep latencies less than 30 minutes. Two subjects were dropped from the study during placebo-baseline due to concern over poor academic performance. No subjects were dropped because of side effects.

Procedure

A parallel, three-phase design was employed. Subjects who qualified as poor sleepers on the screening night went on to complete 11 additional nights of the 12-night protocol (see Table 1).

Following the screening night, subjects received placebos in a single-blind paradigm for 3 consecutive baseline nights. Following the placebo-baseline nights, 10 subjects received 0.5 mg triazolam for 6 nights while the other 10 continued to receive placebo in a double-blind paradigm. After the 6 treatment nights, all subjects received placebo on 2 withdrawal nights. The placebo or drug tablet was given at 2145 hours each night. Lights out was at 2200 and subjects were awakened at 0530.

Each subject slept in an electrically shielded, air-conditioned room with soundproofing. All electrophysiological variables were recorded on an 8-channel Beckman dynograph. The electro-oculogram (EOG) was recorded from biopotential electrodes placed on the outer canthus of each eye. The EEGs were obtained by use of silver chlorided disc electrodes from C₃ and O₁ electrode placements referenced to linked mastoids (A₁ + A₂). Both EOG and EEG time constants were 0.3 seconds. Sleep stages were determined according to standard criteria.²³

TABLE 1
Triazolam Study Protocol

	Condition											
	Placebo-Baseline				Treatment (Placebo or Triazolam)						Placebo-Withdrawal	
Night #	1	2	3	4	5	6	7	8	9	10	11	12
Procedure	(S)	(E)*	A	(C)*	(E)*	A	(E)*	A	(C)*	(P)	(E)*	(E)*

Procedure Code

S = Screening Night

E = All-night EEG

A = Arousal Night

C = AEP Night

P = Performance Night

○ Morning Testing

* Paired-Associates Testing

Subjects were familiarized with all questionnaires and trained on all tasks in a practice session conducted prior to night 1 of the study.

Bedtime and Morning Questionnaires. The subjects completed a Bedtime Questionnaire each evening which required the subject to report side effects, unusual events occurring that day, naps, alcohol consumption, and his readiness for bed. The Stanford Sleepiness Scale (SSS)²⁴ was also included in this questionnaire. Upon awakening at 0530, the subjects completed a Morning Questionnaire which included the SSS, and also required the subject to estimate sleep latency, total sleep time, and number of awakenings, to list any physical complaints, and to rate the effectiveness of the pill and evaluate sleep quality.

Sleep Measures. Sleep latency (time from lights out to the onset of Stage 2) was scored for all study nights. Mean sleep latencies were derived for each subject for each condition: placebo-baseline (nights 2-4), treatment (nights 5-10), and placebo-withdrawal (nights 11-12).

Sleep stage data were obtained for comparison on nights of uninterrupted sleep (*i.e.*, on study nights on which subjects were not awakened to perform tasks or respond to tones or on nights when auditory evoked potentials (AEPs) were obtained). For the three conditions, sleep measures were obtained as follows: placebo-baseline (night 2), treatment (mean of nights 5 and 7), and placebo-withdrawal (mean of nights 11 and 12). Sleep measures were: Total Sleep in minutes (the sum of minutes in Stages 2, 3, 4, and REM); Stage 1 percent (minutes of Stage 1 divided by the total bed time x 100); Stage 2 percent, Stage 3 percent, Stage 4 percent, and Stage REM percent (minutes in each stage divided by Total Sleep x 100); Sleep Efficiency (Total Sleep divided by Total Bedtime x 100); Wake Time (minutes awake while in bed); Wake percent (minutes awake divided by Total Bedtime x 100).

Morning Performance and Mood Testing. Performance and mood test batteries were administered approximately 20-40 minutes after the morning awakening following nights 1, 2, 4, 5, 7, 9, 10, 11, and 12. Data from testing following the screening night (night 1) were not included in the data analysis. Morning batteries included two subjective mood scales, the NHRC Mood Scale and the Profile of Mood States (POMS), and several performance tests, including the Wilkinson 4-Choice Reaction Time Test (performed for 11 minutes) and the Digit Symbol Substitution Test. These tests have been described in detail in a previous publication.²⁵ The test battery also included the Williams Word Memory Test²⁶: this task is a test of short-term memory. Subjects heard a tape-recorded list of 15 words. The voice on the tape said each word, spelled the word, and then repeated each word again. During list presentation, the subject wrote down each word. At the end of the 15-word presentation, the subject was allowed 3 minutes to write down as many of the words as he could recall. Two lists of words were presented during each performance battery. The subjects also performed the Card Sorting Task: this timed task required subjects to sort a deck of 36 numbered playing cards (cards numbered 2-10 in each of 4 suites) first into 4 piles according to suite and then into 9 piles according to number. Time (in seconds) to complete the sorting task was recorded.

The effects of triazolam on retention of material learned prior to drug administration were evaluated through use of a paired associate (P-A) learning task, a modified version of Ekstrand's memory test.²⁷ On study nights 2, 4, 5, 7, 9, 11, and 12, approximately 1 hour prior to drug

administration, the subject learned 10 word pairs from a tape-recorded list. The first word of each pair was a 3-letter noun selected from the 10-40 frequency range²⁸ and the second was an unrelated adjective selected from the 10-20 frequency range. The tape first presented the 10 word-pairs, then presented the first word of each pair in a random order to permit the subject to give the response. If the subject was unable to give all response words correctly, a list of the 10 pairs was presented again. Presentation of the 10 pairs followed by the testing procedure continued until the subject had learned all 10 pairs. Morning recall of the 10 pairs was tested in two ways: in P-A Recall, the subject was required to write down the response word when presented with a list of the 10 nouns. In P-A Matching, the subject was required to match nouns and adjectives of the pairs. For both tasks, number correct was recorded.

Performance and Mood after Arousal from Sleep. On night 10, the performance night, subjects were aroused from Stage 2 sleep during three preestablished time windows--90-100 minutes, 180-200 minutes, and 270-300 minutes after lights out--to complete performance and mood tests. Performance and mood testing during these time windows generally occurred at times 1.5, 3, and 5 hours post-drug ingestion. These nighttime batteries differed from the morning batteries in two aspects only: the SSS was administered in place of the POMS and the 4-Choice Reaction Time Test was performed for 6 rather than 11 minutes.

Following completion of each test battery, the subjects were instructed to go back to sleep. Latency of the return to sleep was also recorded.

In the morning following night 10, subjects were presented with the Memory Checklist. This list contained the 90 words which were presented in the Williams Word Memory Task during nighttime test sessions plus 90 filler words. Subjects were instructed to pick out the words they remembered from the nighttime sessions.

Arousal Threshold Procedure. The threshold for arousal from sleep was obtained on 3 recording nights: night 3 (placebo-baseline), night 6 (second treatment night), and night 8 (fourth treatment night). Tones were delivered over a loudspeaker positioned approximately 46 cm above the sleeper's head. The subject's threshold for tones while awake was obtained prior to lights out. During arousals from sleep, tones were begun at 20 dB above the awake threshold and were incremented in 5 dB steps until the subject made the behavioral (three button pushes) and verbal ("I'm awake") responses. Tones were 2 seconds long and occurred at 16-second intervals. Arousals were scheduled to reveal the time course of action of triazolam and were performed six times: #1: during the first Stage 2 sleep, 5 minutes after the sleep onset; #2: during the first SWS (Stage 3 or Stage 4), 20 minutes after the return to sleep following the first arousal; #3: in Stage 2, 150-210 minutes after lights out (0030-0130); #4: in Stage 2, 270-330 minutes after lights out (0200-0330); #5: in Stage 2, 370-430 minutes after lights out (0410-0510); #6: the morning arousal, at 0530.

Additional criteria which had to be met to initiate arousal procedures were: (1) there could be no major (8 seconds or longer) body movement for 10 minutes prior to the arousal. (2) Stage 2 or SWS had to be well-defined for 5 minutes prior to the arousal. After the subject made the appropriate response, he was told to go back to sleep. The dB level for the highest tone presented and the latency (in minutes) from the time of the awakening to the return to sleep were recorded.

EEG Parameters. On-line EEG detection of spindles and delta was performed for nights 2, 4, 5, 7, 9, 11, and 12. AEPs were recorded on nights 4 and 9. These EEG findings are reported in detail elsewhere.²⁹

Statistical Analysis. Mean placebo-baseline, mean treatment, and mean placebo-withdrawal scores were derived for each subject for the various measures. Mean data were statistically evaluated using between-groups *t*-tests on difference scores, derived by subtracting each subject's mean treatment or mean placebo-withdrawal score from his mean placebo-baseline score. For night 10 data, Analysis of Variance for Repeated Measures, with "Sessions" and "Groups" as factors, was used. Night-by-night data were also plotted, inspected, and appropriately tested if there was evidence of night-by-night effects. Additional analyses for specific measures are described in the results section. All tests were one-tailed unless otherwise stated.

RESULTS

Effects on Sleep. A night-by-night plot of mean sleep latencies is presented in Fig. 1. Mean sleep data by conditions are presented in Table 2. Triazolam significantly reduced sleep latency during treatment ($t_{18} = 2.1051$, $P < 0.025$). The medication also increased total sleep time ($t_{18} = 2.3896$, $P < 0.025$), sleep efficiency ($t_{18} = 2.4279$, $P < 0.025$), and decreased wake time ($t_{18} = 2.32469$, $P < 0.025$) and wake percent ($t_{18} = 1.79634$, $P < 0.05$) during treatment.

Stage 2 percent was increased ($t_{18} = 4.3160$, $P < 0.0005$) and Stage 4 percent was decreased ($t_{18} = 2.1224$, $P < 0.025$). Brain electrical activity was correspondingly altered: spindle rate per minute of NREM was elevated ($t_{18} = 4.04$, $P < 0.0005$) and delta rate per minute NREM sleep was reduced ($t_{18} = 3.88$, $P < 0.005$). (See Johnson and Spinweber²⁹ for a further discussion of spindle and delta count findings.) REM percent was reduced in the triazolam group during treatment ($t_{19} = 3.0391$, $P < 0.005$).




Fig. 1. Mean sleep latencies for nights 2-12.

TABLE 2
Mean Sleep Data by Conditions^a

	Placebo Group		
	Placebo-Baseline $\bar{X}$ ($\pm$ SD)	Treatment $\bar{X}$ ($\pm$ SD)	Placebo-Withdrawal $\bar{X}$ ($\pm$ SD)
Sleep latency (minutes) ^c	26.65 (7.56)	33.82 (17.95)	26.55 (9.09)
Spindles/minute NREM ^e	3.36 (1.31)	3.24 (1.57)	3.55 (1.41)
Delta/minute NREM ^d	40.92 (20.83)	39.65 (19.89)	39.66 (17.72)
Total sleep (minutes) ^c	389.65 (21.00)	392.07 (18.64)	398.87 (19.78)
Stage 1 percent	3.82 (1.54)	4.21 (1.56)	3.86 (2.02)
Stage 2 percent ^e	50.72 (9.03)	52.57 (8.98)	53.27 (6.40)
Stage 3 percent	8.03 (2.23)	7.59 (1.54)	7.42 (1.60)
Stage 4 percent ^e	11.00 (6.59)	9.36 (6.00)	9.72 (5.95)
REM percent ^d	30.25 (6.11)	30.48 (6.38)	29.58 (4.06)
Sleep efficiency ^c	87.04 (4.74)	87.60 (4.03)	89.20 (4.30)
Wake time (minutes) ^c	34.30 (19.23)	27.40 (16.27)	21.72 (9.70)
Wake percent ^b	7.66 (4.30)	6.13 (3.65)	4.86 (2.19)

	Triazolam Group		
	Placebo-Baseline $\bar{X}$ ($\pm$ SD)	Treatment $\bar{X}$ ($\pm$ SD)	Placebo-Withdrawal $\bar{X}$ ($\pm$ SD)
Sleep latency (minutes) ^c	35.54 (23.96)	26.67 (7.34)	44.88 (35.30)
Spindles/minute NREM ^e	3.97 (2.23)	5.69 (2.69)	3.86 (1.95)
Delta/minute NREM ^d	47.52 (25.90)	32.08 (23.89)	37.66 (27.39)
Total sleep (minutes) ^c	373.95 (32.53)	406.45 (6.81)	381.12 (29.95)
Stage 1 percent	3.57 (1.61)	2.94 (1.57)	3.78 (1.30)
Stage 2 percent ^e	48.70 (7.45)	60.67 (5.26)	50.12 (7.66)
Stage 3 percent	10.95 (3.91)	9.03 (4.33)	9.30 (3.28)
Stage 4 percent ^e	9.45 (7.23)	4.89 (7.00)	8.09 (6.75)
REM percent ^d	30.89 (4.57)	25.40 (3.09)	32.48 (6.03)
Sleep efficiency ^c	83.43 (7.10)	90.77 (1.50)	85.04 (6.59)
Wake time (minutes) ^c	50.30 (35.87)	21.95 (8.35)	41.15 (34.53)
Wake percent ^b	11.24 (8.05)	4.90 (1.86)	9.19 (7.72)

^a For sleep latency, means for conditions were computed as follows: Placebo-Baseline = Nights 2-4; Treatment = Nights 5-10; Placebo-Withdrawal = Nights 11-12; For other measures: Placebo-Baseline = Night 2; Treatment = Nights 5 and 7; Placebo-Withdrawal = Nights 11-12.

^b $P < 0.05$, one-tailed (comparison of Placebo-Baseline and Treatment difference scores).

^c $P < 0.025$, one-tailed (comparison of Placebo-Baseline and Treatment difference scores).

^d $P < 0.005$, one-tailed (comparison of Placebo-Baseline and Treatment difference scores).

^e $P < 0.0005$, one-tailed (comparison of Placebo-Baseline and Treatment difference scores).

In order to determine if triazolam produced "rebound insomnia" on the first withdrawal night, between-groups t -tests were performed on difference scores derived by subtracting each subject's night 11 data from his night 2 (placebo-baseline) data. No significant differences between placebo-baseline and the first withdrawal night were found on measures of sleep latency, total time awake, and total sleep time. In addition, a within-group paired t -test was performed on mean sleep latencies for the placebo-baseline and placebo-withdrawal conditions for the triazolam subjects alone. It was found that sleep latency returned to placebo-baseline values but did not exceed these values during withdrawal.

All sleep parameters which were significantly altered by triazolam during treatment returned to placebo-baseline levels during withdrawal. Consistent with the EEG data, triazolam subjects reported a significantly reduced subjective estimate of sleep latency during treatment ($t_9 = 2.4528$, $P < 0.025$).

Effects on Morning Performance and Mood. At the time of morning test sessions, approximately 8.25 hours post-administration, there were no significant differences between the placebo and triazolam groups on performance and mood measures.

Time Course of Acute Effects

Performance. Table 3 summarizes performance data for test sessions conducted during the three arousals from sleep on night 10. In addition, morning performance data obtained in testing following night 10 are presented. These morning test results are representative of all morning test sessions. For one subject in the placebo group, the second performance battery of the night was not performed, due to procedural error, thus altering the degrees of freedom reported below. Significant main effects of Groups were found for card sorting time ($F_{1,17} = 11.14$, $P < 0.004$), total correct on digit substitution ($F_{1,17} = 10.19$, $P < 0.005$), and number correct on Williams Word Memory Test (for the first list presented in each performance battery $F_{1,17} = 13.02$, $P < 0.002$ and, for the second list, $F_{1,17} = 5.81$, $P < 0.028$). Post-hoc between-groups t -tests for each test session indicated that the triazolam group performed significantly worse on each performance measure during each of the three test sessions. On the 4-Choice Reaction Time Test, data from the third nighttime test session and the morning session following night 10 on one triazolam subject were lost due to a faulty cassette tape. Because of this additional data loss, the omnibus F test was not performed for the 4-choice test but between-groups t -tests indicated that reaction time was significantly slower for triazolam subjects at each test session.

Performance data from all four tasks show a gradual recovery which begins at the time of the third test session, approximately 5 hours post-drug and, in comparison with morning performance on placebo, full recovery is seen by the time of morning testing. For example, on the Wilkinson 4-Choice Reaction Time Task, curves for the placebo group and triazolam group converge by the time of morning testing (Fig. 2). There were no significant differences between the two groups on subjective (SSS, and NHRC Mood Scale) measures.

Latency of Return to Sleep. The latency of return to sleep following nighttime testing was significantly shorter for the triazolam group for each of the three sessions (Fig. 3). The average latency of return to sleep following nighttime performance sessions was 5.10 ± 2.30 minutes for the triazolam subjects and 11.89 ± 6.87 minutes for placebo subjects ($t_{18} = 2.9641$, $P < 0.005$).




Fig. 2. Mean reaction time on the Wilkinson 4-Choice Reaction Time Task during three arousals from sleep (at 1.5, 3, and 5 hours post-drug) and for the morning test session (AM) at 8.25 hours post-ingestion.

SLEEP LATENCIES AFTER PERFORMANCE TESTING




Fig. 3. Mean latencies of return to sleep following nighttime test sessions on night 10. Sleep latency is significantly shorter for the triazolam group at 1.5 hours ($t_{18} = 1.7629$, $P < 0.05$), at 3 hours ($t_{17} = 3.7437$, $P < 0.005$), and at 5 hours ($t_{18} = 2.4324$, $P < 0.025$).

TABLE 3
Performance During Arousals from Sleep and Morning Performance Following Night 10

Task	Hours Post-Drug	Triazolam $\bar{X}$ ($\pm$ SD)	Placebo $\bar{X}$ ($\pm$ SD)	<i>t</i>	<i>df</i>	<i>P</i> <
Four-Choice Reaction Time (ms)	1.5	1020. (279.)	650. (127.)	3.795 ^a	17	0.005
	3	1139. (607.)	676. (101.)	2.382 ^a	17	0.025
	5	908. (129.)	717. (172.)	2.599 ^a	16	0.01
	8.25	598. (86.)	558. (50.)	0.0188 ^b	16	n.s.
Digit Symbol Substitution (score)	1.5	32.65 (14.31)	48.89 (9.73)	2.9627 ^a	18	0.005
	3	27.75 (13.44)	48.22 (7.38)	4.0460 ^a	17	0.0005
	5	36.45 (15.93)	46.67 (9.22)	1.8027 ^a	18	0.05
	8.25	52.70 (11.38)	59.70 (11.65)	0.1542 ^b	18	n.s.
Williams Word Memory Test (No. correct)	1.5	8.70 (4.27)	13.60 (2.32)	3.1889 ^a	18	0.005
	3	8.80 (3.68)	14.00 (3.46)	3.1633 ^a	17	0.005
	5	11.00 (4.11)	14.40 (2.59)	2.2132 ^a	18	0.025
	8.25	16.30 (3.56)	15.70 (2.45)	0.8961 ^b	18	n.s.
Card Sorting Time (seconds)	1.5	171.35 (37.16)	119.28 (19.06)	3.8355 ^a	18	0.005
	3	176.70 (63.92)	123.78 (22.95)	2.3458 ^a	17	0.025
	5	155.15 (31.74)	120.00 (21.91)	2.7420 ^a	18	0.01
	8.25	110.17 (13.22)	97.75 (9.32)	0.6607 ^b	17	n.s.

^a Results of between-groups *t*-tests.

^b Results of between-groups *t*-tests on difference scores (mean Placebo-Baseline minus Morning score).

Effects on Memory

In the morning following night 10, triazolam subjects had a significantly lower score (number correct) on the Memory Checklist ($\bar{X}_{\text{drug}} = 17.30 \pm 8.38$ vs. $\bar{X}_{\text{placebo}} = 40.40 \pm 18.67$, $t_{18} = 3.5689$, $P < 0.005$). Since the target words on this checklist were words presented during the previous night on the Williams Word Memory Test, further analyses were performed to determine when memory loss had occurred. In a procedure similar to that described by Roth et al.,²² a total night recall score for each subject was derived by summing the total number of words correctly recalled during the night on the presentations of the Williams Word Memory Test lists and then obtaining a mean percentage loss for the night. For the triazolam group, immediate loss during the night was $69.89 \pm 10.07\%$ and for the placebo group, $53.67 \pm 6.75\%$ ($t_{18} = 4.2367$, $P < 0.0005$). An inspection of the words correctly identified on the Memory Checklist the following morning indicated that, in addition to recognizing some words which they had correctly recalled during the night, subjects in

both groups correctly recognized target words in the morning which they had not been able to recall during nighttime testing. In order to quantify additional memory loss occurring across the night, we considered only the pool of words that the subject had correctly recalled during the night and calculated the percentage of loss from that pool--i.e., the words subjects recalled correctly at night but were unable to recognize correctly on the Memory Checklist in the morning. For the triazolam group, the loss from nighttime testing to morning testing was $66.92 \pm 12.11\%$ and for the placebo group, $40.02 \pm 21.92\%$ ($t_{18} = 3.3971$, $P < 0.0005$).

Correlational analysis revealed a significant positive relationship between latency of return to sleep (during night 10) and Memory Checklist score in the morning. Over all subjects, the correlation coefficient, Pearson r , was 0.53 ($P < 0.025$). This relationship was significant within the triazolam group ($r = 0.72$, $P < 0.025$) but not within the placebo group.

Time Course of Effects on Arousal Threshold

Arousal threshold was significantly higher during treatment for triazolam subjects at the time of the first ($t_{18} = 2.4370$, $P < 0.025$), second ($t_{16} = 5.6518$, $P < 0.0005$), and third ($t_{17} = 2.9334$, $P < 0.005$) arousals (Fig. 4). (Not all subjects met the criteria for all arousals, thus altering the degrees of freedom reported above.) Within-groups analyses revealed that triazolam significantly raised arousal threshold for the SWS arousal ($t_9 = 3.4001$, $P < 0.005$); it was also found that placebo group subjects became more sensitive to the tone with repeated experience and had significantly reduced arousal threshold levels during the treatment condition for the first, second, and third arousals. Consistent with the arousal threshold data, latency of return to sleep following arousal by tones was significantly reduced for the first through third arousals in triazolam subjects during treatment.




Fig. 4. Mean arousal thresholds for the placebo-baseline night (n3) and for the mean of 2 treatment nights (n6&8).

DISCUSSION

Our results replicate previous reports of the hypnotic efficacy of triazolam (0.5 mg) and further confirm the well-described effects of benzodiazepines on sleep and brain activity during sleep. Previous reports of the effects of triazolam on SWS have been inconsistent,^{1,3-7,9} but, in our subjects who were screened to insure that they had adequate amounts of SWS to show an alteration, there was a highly significant reduction in Stage 4 percent. Even though the metabolic pathway of triazole benzodiazepines is different from that of flurazepam, its effects on sleep stages and spindle and delta activity are similar. The major difference between triazolam and longer-acting hypnotics in terms of sleep-related effects is that sleep measures return to baseline levels during short-term withdrawal, a finding which emphasizes, in behavioral terms, the fact that triazolam and its major metabolite, 7- α -hydroxy triazolam, are rapidly metabolized and do not appear to produce any substantial cumulative effects.

The advantages of use of a parallel, three-phase design are also emphasized by our findings. For example, in the night-by-night sleep latency data, we note the previously described but substantial effects of placebo administration in reduction of sleep latency in both groups during the placebo condition. In longer administration, placebo loses its effectiveness and sleep latencies in the placebo group become highly variable over nights. Triazolam, however, maintains the reduction in sleep latency during treatment. A finding which deserves emphasis is that our data do not show "rebound insomnia" in subjective and objective measures of sleep latency during withdrawal as has been found in previous research.³⁰

The major goals of this study were (1) to determine if bedtime administration of triazolam would cause a performance decrement in the morning, and (2) to determine the behavioral time course during the night of acute effects on performance. Our results demonstrate that morning mood and performance are not deleteriously affected by administration of triazolam (0.5 mg) at bedtime. Thus, triazolam is suitable for use when the individual will not be required to perform these tasks for 8.25 hours post-administration. However, assessment of the acute effects of triazolam administration--at approximately 1.5, 3, and 5 hours post-ingestion--reveals that cognitive, memory, and visuo-motor performance are impaired. To our knowledge, this study was the first to measure the time course of drug effect during sleep on the EEG and on behavioral response through systematic arousals of the sleeping subject. Most sleep studies have looked at early morning and subsequent daytime performance. The time course of the behavioral effects following drug ingestion has also been studied in subjects who remain awake during the entire study period. It is of interest that, for triazolam, at least, the time course of drug effects on performance tasks is the same regardless of whether the subject remains awake, or is allowed to sleep and is awakened to perform. Nicholson and Stone,⁷ using a 0.25 mg triazolam dose level, and a 0.4 mg brotizolam dose level,³¹ reported a similar temporal change on a visual motor (tracking) task. Though the Nicholson et al. findings and ours cannot be directly compared, the similarity of results on visual motor tasks suggests that whether the subject is awake or asleep is not a major factor in the pattern of the performance change post-drug and, perhaps, in the rate of drug metabolism.

Our findings on the effects of triazolam on memory are consistent with previous reports of anterograde amnesia produced by presurgical administration of benzodiazepines³²⁻³⁵ and also with

previous reports of anterograde amnesic effects of benzodiazepines administered as hypnotics.^{22,36} The memory impairment associated with hypnotic use could be the result of several underlying mechanisms: a lowered arousal level during stimulus presentation, disruption of memory consolidation, or interference with retrieval mechanisms. Roth et al.²² have recently presented data supporting the hypothesis that the mechanism of memory loss is an impairment of consolidation caused by a more rapid return to sleep following stimulus presentation on the drug nights. In our procedures, the Williams Word Memory Task was the last task presented during nighttime test sessions. Following completion of this test, the lights were turned out and subjects were instructed to return to sleep; thus, differences in wake time after completion of the task were almost entirely due to sleep latency. The significant correlation between sleep latencies during night 10 and number correct on the morning Memory Checklist indicates that rapid return to sleep may be a factor in the memory loss across the night. The time period during which memory systems are disrupted by rapid return to sleep is quite short, since, in placebo subjects who remained awake on the average about 6 minutes longer, there was no significant correlation between latency of return to sleep and morning recall.

However, the interpretation of the findings on memory are further complicated by the fact that our triazolam subjects showed a significantly larger percentage loss in immediate recall during nighttime testing on the Williams Word Memory Tests. Because of the nature of this task, in which subjects must write down each word during stimulus presentation, subjects had to attend to the stimulus material and the material received at least preliminary processing. This larger short-term loss raises the possibility that triazolam disrupts entrance of material into short-term store or reduces the number of items which can be held in short-term store for immediate retrieval. Since a wide variety of performance measures, including visuo-motor and other cognitive tasks, showed impairment during the night, a parsimonious explanation might be that arousal level is reduced due to the presence of the hypnotic in the brain. Our finding that morning recall of word pairs on the P-A task was not impaired during treatment demonstrates that triazolam administration does not produce retrograde amnesia.

Consistent with the findings describing the time course of acute effects of triazolam on performance during the night was our finding of similar effects on arousal threshold and latency of return to sleep after arousals. Our parallel design permitted us to identify the fact that with repeated exposure to the arousal tones, placebo subjects became sensitized to the tone and were aroused by tones of lower dB level. On the other hand, for triazolam subjects, the tone did not become more salient with repeated exposure and, in fact, their arousal threshold was significantly elevated above placebo values during SWS.

The focus of recent evaluations of sleep medications has been on performance effects and how long these effects persist, as well as on hypnotic efficacy. Although triazolam did not produce morning performance decrements in our study, acute effects on performance during arousals from sleep, on arousal threshold, and memory were evident. While the predictive validity of performance on laboratory tasks to real-world performance may be debated, our findings indicate that poor sleepers who take triazolam at bedtime may not respond as readily to salient signals which occur during sleep, such as smoke detectors or other alarms. Our data suggest that behaviors performed during arousals from sleep during the night, such as taking telephone calls or ingesting medications (as has been discussed by Roth et al.²²), may be forgotten.

REFERENCES

1. Roth, T., M. Kramer, & J.L. Schwartz. Triazolam: A sleep laboratory study of a new benzodiazepine hypnotic. *Current Therapeutic Research*, 1974, 16: 117-123.
2. Roth, T., M. Kramer, & T. Lutz. The effects of hypnotics on sleep, performance, and subjective state. *Drugs Under Experimental & Clinical Research*, 1977, 1: 279-286.
3. Roth, T., M. Kramer, & T. Lutz. The effects of triazolam (.25 mg.) on the sleep of insomniac subjects. *Drugs Under Experimental & Clinical Research*, 1977, 1: 271-277.
4. Vogel, G., A. Thurmond, P. Gibbons, K. Edwards, K.B. Sloan, & K. Sexton. The effect of triazolam on the sleep of insomniacs. *Psychopharmacologia (Berlin)*, 1975, 41: 65-69.
5. Vogel, G.W., K. Barker, P. Gibbons, & A. Thurmond. A comparison of the effects of flurazepam 30 mg and triazolam 0.5 mg on the sleep of insomniacs. *Psychopharmacology*, 1976, 47: 81-86.
6. Kales, A., J.D. Kales, E.O. Bixler, M.B. Scharf, & E. Russek. Hypnotic efficacy of triazolam: Sleep laboratory evaluation of intermediate-term effectiveness. *Journal of Clinical Pharmacology*, 1976, 16: 399-406.
7. Nicholson, A.N. & B.M. Stone. Activity of the hypnotics, flunitrazepam and triazolam, in man. *British Journal of Clinical Pharmacology*, 1980, 9: 187-194.
8. Ogura, C., K. Nakazawa, K. Majima, K. Nakamura, H. Ueda, Y. Umezawa, & W.M. Wardell. Residual effects of hypnotics: Triazolam, flurazepam, and nitrazepam. *Psychopharmacology*, 1980, 68: 61-65.
9. Pegram, V., P. Hyde, & P. Linton. Chronic use of triazolam: The effects on the sleep patterns of insomniacs. *Journal of International Medical Research*, 1980, 8: 224-231.
10. Veldkamp, W., R.N. Straw, C.M. Metzler, & H.V. Demissianos. Efficacy and residual effect evaluation of a new hypnotic, triazolam. *Journal of Clinical Pharmacology*, 1974, 14: 102-111.
11. Fabre, L.F., L. Gross, V. Passignajen, & C. Metzler. Multiclinic double-blind comparison of triazolam and flurazepam for seven nights in outpatients with insomnia. *Journal of Clinical Pharmacology*, 1977, 17: 402-409.
12. Fabre, L.F. & W.T. Smith. Multi-clinic cross-over comparison of triazolam (Halcion®) and placebo in the treatment of co-existing insomnia and anxiety in anxious out-patients. *Diseases of the Nervous System*, 1977, 38: 487-491.
13. Leibowitz, M. & A. Sunshine. Long-term hypnotic efficacy and safety of triazolam and flurazepam. *Journal of Clinical Pharmacology*, 1978, 18: 302-309.
14. Nair, N.P.V. & G. Schwartz. Triazolam in insomnia: A standard-controlled trial. *Current Therapeutic Research*, 1978, 23: 388-392.
15. Deberdt, R. Long-term use of triazolam, a benzodiazepine hypnotic. *Current Therapeutic Research*, 1979, 26: 1005-1014.
16. Sundaresan, P.R., W.M. Wardell, M. Weintraub, & L. Lasagna. Methodology for demonstrating sustained efficacy of hypnotics: A comparative study of triazolam and flurazepam. *Clinical Pharmacology & Therapeutics*, 1979, 25: 391-398.

17. Hindmarch, I. & C.A. Clyde. The effects of triazolam and nitrazepam on sleep quality, morning vigilance, and psychomotor performance. *Arzneimittel-Forschung/Drug Research*, 1980, 30(II): 1163-1166.
18. Piccione, P., F. Zorick, T. Lutz, T. Grissom, M. Kramer, & T. Roth. The efficacy of triazolam and chloral hydrate in geriatric insomniacs. *Journal of International Medical Research*, 1980, 8: 361-367.
19. Hare, S.A. A comparison of three benzodiazepine hypnotics as oral preanesthetic medication. *South African Medical Journal*, 1975, 49: 1883-1884.
20. Eberts, F.S., H. Ko, & R.C. Thomas. Metabolism and pharmacokinetics of triazolam. Cited in D.D. Breimer: Pharmacokinetics and metabolism of various benzodiazepines used as hypnotics. *British Journal of Clinical Pharmacology*, 1979, 8: 7S-13S.
21. Greenblatt, D.J., M. Divoll, L.J. Moschitto, & R.I. Shader. Electron-capture gas chromatographic analysis of triazolobenzodiazepines, alprazolam and triazolam. *Psychopharmacology*, in press.
22. Roth, T., K.M. Hartse, P.G. Saab, P.M. Piccione, & M. Kramer. The effects of flurazepam, lorazepam, and triazolam on sleep and memory. *Psychopharmacology*, 1980, 70: 231-237.
23. Rechtschaffen, A. & A. Kales (Eds.). A manual of standardized terminology, techniques and scoring system for sleep stages of human subjects. Washington, D.C.: Public Health Service, U.S. Government Printing Office, 1968.
24. Hoddes, E., V. Zarcone, H. Smythe, R. Phillips, & W.C. Dement. Quantification of sleepiness: A new approach. *Psychophysiology*, 1973, 10: 431-436.
25. Church, M.W. & L.C. Johnson. Mood and performance of poor sleepers during repeated use of flurazepam. *Psychopharmacology*, 1979, 61: 309-316.
26. Williams, H.L. & C.L. Williams. Nocturnal EEG profiles and performance. *Psychophysiology*, 1966, 3: 164-175.
27. Ekstrand, B.R. Effect of sleep on memory. *Journal of Experimental Psychology*, 1967, 75: 64-72.
28. Eaton, H.S. An English-French-German-Spanish Word Frequency Dictionary. New York: Dover Publications, 1961.
29. Johnson, L.C. & C.L. Spinweber. Effects of a short-acting benzodiazepine on brain electrical activity during sleep. *Electroencephalography & Clinical Neurophysiology*, in press.
30. Kales, A., M.B. Scharf, & J.D. Kales. Rebound insomnia: A new clinical syndrome. *Science*, 1978, 201: 1039-1041.
31. Nicholson, A.N., B.M. Stone, & P.A. Pascoe. Studies on sleep and performance with a triazolo-1, 4-thienodiazepine (brotizolam). *British Journal of Clinical Pharmacology*, 1980, 10: 75-81.
32. Clarke, P.R.F., P.S. Eccersley, J.P. Frisby, & J.A. Thornton. The amnesic effect of diazepam (Valium). *British Journal of Anaesthesia*, 1970, 42: 690-697.
33. Pandit, S.K., J.W. Dundee, & S.R. Keilty. Amnesia studies with intravenous premedication. *Anaesthesia*, 1971, 26: 421-428.

34. Heisterkamp, D.V. & P.J. Cohen. The effect of intravenous premedication with lorazepam (Ativan), pentobarbitone or diazepam on recall. *British Journal of Anaesthesia*, 1975, 47: 79-81.
35. Dundee, J.W. & D.A. George. The amnesic action of diazepam, flunitrazepam and lorazepam in man. *Acta Anaesthesiologia Belgica*, 1976, [Suppl.] 27: 3-11.
36. Bixler, E.O., M.B. Scharf, C.R. Soldatos, D.J. Mitsky, & A. Kales. Effects of hypnotic drugs on memory. *Life Sciences*, 1979, 25: 1379-1388.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 81-1	2. GOVT ACCESSION NO. AD-A101785	3. RECIPIENT'S CATALOG NUMBER (1)
4. TITLE (and Subtitle) (U) Effects of Triazolam (0.5 mg) on Sleep, Performance, Memory, and Arousal Threshold	5. TYPE OF REPORT & PERIOD COVERED Interim rept.	
6. PERFORMING ORG. REPORT NUMBER		7. AUTHOR(s) Cheryl L. SPINWEBER and Laverne C. JOHNSON
8. CONTRACT OR GRANT NUMBER(s)		9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Health Research Center P.O. Box 85122 San Diego, CA 92138
10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 17 MR041 01 003-0157		11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research & Development Command National Naval Medical Center Bethesda, MD 20014
12. REPORT DATE 11 March 1981		13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Bureau of Medicine and Surgery Department of the Navy Washington, DC 20372 12 24		15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 16 MR041 01		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 14 NAHLTHRSCHC-81-1		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Benzodiazepines Anterograde amnesia Triazolam Arousal threshold Humans Sleep Performance		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) (U) The effects of a short-acting benzodiazepine hypnotic, triazolam (0.5 mg), on sleep, performance, and arousal threshold were assessed in 20 male poor sleepers (age 21 ± 2.37 years). Following a laboratory screening night, all subjects received placebo for 3 nights (single-blind), 10 received triazolam and 10 placebo for 6 nights (double-blind), and all received placebo on 2 withdrawal nights (single-blind). All effects described below were statistically significant. Triazolam reduced sleep latency and increased total sleep time and sleep (continued on reverse side)		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

391642

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. ABSTRACT (continued)

efficiency. Stage 2% was increased and Stage 4% was reduced during treatment. Morning performance, measured 8.25 hours post-drug, showed no decrements. Acute effects were assessed on the 6th treatment night during arousals from sleep at 1.5, 3, and 5 hours post-administration: performance was impaired in triazolam subjects on the Wilkinson 4-Choice Reaction Time Test, Digit Symbol Substitution Test, Williams Word Memory Test, and Card Sorting Task. In the morning following the 6th treatment night, long-term memory was tested using a recognition task requiring subjects to identify words presented during nighttime test batteries: triazolam subjects correctly identified fewer target words. Triazolam administration produced anterograde amnesic effects. But, in a Paired Associates Test learned prior to drug ingestion on the previous evening, triazolam did not impair morning recall of word pairs. Threshold for arousal from slow wave sleep was elevated during treatment, and triazolam subjects did not show increased sensitivity to the arousing tone over nights as did placebo subjects.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)