

AD-A101 991

PURDUE UNIV LAFAYETTE IN DEPT OF STATISTICS

F/G 12/1

ASYMPTOTIC (BETA) SUB D JOINT NORMALITY OF SAMPLE QUANTILES.(U)

JUN 81 S IKEDA

N00014-75-C-0455

UNCLASSIFIED

MMS-81-20

NL

1 of 1

20

20/1/81




END
DATE
FILMED
8 81
DTIC

AD A101991

LEVEL

(1)

PURDUE UNIVERSITY



DTIC
ELECTE
JUL 24 1981

DEPARTMENT OF STATISTICS

DIVISION OF MATHEMATICAL SCIENCES

DTIC FILE COPY

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

81 7 21 040

(13)

(b) (a)
ASYMPTOTIC (B) JOINT NORMALITY
OF SAMPLE QUANTILES

By

(10) Sadao Ikeda

Department of Economics, Soka University
and
Department of Statistics, Purdue University

9 Mimeograph Series, 81-20 (14) 11/12-1

DTIC
ELECTE
JUL 24 1981

Department of Statistics
Division of Mathematical Sciences

(11) June 1981

(15) 16

*) The writing of this paper was facilitated while the author was
visiting Purdue University in part under the ONR Contract N00014-75-C-0455

(13)

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

371730

15

ASYMPTOTIC $(\mathfrak{B})_d$ JOINT NORMALITY OF SAMPLE QUANTILES*

By

Sadao Ikeda
 Dept. of Economics, Soka Univ., and Dept. of Statistics, Purdue Univ.

Summary

Notions of asymptotic equivalence of probability distributions and some of their properties are briefly presented. By applying the results on type $(\mathfrak{B})_d$ asymptotic equivalence, asymptotic $(\mathfrak{B})_d$ joint normality of a set of increasing number of sample quantiles are discussed, which improves and refines the previous work by Ikeda and Matsunawa (1972).

1. Introduction

Consider two sequences of random variables $\{X_t; t \rightarrow t_0\}$ and $\{Y_t; t \rightarrow t_0\}$, X_t and Y_t belonging to $\mathcal{P}(R_t, \mathfrak{B}_t)$ for each t , where t is a parameter taking values in a given metric space, R_t is any given abstract space, $\mathfrak{B}_t$ is a σ -field of subsets of R_t , and finally $\mathcal{P}(R_t, \mathfrak{B}_t)$ designates the class of all random variables distributed over the measurable space $(R_t, \mathfrak{B}_t)$.

Let us consider a sequence, $\{C_t; t \rightarrow t_0\}$, of subclasses of corresponding R_t 's, and denote it by (C) .

*The writing of this paper was facilitated while the author was visiting Purdue University in part under the ONR Contract N00014-75-C-0455.

Keywords: Asymptotic theory, order statistics, normal distribution.

Accession for	
NTIS GRA&I	☐
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Special	
Dist	

Under the above situation, two kinds of notions are defined:

Definition 1.1 (a) Two sequences of random variables $\{X_t; t \rightarrow t_0\}$ and $\{Y_t; t \rightarrow t_0\}$, are said to be asymptotically equivalent in the sense of type (C)_d, and briefly denoted as

$$(1.1) \quad X_t \sim Y_t(C)_d, (t \rightarrow t_0),$$

provided that the following condition holds:

$$(1.2) \quad \delta_d(X_t, Y_t; C_t) \equiv \sup_{E \in C_t} |P^{X_t}(E) - P^{Y_t}(E)| \rightarrow 0, (t \rightarrow t_0).$$

(b) The two sequences are said to be asymptotically equivalent in the sense of type (C)_r, and denoted as

$$(1.3) \quad X_t \sim Y_t(C)_r, (t \rightarrow t_0),$$

provided that

$$(1.4) \quad \delta_r(X_t, Y_t; C_t) \equiv \sup_{E \in C_t} \left| \frac{P^{X_t}(E)}{P^{Y_t}(E)} - 1 \right| \rightarrow 0, (t \rightarrow t_0)$$

In case where $(R_t, \mathcal{B}_t)$ and C_t are fixed independently of t , a weaker notion of asymptotic equivalence is defined corresponding to each types of the above definition. Let $(R_t, \mathcal{B}_t) = (R, \mathcal{B})$ and $C_t = C$.

Defintion 1.2. (a) $\{X_t; t \rightarrow t_0\}$ and $\{Y_t; t \rightarrow t_0\}$ are said to be asymptotically equivalent in the sense of type ((C))_d, and denoted as

$$(1.5) \quad X_t \sim Y_t((C))_d, (t \rightarrow t_0),$$

provided that

$$(1.6) \quad |P^{X_t}(E) - P^{Y_t}(E)| \rightarrow 0, (t \rightarrow t_0)$$

for each E belonging to C .

(b) The sequences are said to be asymptotically equivalent in the sense of type $((C))_r$, and denoted as

$$(1.7) \quad X_t \sim Y_t ((C))_r, (t \rightarrow t_0),$$

provided that

$$(1.8) \quad \left| \frac{P^X_t(E)}{P^Y_t(E)} - 1 \right| \rightarrow 0, (t \rightarrow t_0)$$

for each E belonging to $\mathcal{C}$.

Each of the four types of asymptotic equivalence defines the corresponding type of notions of asymptotic independence and convergence. (see Ikeda 1963, 1968).

Among various types of asymptotic equivalence defined above, type $(\mathcal{B})_d$ asymptotic equivalence enjoys very nice properties. By Def. 1.1, $X_t \sim Y_t (\mathcal{B})_d$, $(t \rightarrow t_0)$, if and only if

$$(1.9) \quad \delta_d(X_t, Y_t; \mathcal{B}_t) \equiv \sup_{E \in \mathcal{B}_t} |P^{X_t}_t(E) - P^{Y_t}_t(E)| \rightarrow 0, (t \rightarrow t_0).$$

First we state the following:

Proposition 1.1 Let $\varphi_t(z)$ be any given measurable transformation from $(R_t, \mathcal{B}_t)$ to $(\bar{R}_t, \bar{\mathcal{B}}_t)$ for each t, and put $\bar{X}_t = \varphi_t(X_t)$ and $\bar{Y}_t = \varphi_t(Y_t)$. Then, $X_t \sim Y_t (\mathcal{B})_d$ implies $\bar{X}_t \sim \bar{Y}_t (\bar{\mathcal{B}})_d$, as $t \rightarrow t_0$. If, in particular, both the spaces are identical and φ_t is non-singular, then $X_t \sim Y_t (\mathcal{B})_d$ is equivalent to $\bar{X}_t \sim \bar{Y}_t (\mathcal{B})_d$, as $t \rightarrow t_0$.

Proposition 1.2 Let $h_t(z)$ be a real, single valued function defined over R_t , for each t. If $\mathcal{E}|h_t(X_t)|^{1+\delta}$ and $\mathcal{E}|h_t(Y_t)|^{1+\delta}$ are both uniformly bounded in t for some $\delta > 0$, then the condition $X_t \sim Y_t (\mathcal{B})_d$, $(t \rightarrow t_0)$, implies that $|\mathcal{E} h_t(X_t) - \mathcal{E} h_t(Y_t)| \rightarrow 0$, $(t \rightarrow t_0)$, where in general $\mathcal{E} h(X) = \int_R h d P^X$.

Some useful criteria for type $(\mathcal{B})_d$ asymptotic equivalence have been given:

Proposition 1.3 (a) For $X_t \sim Y_t (\mathcal{B})_d$, $(t \rightarrow t_0)$, it is sufficient that either one

of the following conditions holds:

$$(1.10) \quad \min\{I(X_t:Y_t;A_t), I(Y_t:X_t;A_t)\} \rightarrow 0, (t \rightarrow t_0).$$

(1.11) $\min\{W(X_t:Y_t;A_t), W(Y_t:X_t;A_t)\} \rightarrow 0, (t \rightarrow t_0)$, where $\{A_t; t \rightarrow t_0\}$ is a sequence of subsets such that $P^{X_t}(A_t) \rightarrow 1$ or $P^{Y_t}(A_t) \rightarrow 1$ as $t \rightarrow t_0$, and I and W are defined by

$$(1.12) \quad I(X_t:Y_t;A_t) = \int_{A_t} f_t \log \frac{f_t}{g_t} dv_t,$$

and

$$(1.13) \quad W(X_t:Y_t;A_t) = \int_{A_t} g_t \left(\frac{f_t}{g_t} - 1 \right)^2 dv_t.$$

(b) For $X_t \sim Y_t (B)_d, (t \rightarrow t_0)$, it is necessary and sufficient that

$$(1.14) \quad \rho(X_t, Y_t; A_t) \equiv \int_{A_t} \sqrt{f_t g_t} dv_t \rightarrow 0, (t \rightarrow t_0)$$

where $\{A_t, t \rightarrow t_0\}$ is the same as in (a) above.

Evaluation of approximation error, $\delta_d(X_t, Y_t; B_t)$, can be done by using either one of the following inequalities:

Proposition 1.4

$$(1.15) \quad 0 \leq 1 - \rho(X_t, Y_t; R_t) \leq \delta_d(X_t, Y_t; B_t) \leq \{1 - \rho^2(X_t, Y_t; R_t)\}^{1/2}$$

$$\leq \{\min\{I(X_t:Y_t;R_t), I(Y_t:X_t;R_t)\}\}^{1/2}$$

$$\leq \{\min\{\log(1+W(X_t:Y_t;R_t)), \log(1+W(Y_t:X_t;R_t))\}\}^{1/2}$$

$$(1.16) \quad \delta_d(X_t, Y_t; B_t) \leq \left\{ \frac{3}{4} \left(\left[1 + \frac{4}{3} \min\{I(X_t:Y_t;R_t), I(Y_t:X_t;R_t)\} \right]^{1/2} - 1 \right) \right\}^{1/2}$$

$$\leq \left(\frac{3}{8} \min\{I(X_t:Y_t;R_t), I(Y_t:X_t;R_t)\} \right)^{1/2}.$$

Recently, Matsunawa and Ikeda (1981) have obtained a necessary and sufficient condition for $X_t \sim Y_t(B)_d, (t \rightarrow t_0)$ to hold in terms of the quantity $I(X_t:Y_t;A_t)$.

In the following section, we shall specialize the notions of asymptotic equivalence of type $(C)_d$ and of type $((C))_d$ to a special set of subclasses in real case, which are of common interest in statistics.

2. Asymptotic equivalence in real case.

Let $\{X_{s(n_s)}; s \rightarrow \infty\}$ and $\{Y_{s(n_s)}; s \rightarrow \infty\}$ be two sequences of real random variables, $X_{s(n_s)}$ and $Y_{s(n_s)}$ belonging to $\rho(R_{(n_s)}, \mathcal{B}_{(n_s)})$, the class of all random variables distributed over the n_s -dimensional Euclidean space, $R_{(n_s)}$, with the usual Borel field $\mathcal{B}_{(n_s)}$.

We take the following sequences of subclasses:

$$(2.1) \quad \{\mathcal{M}_{(n_s)}; s=1,2,\dots\}, \{\mathcal{S}_{(n_s)}; s=1,2,\dots\}, \{\mathcal{A}_{(n_s)}; s=1,2,\dots\}, \\ \{\mathcal{L}_{(n_s)}; s=1,2,\dots\} \text{ and } \{\mathcal{B}_{(n_s)}; s=1,2,\dots\},$$

where we have defined:

$$(2.2) \quad \mathcal{M}_{(n)} = \{Z_{(n)} \mid -\infty < Z_i < b_i; i = 1,2,\dots,n; b_i: \text{extended real}\}, \\ \mathcal{S}_{(n)} = \{Z_{(n)} \mid a_i \leq Z_i < b_i; i = 1,2,\dots,n; a_i, b_i: \text{extended real}\}, \\ \mathcal{A}_{(n)} = \left\{ \sum_{j=1}^N E_j(\text{disjointsum}) \mid E_j \in \mathcal{S}_{(n)}; j=1,\dots,N; N: \text{any positive integer} \right\}, \\ \mathcal{L}_{(n)} = \text{class of all open subsets of } R_{(n)}.$$

Among the five types of asymptotic equivalence, $(\mathcal{M})_d, (\mathcal{S})_d, (\mathcal{A})_d, (\mathcal{L})_d$ and $(\mathcal{B})_d$, corresponding to the sequences in (2.1), the following implication relations hold:

In case of general basic spaces,

$$(2.3) \quad \{(\mathcal{B})_d, (A)_d, (\mathcal{L})_d\} \xrightarrow{\rightarrow} \{(\mathcal{S})_d\} \xrightarrow{\rightarrow} \{(\mathcal{M})_d\},$$

and in case of identical (or equal) basic spaces where $(R_{(n_s)}, \mathcal{B}_{(n_s)}) = (R_{(n)}, \mathcal{B}_{(n)})$ for all s ,

$$(2.4) \quad \{(\mathcal{B})_d, (A)_d, (\mathcal{L})_d\} \xrightarrow{\rightarrow} \{(\mathcal{S})_d, (\mathcal{M})_d\}$$

where $\{ \}$ designates a group of equivalent notions, $\rightarrow$ "imply", and $\nrightarrow$ "not necessarily imply, counter example shown".

Among the five weaker types of notions, the following implication diagram is obtained:

$$(2.5) \quad \{((\mathcal{B}))_d\} \rightarrow \{((\mathcal{L}))_d\} \xrightarrow{\rightarrow} \{((A))_d, ((\mathcal{S}))_d, ((\mathcal{M}))_d\}.$$

Some conditional implication relations have been also obtained (see Ikeda 1968). Note that type $((\mathcal{M}))_d$ convergence is equivalent to the usual in law convergence, provided the limiting distribution is of continuous type, in which case, it is also shown that $((\mathcal{M}))_d$ and $(\mathcal{M})_d$ are mutually equivalent, as in the Central Limit Theorem. Also, in many cases of in law convergence, it turns out that the convergence is of type $(\mathcal{B})_d$.

In the following section, we shall apply the type $(\mathcal{B})_d$ asymptotic equivalence notion to the asymptotic normality of a set of increasing number of selected order statistics.

3. Asymptotic $(\mathcal{B})_d$ joint normality of sample quantiles.

Mosteller (1946) has shown, under mild conditions, that given a spacing $0 < \lambda_1 < \dots < \lambda_k < 1$, the corresponding set of order statistics,

$X_{n,n_1} < X_{n,n_2} < \dots < X_{n,n_k}$, $n_i = [\lambda_i n] + 1$, are asymptotically jointly normally distributed, in a sense of type $((\mathcal{M}))_d$ in our present terminology, with mean vector $(F^{-1}(\lambda_1), \dots, F^{-1}(\lambda_k))$ and covariance matrix

$$(3.1) \quad \frac{1}{n} \begin{bmatrix} \frac{\lambda_1(1-\lambda_1)}{f_1^2} & \frac{\lambda_1(1-\lambda_2)}{f_1 f_2} & \dots & \frac{\lambda_1(1-\lambda_k)}{f_1 f_k} \\ & \frac{\lambda_2(1-\lambda_2)}{f_2^2} & \dots & \frac{\lambda_2(1-\lambda_k)}{f_2 f_k} \\ & & \ddots & \vdots \\ & & & \frac{\lambda_k(1-\lambda_k)}{f_k^2} \end{bmatrix}, \quad f_i = f(F^{-1}(\lambda_i)), i=1, \dots, k$$

Later on, Weiss (1969) has tried to derive an asymptotic distribution, in a strong sense, of a set of increasing number of sample quantiles, and has proved a result, which is a special case of Ikeda and Matsunawa (1972).

Now, we begin with the case of uniform distribution over the unit interval, $(0,1)$. Let $U_{n,1} < U_{n,2} < \dots < U_{n,n}$ be order statistics based on a random sample of size n drawn from a uniform distribution over $(0,1)$. Select k order statistics, $U_{n,n_1} < U_{n,n_2} < \dots < U_{n,n_k}$, and put

$$(3.2) \quad U_{n(k)} = (U_{n,n_1}, U_{n,n_2}, \dots, U_{n,n_k})';$$

where k and $(n_1, n_2, \dots, n_k)$ may depend on n as $n \rightarrow \infty$.

By applying the criterion (1.10), Ikeda and Matsunawa (1972) have shown that the following theorem holds:

Theorem 3.1 If the condition

$$(3.3) \quad w_n \equiv \frac{k}{\min_{1 \leq i \leq k+1} (n_i - n_{i-1})} \rightarrow 0, \quad (n \rightarrow \infty),$$

holds, then $U_{n(k)}$ and $Z_{n(k)}$ are asymptotically equivalent in the sense of type $(B)_d$ as $n \rightarrow \infty$, where $Z_{n(k)}$ stands for a normal random variable with mean vector

$$(3.4) \quad \ell_{n(k)} = (\ell_{n1}, \ell_{n2}, \dots, \ell_{nk})', \text{ with } \ell_{ni} = \frac{n_i}{n+1}, \quad i=1, 2, \dots, k,$$

and covariance matrix

$$(3.5) \quad L_{n(k)} = \frac{1}{n+2} \begin{bmatrix} \ell_{n1}(1-\ell_{n1}) & \ell_{n1}(1-\ell_{n2}) & \dots & \ell_{n1}(1-\ell_{nk}) \\ & \ell_{n2}(1-\ell_{n2}) & \dots & \ell_{n2}(1-\ell_{nk}) \\ & * & \dots & \ell_{nk}(1-\ell_{nk}) \end{bmatrix}.$$

Here we have taken a convention $n_0=0$, $n_{k+1}=n+1$.

In this case an upper bound for the quantity $\delta_d(U_{n(k)}, Z_{n(k)}; B_{(k)})$ is given by (1.16):

$$(3.6) \quad \delta_d(U_{n(k)}; Z_{n(k)}; B_{(k)}) \leq \left(\frac{3}{8} I(U_{n(k)}; Z_{n(k)}; R_{(k)}) \right)^{1/2},$$

with

$$(3.7) \quad I(U_{n(k)}; Z_{n(k)}; R_{(k)}) = w_n + o(w_n).$$

The result obtained in Theorem 3.1 is, of course, extended to a uniform distribution over any given finite interval, by virtue of Prop.1.1.

Ikeda and Matsunawa (1972) have also obtained a result in case of general basic distribution. However, the conditions there are somewhat gloomy and not convenient to practical use. Moreover, the conditions include a stronger condition for the spacing of $(n_1, n_2, \dots, n_k)$ than (3.3):

$$(3.8) \quad \frac{k^2}{\min_{1 \leq i \leq k+1} (n_i - n_{i-1})} \rightarrow 0, \quad (n \rightarrow \infty).$$

Recently, Ikeda and Nonaka (1981) have obtained a refined result, which improves the earlier one.

Let $X_{n,1} < X_{n,2} < \dots < X_{n,n}$ be order statistics from a continuous distribution over the real line, whose pdf. and cdf. being given by $f(x)$ and $F(x)$, respectively. Choose k out of the order statistics, and put

$$(3.9) \quad X_{n(k)} = (X_{n,n_1}, X_{n,n_2}, \dots, X_{n,n_k})'$$

We shall first consider the case where the support of $f(x)$ is identical with the entire real line: $D_f = (-\infty, \infty)$.

Then, the transformed variable

$$(3.10) \quad F(X_{n(k)}) = (F(X_{n,n_1}), F(X_{n,n_2}), \dots, F(X_{n,n_k}))'$$

is identically distributed with $U_{n(k)}$ given in (3.2), or,

$$(3.11) \quad F^{-1}(U_{n(k)}) = (F^{-1}(U_{n,n_1}), F^{-1}(U_{n,n_2}), \dots, F^{-1}(U_{n,n_k}))'$$

is identically distributed with $X_{n(k)}$.

Let us consider a truncation of $Z_{n(k)}$ over the domain $A_{(k)} = \{z_{(k)} | 0 < z_i < 1; i = 1, 2, \dots, k\}$, and denote it by $Z_{n(k)}^*$. It is then evident that, under the condition (3.3), $Z_{n(k)} \sim Z_{n(k)}^* (B)_d, (n \rightarrow \infty)$. Further, let us put

$$(3.12) \quad Y_{n(k)}^* = F^{-1}(Z_{n(k)}^*) \equiv (F^{-1}(Z_{n,1}^*), F^{-1}(Z_{n,2}^*), \dots, F^{-1}(Z_{n,k}^*))',$$

and finally let $Y_{n(k)}$ be a normal random variable with mean vector

$$(3.13) \quad s_{n(k)} = (s_{n1}, s_{n2}, \dots, s_{nk})', \text{ with } s_{ni} = F^{-1}(\ell_{ni}), i=1, 2, \dots, k,$$

and covariance matrix

$$(3.14) \quad S_{n(k)} = \frac{1}{n+2} \begin{bmatrix} \frac{\ell_{n1}(1-\ell_{n1})}{f_{n1}^2} & \frac{\ell_{n1}(1-\ell_{n2})}{f_{n1}f_{n2}} & \dots & \frac{\ell_{n1}(1-\ell_{nk})}{f_{n1}f_{nk}} \\ * & \frac{\ell_{n2}(1-\ell_{n2})}{f_{n2}^2} & \dots & \frac{\ell_{n2}(1-\ell_{nk})}{f_{n2}f_{nk}} \\ & & & \frac{\ell_{nk}(1-\ell_{nk})}{f_{nk}^2} \end{bmatrix}$$

$$= D^{-1}(f_{n1}, \dots, f_{nk}) L_{n(k)} D^{-1}(f_{n1}, \dots, f_{nk}),$$

where we have put $f_{ni} = f(s_{ni}) = f(F^{-1}(\ell_{ni}))$, $i=1, 2, \dots, k$.

The following diagram indicate the relations among the variables thus defined:

$$(3.15) \quad \begin{array}{ccc} U_{n(k)} & \overset{(B)_d}{\sim} & Z_{n(k)} \overset{(B)_d}{\sim} Z_{n(k)}^* \\ F^{-1} \searrow & & \swarrow F^{-1} \\ X_{n(k)} & \overset{(B)_d}{\sim} & Y_{n(k)}^* \overset{(B)_d}{\sim} Y_{n(k)} \end{array}$$

Under the condition (3.3), it holds that $U_{n(k)} \sim Z_{n(k)}^* (\mathcal{B})_d$ and therefore, by Prop. 1.1, $X_{n(k)} \sim Y_{n(k)}^* (\mathcal{B})_d$, as $n \rightarrow \infty$. Hence, if one can show that $Y_{n(k)} \sim Y_{n(k)}^* (\mathcal{B})_d$, ($n \rightarrow \infty$), with possibly some additional conditions, then it holds that $X_{n(k)} \sim Y_{n(k)} (\mathcal{B})_d$, i.e., $X_{n(k)}$ would be distributed as normal with mean vector $s_{n(k)}$ and covariance matrix $S_{n(k)}$.

Ikeda and Nonaka (1981) investigated the quantity $I(Y_{n(k)}; Y_{n(k)}^*; R_{(k)})$ for criticizing the type $(\mathcal{B})_d$ asymptotic equivalence between $Y_{n(k)}$ and $Y_{n(k)}^*$, to get the following theorem.

Theorem 3.2 Suppose that the following assumptions are fulfilled:

- (i) The support of $f(x)$ is identical to the entire real line: $D_f = (-\infty, \infty)$.
- (ii) $f(x)$ is twice differentiable and $f''(x)$ is bounded and continuous over the entire real line.
- (iii) The function, $\varphi(x) \equiv \{f(x)f''(x) - f'(x)^2\} / f(x)^2$, is bounded uniformly for all x in $(-\infty, \infty)$.

Then, in order that $X_{n(k)} \sim Y_{n(k)} (\mathcal{B})_d$, ($n \rightarrow \infty$), it is sufficient that the following conditions are satisfied simultaneously:

$$(3.16) \quad w_n = \frac{k}{\min_{1 \leq i \leq k+1} (n_i - n_{i-1})} \rightarrow 0, \quad (n \rightarrow \infty),$$

and

$$(3.17) \quad w_n \cdot \max\left\{\sigma_n^4, \frac{\sigma_n^5}{\sqrt{n+2}}, \frac{\sigma_n^6}{n+2}\right\} \rightarrow 0, \quad (n \rightarrow \infty),$$

where we have put

$$(3.18) \quad \sigma_n^2 = \max \sigma_{ni}^2, \quad \sigma_{ni}^2 = \frac{\ell_{ni}(1 - \ell_{ni})}{f_{ni}^2}, \quad i=1, 2, \dots, k.$$

In case where k and x_{ni} 's are fixed, the conditions (3.16) and (3.17) are automatically fulfilled, in which case, however, the assumptions (i)–(iii) happen to be slightly stronger than those by Mosteller (1946). In this case, more direct calculation would be possible, which will be left open.

Ikeda and Nonaka (1981) have given a more general result than the above theorem, which states the following result:

Theorem 3.3 Suppose that the following assumptions are fulfilled:

- (i) The support of $f(x)$ is identical to an open interval: $D_f = (a, b)$, where a and b are extended real.
- (ii) $f(x)$ is twice differentiable and $f''(x)$ is bounded and continuous over (a, b) .
- (iii) The function, $\varphi(x) \equiv \{f(x)f''(x) - f'(x)^2\} / f(x)^2$, is bounded uniformly for all x in (a, b) .

Then, in order that $X_{n(k)} \sim Y_{n(k)} (\mathcal{B})_d$, $(n \rightarrow \infty)$, it is sufficient that the following conditions are satisfied simultaneously:

$$(3.19) \quad w_n \equiv \frac{k}{\min_{1 \leq i \leq k+1} (n_i - n_{i-1})} \rightarrow 0, \quad (n \rightarrow \infty).$$

$$(3.20) \quad w_n \cdot \left\{ \frac{4}{\sigma_n}, \frac{\sigma_n^5}{\sqrt{n+2}}, \frac{\sigma_n^6}{n+2} \right\} \rightarrow 0, \quad (n \rightarrow \infty).$$

and

$$(3.21) \quad w_n \cdot \max_{1 \leq i \leq k} \left[\frac{\sigma_{ni}^4}{\min\{|a-s_{ni}|^2, |b-s_{ni}|^2\}} \right] \rightarrow 0, \quad (n \rightarrow \infty).$$

It is evident that this theorem implies the result in the preceding theorem. Also, it should be noted that, in order to obtain an error estimation

for $\delta_d(X_{n(k)}, Y_{n(k)}; R_{(k)})$, an evaluation of the K-L information $I(X_{n(k)}; Y_{n(k)}; R_{(k)})$ should be done directly.

References

- Ikeda S. (1963). Asymptotic equivalence of probability distributions with applications to some problems of asymptotic independence. Ann. Inst. Stat. Math., Tokyo, 15, 87-116
- Ikeda S. (1968). Asymptotic equivalence of real probability distributions. Ann. Inst. Stat. Math., Tokyo, 20, 339-362.
- Ikeda, S. and Matsunawa, T. (1972). On the uniform asymptotic joint normality of sample quantiles. Ann. Inst. Stat. Math., Tokyo, 24, 435-449.
- Ikeda, S. and Nonaka, Y. (1981). Uniform asymptotic joint normality of a set of increasing number of sample quantiles. (not published yet).
- Matsunawa, T. and Ikeda, S. (1981). Necessary and sufficient conditions for type $(\theta)_d$ asymptotic equivalence with applications (not published yet).
- Mosteller, F. (1946). On the useful "inefficient" statistics. Ann. Math. Statist., 17, 377-407.
- Weiss, L. (1969). The asymptotic joint distribution of an increasing number of sample quantiles. J. Roy. Stat. Soc., B. 30, 570-575.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Mimeograph Series 81-20	2. GOVT ACCESSION NO. AD-A101 991	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Asymptotic $(\beta)_d$ Joint Normality of Sample Quantiles		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) Sadao Ikeda		6. PERFORMING ORG. REPORT NUMBER Mimeo. Series 81-20
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Economics, Soka University and Department of Statistics, Purdue University		8. CONTRACT OR GRANT NUMBER(s) ONR N0014-75-C-0455
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, D.C.		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1981
		13. NUMBER OF PAGES 14
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION, DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES YES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Asymptotic theory, order statistics, normal distribution		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Notions of asymptotic equivalence of probability distributions and some of their properties are briefly presented. By applying the results on type $(\beta)_d$ asymptotic equivalence, asymptotic $(\beta)_d$ joint normality of a set of increasing number of sample quantiles are discussed, which improves and refines the previous work by Ikeda and Matsunawa (1972).		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)