

AD-A102 232

ILLINOIS UNIV AT URBANA DEPT OF COMPUTER SCIENCE
A NOTE ON AN NP-COMPLETE PROBLEM, (U)
1979 J L LEWANDOWSKI

F/6 12/1

N00014-79-C-0775
NL

UNCLASSIFIED

1-1

2-1



END

DATE

FORMED

BY

DTIC

AD A102232

LEVEL II

①

⑥

A Note on an NP-Complete Problem

⑩

J. L. Lewandowski
Department of Computer Science
University of Illinois
Urbana, IL 61801

⑪ 1977

DTIC ELECTE
S JUL 30 1981 D

E
⑫ 5

We show in this note that the problem of two-coloring a graph $G = (V, E)$ with a distinguished vertex p and two distinguished subset $A(p)$ and $B(p)$ of V , such that there is a red connected subgraph containing $A + p$, and a blue connected subgraph containing $B + p$, is NP-Complete. We prove this by reduction of 3-satisfiability [Aho] by constructing, for a given boolean expression in disjunctive normal form with three variables in each term, a graph G with a distinguished vertex D and two subsets $A(D)$ and $B(D)$ such that the expression is satisfiable if and only if the graph has a two-coloring with the desired monochromatic subtrees.

This work was partially supported by the Office of Naval Research under Contract No. ONR-N00014-79-C-0775

⑮

DTIC FILE COPY

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

176014

81 7 30 06

In constructing our graph, for a given expression E in variables x_1, x_2, \dots, x_n there will be two types of configurations. The first one corresponds to a variable (Figure 1). For each variable, we construct one of these configurations and connect it to what will be our distinguished vertex D by an edge from the top vertex. In this construction, we let the edge on the left side of the configuration correspond to the un-complemented form x_i of the i th variable x_i and the right side correspond to the complemented form x'_i . Now, for each term t_j in the expression, we construct a copy of the second configuration (Figure 2) and again connect the top vertex of the configuration to D by an edge. Then we identify the three edges in the middle of the configuration to three of the middle edges which are in the "variable" configurations in the following way: If the i th term contains the three variable forms y_{ia}, y_{ib} , and y_{ic} : where, for example, y_{ia} is either the complemented or un-complemented form of some variable x_j , then we identify the edge y_{ia} to the corresponding edge in the x_j "variable" configuration. This implies that the two vertices connected to this edge are also identified everywhere the variable form y_{ia} appears.

So we have constructed a graph (Figure 3) which corresponds to the given expression. We have here drawn the edges in the two sides of the graph as disjoint for simplicity; but in the graph, we must remember that all of the edges which correspond to a particular form of a variable are identified. We note here that this construction is

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<i>in file</i>
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	




Figure 1 Variable




Figure 2 Term

clearly polynomial in time, (and also space). It should also be noted that our problem is clearly in NP, since if we are given an alleged solution, we can easily check to determine if all of the communication constraints are satisfied by performing a depth first search from D to see which vertices are reachable from it by blue and red paths, and then to check if the two distinguished subsets are included in the reachable vertices for the corresponding color.

We now form the two subsets of vertices in the following way: A (D) (or simply A) consists of all of the "round" vertices, and B (D) (or simply B) of the "square" ones. We note that none of these vertices is incident to an edge which has been identified.




Figure 3. The Corresponding Graph

"variable" configuration (say x_i) both of the edges are blue ("true") and if there is to be a red path from the "round" vertex V_i in the i th "variable" configuration to D , it must "sneak through" to D through the "expression" side of the graph. If we assume that it starts this "sneak path" through the term t_j , it must go through the vertex T_j since it can't go directly upwards because this edge is, by hypothesis, blue. This red path will isolate some "square" vertex by two red edges (and note that these are the only edges leading to this "square" vertex). However, this contradicts the fact that there is a blue path from this vertex to D . Therefore one of the edges in each "variable" configuration must be red.

It may be the case that both of these edges might be red, but this indicates that the variable has nothing to do with the truth value of the expression, it cannot appear in any blue path, and is a "don't care" which can have no effect on the satisfiability of the expression. This completes the proof to show that the graph has a suitable orientation if and only if the expression is satisfiable; and therefore, this edge-colorability problem is NP-Complete.