

AD-A103 467

WASHINGTON UNIV SEATTLE DEPT OF CHEMISTRY
HIGH TEMPERATURE GAS ENERGY TRANSFER.(U)
AUG 81 B S RABINOVITCH

F/G 7/4

N00014-75-C-0690

NL

UNCLASSIFIED

1 of 1
AD A
63467



END
DAIL
FILMED
10-81
DTIC

LEVEL

(12)

ANNUAL SUMMARY REPORT, 1 August 1980 - 31 July 1981

High Temperature Gas Energy Transfer

B. S. Rabinovitch, Principal Investigator
Department of Chemistry BG-10
University of Washington
Seattle, Washington 98195

Accepted

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0690

Project NR 092-549

9 August 1981

DTIC
ELECTE
S **D**
AUG 31 1981
A

DTIC FILE COPY

This document has been approved for public release; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government. This research was sponsored by the Office of Naval Research.

81-8 31 130

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A103467	
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
High Temperature Gas Energy Transfer.	Annual Summary, 1 Aug 1980-31 Jul 1981	
6. AUTHOR(s)	7. PERFORMING ORG. REPORT NUMBER	
B. S. Rabinovitch		
8. CONTRACT OR GRANT NUMBER(s)	9. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
N00014-75-C-0690 NR 092 549		
10. CONTROLLING OFFICE NAME AND ADDRESS	11. REPORT DATE	
Department of Chemistry BG-10 University of Washington Seattle, WA 98195	9 Aug 1981	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES	
14. SECURITY CLASS. (of this report)	15. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)		
This document has been approved for public release; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Diffusion Cloud Method Thermal System Energy Transfer Transients Gases Variable Encounter Method High Temperature Vibrational Energy Wall Activation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Work under this contract is summarized. Five technical reports have been issued. The effect of molecular parameters - vibrational eigenstate density and intermolecular attractive potential - on gas-surface vibrational energy transfer - has been documented. The VEM technique has been refined to single collision measurements that provide a more sensitive test of energy transfer models. Study of cyclopropane to 1300 K shows a decline in collision efficiency with rise of temperature. Strong collision behavior is found at low gas-surface temperatures.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102 LF 014 6601

Unclassified 370274
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

- I. Summary of Progress This report covers the period 1 August 1980 - 31 July 1981. The statement below summarizes the most important results and conclusions of the work in this period

1. Collisional efficiency in transfer of vibrational energy from a hot molecule to a surface (or another molecule) declines with increase in vibrational eigenstate density (i.e., with increase of number, and decrease in frequency, of the number of internal vibrational modes) of the hot molecule. However, paradoxically, the efficiency for collisional activation of reaction increases with increase in eigenstate density of the substrate molecule. These findings represent statistical effects and illustrate the statistical theory of energy accommodation of Lin and Rabinovitch. These matters are discussed in Technical Reports 16 and 17 for cyclopropane, methyl cyclopropane and cyclobutane, and include work in progress on cyclopropane-d₆.

2. In the higher temperature region, up to the highest temperature so far investigated, 1325 K, the following results are found and conclusions confirmed: a) Collisional efficiency for vibrational energy transfer declines with increase of temperature of the heat bath or surface acting as collision partner (TR17, TR18); b) the decline in efficiency is most marked at lower temperatures (below 1000 K) and itself decreases above that temperature (TR19). Unfortunately, the general behavior for polyatomics in the region 1000-2000 is, however, still a closed book even though such knowledge is essential for the understanding and design of combustion, shock, laser, etc. systems. The model for energy transfer — attractive interaction with statistical accommodation, as opposed to a repulsive interaction and an impulsive phenomenon — may change in this region.

3. The ordinary surface of flame and combustion systems behaves as a strong collider for vibrational energy transfer involving polyatomic molecules (TR18) at low temperatures (~ 450 K).

4. The Variable Encounter Method (VEM) has been refined to single-collision conditions. These are the first studies of single collisional activation of reacting polyatomic molecules (by energy transport, not catalysis). The method provides a much more refined probe of the form of the collision transition probability matrix elements p_{ij} (TR20).

Codes	
Dist	Avail and/or Special
A	

II. Complete listing of Technical Reports:

1. Technical Report TR01, 7 January 1976.

Vibrational Energy Transfer in Reaction Systems at Elevated Temperatures by B. S. Rabinovitch, D. G. Keil, J. F. Burkhalter and G. B. Skinner.

In this report a theoretical analysis was given of shock tube data for high temperature gas reactions. The data were analyzed to show that collisional energy transfer efficiency appears to decrease at high temperatures.

2. Technical Report TR02, 20 April 1976

Temperature Dependence of the Arrhenius Activation Energy. High Temperature Limit, J. F. Burkhalter and B. S. Rabinovitch.

In this report a conventional approximation to the high temperature vibrational partition function is shown to be defective, and the usual high temperature expression for the Arrhenius activation is in error. The variation of the activation energy for unimolecular reactions as a function of temperature is calculated for various representative systems.

3. Technical Report TR03, 15 October 1976

Intermolecular Vibrational Energy Transfer in Thermal Unimolecular Systems, D. C. Tardy and B. S. Rabinovitch.

This report gives a comprehensive and detailed survey and analysis of energy transfer in thermal gaseous systems. The present status of the field is delineated and future areas of work indicated.

4. Technical Report TR04, 15 April 1977

On the Use of Exact Vibrational State Counting Methods in RRKM Rate Calculations, S. E. Stein and B. S. Rabinovitch.

This report describes an algorithm for accurate state sum and density calculations and refutes an error in the literature.

5. Technical Report TR05, 15 August 1977

Vibrational Translational Energy Transfer in Atom-Polyatomic Molecule Collisions in Thermal Reaction Systems by I. Oref and B. S. Rabinovitch.

In this report a simple theoretical model describing energy transfer probabilities is given. Conservation of angular momentum was imposed. The importance of completeness and detailed balance conditions and correction of an earlier model in the literature are illustrated. The model is applied to experimental data.

6. Technical Report TR06, 1 September 1977

Problems of Diffusion in a Low Pressure Gas Stream as Related to the Polanyi Diffusion Flame (Cloud) Method by D. G. Keil, J. F. Burkhalter and B. S. Rabinovitch.

A theoretical and experimental study was made of flow and diffusion of a reactant in a low pressure gas stream. Connection is made with earlier theoretical analysis.

7. Technical Report TR07, 10 October 1978

Do Highly Excited Reactive Polyatomic Molecules Behave Ergodically?
I. Oref and B. S. Rabinovitch.

The literature is surveyed and analyzed. A variety of excitation techniques including crossed molecular beam, laser, chemical activation, photochemical and thermal are considered. The answer to the title question is "yes." The relationship to the present work on intermolecular transfer is the following: multiphoton laser experiments frequently involve collisional heating of the gas. In order to understand the results, one must have a good basis for interpretation of data. In this survey, some suggested models for energy relaxation by molecular collisions are criticized and data reinterpreted in light of results obtained in this laboratory.

8. Technical Report TR08. 1 November 1978

Vibrational Energy Transfer in Thermal Unimolecular Systems by the Diffusion Cloud Method. Cyclopropane. E. Kamaratos, J. D. Burkhalter, D. G. Keil and B. S. Rabinovitch.

A study of vibrational energy transfer by the Diffusion Cloud Method has been made at temperatures from 975 K to 1175 K in the cyclopropane isomerization system. H_2 , He, N_2 and CO_2 were studied as inert bath gases. Their relative efficiencies increase in that order. Values of $\langle \Delta E \rangle$ vary from 150 cm^{-1} to 1100 cm^{-1} at 975 K. It was shown that these values decline, and the collisional efficiencies, β_c , decrease markedly with rise of temperature to 1175 K. This is only the second measurement of such a phenomenon.

9. Technical Report TR09. 15 March 1979

Collisional Relaxation of Non-Equilibrium Vibrational Energy Distributions in a Thermal Unimolecular System. Surface Collisions. D. F. Kelley, B. D. Barton, L. Zalotai, and B. S. Rabinovitch.

A novel, simple technique, the Variable Encounter Method, is described for obtaining information on energy transfer efficiency between a gas and a solid in the transient region, and as a function of temperature. The method is as powerful as it is simple. Results for cyclopropane are described. Decrease

in collisional efficiency $\langle \Delta E \rangle$ with rise of temperature is confirmed, and the form of the transfer probability matrix $\tilde{P}$ is discovered. This method and these results are considered to be some of the most innovative and important that have ever been made in this field.

10. Technical Report TR10. 15 June 1979

Transients in Vibrational Relaxation of Polyatomic Molecules at High Temperatures by VEM. M. C. Flowers, D. F. Kelley, F. C. Wolters, and B. S. Rabinovitch.

Extension of the VEM technique to cyclobutane is described. An exponential form of $\tilde{P}$ is required to fit the data. The value of $\langle \Delta E \rangle$ declines with rise of temperature.

11. Technical Report TR11; 15 September 1979

Collisional Relaxation of Transient Vibrational Energy Distributions in a Thermal Unimolecular System. The Variable Encounter Method. D. F. Kelley, L. Zalotai and B. S. Rabinovitch.

The Variable Encounter Method permits the study of the transient in the relaxation of an initial vibrationally cold ensemble of molecules in a vibrationally hot distribution by a known and variable number of successive collisions with a hot wall. The system studied was the isomerization of 1,1-cyclopropane- d_2 with a fused quartz wall temperature of 800 K to 1175 K, and average number of collisions from 2.3 to 22.3. The detailed theory of the experiment was presented, together with the method of data deconvolution. Various modified gaussian and exponential models of energy transfer were found to give agreement with the data. The average down-step size was found to decline from $\leq 3500 \text{ cm}^{-1}$ at the lowest temperature to $\sim 2500 \text{ cm}^{-1}$ at the highest on the basis of a gaussian model, but heterogeneous energy transfer is more efficient than homogeneous gas-gas collisional transfer. A mathematical analysis of the relation between mean first-passage times and incubation times is given. Incubation times increase from ~ 7 to ~ 12 collisions with increasing temperature. Transient population distributions and the sequential reaction probabilities as a function of collision number are calculated.

12. Technical Report TR12. 10 October 1979

Vibrational Energy Transfer in a Diffusion-Flow Cyclopropane- d_2 System. J. F. Burkhalter, E. Kamaratos, and B. S. Rabinovitch.

In this study, absolute rate constants for isomerization of cyclopropane- d_2 and high temperature homogeneous energy transfer were determined in a diffusion-flow system. The relative rates of the competitive isotopic isomerization channels

were measured for two bath gases, N_2 and He, at two temperatures, 973 K and 1073 K. Values of the average energy down-jump size $\langle \Delta E \rangle$ were computed from both the absolute rates and the isotopic relative rates by suitable modelling by a stochastic calculation. The results confirm earlier diffusion cloud measurements.

13. Technical Report TR13, 10 October 1979

Transients in the Vibrational Excitation of Cyclobutane Decomposition Using the Variable Encounter Method. M. C. Flowers, F. C. Wolters, B. D. Barton and B. S. Rabinovitch.

The probability of reaction of cyclobutane molecules in a fixed-time interval after experiencing a known number of collisions with a hot surface at temperatures between 749 K and 1126 K was determined using the Variable Encounter Method. Calculations utilizing exponential or gaussian models for energy transfer enabled the average amounts of energy transferred for deactivating collisions, $\langle \Delta E' \rangle$, to be estimated. The exponential model fits the experimental data best and, using this model, $\langle \Delta E' \rangle$ is 2430 cm^{-1} at 748 K and decreases to 1470 cm^{-1} at 1123 K. Surface collisions are more efficient than binary gas collisions; but cyclobutane is a less efficient partner than cyclopropane- d_2 (TR No. 11). The results follow from a quasi-statistical model of energy accommodation.

14. Technical Report TR14, 1 December 1979

A Crucial Demonstration of Strong Collisional Behavior of Vibrational Energy for Gas-Surface Collisions. M. C. Flowers, F. C. Wolters, D. F. Kelley and B. S. Rabinovitch.

Application of the VEM technique was made to the thermal cyclobutene decomposition system. Vibrational energy transfer in the cyclobutene - seasoned quartz surface system was studied. Collisional efficiency again declines with rise of temperature, but the occurrence of strong collisions was demonstrated in crucial manner at temperatures below 450 K. The wall appears to behave like a theoretical strong collider below 450 K. This is a very novel finding.

15. Technical Report TR15, 31 March 1980

Gas-Surface Vibrational Energy Transfer in the Transient Region of a Low-Pressure Unimolecular Reaction. B. D. Barton, D. F. Kelley and B. S. Rabinovitch.

The Variable Encounter Method, for the study of gas-wall vibrational energy transfer in the transient region of a unimolecular reaction was applied to the

isomerization of cyclopropane to propylene. Temperatures in the range 900 K - 1125 K were employed. The average probability of reaction per collision, $\bar{P}_c(m)$, was deduced from the data and compared with a theoretical stochastic calculation based on both gaussian and exponential models for the energy transfer probabilities. The former model is more appropriate. The efficiency of a seasoned quartz wall was greater than gas-gas collisions of substrate and the efficiency declined with increase of temperature. The steady state is closely approached (90%) in a comparatively small number of gas-wall collisions - 10-20, approximately. The efficiency of transfer by cyclopropane is comparable with that by cyclopropane- d_2 (TR 11).

The following were submitted in the present report period.

16. Technical Report TR16, 1 August 1980

Collisional Relaxation of Vibrational Energy Transients in the Methyl Cyclopropane System. D. F. Kelley, T. Kasai and B. S. Rabinovitch.

The variable encounter method has been used to study vibrational energy transients in the isomerization of methylcyclopropane to various butenes. This molecule, which has a hindered methyl rotation mode, has the highest vibrational energy level density of any molecule - cyclopropane, cyclobutene, cyclobutane - that has been studied so far in testing the variation of energy transfer efficiency with molecular structure. This system was studied with reactor surface temperatures of 800 to 1130 K and average numbers of collisions per encounter with the reactor of 5.6 and 20.0. An exponential model of energy transfer was found to give the best fit to the data with the average down step energy $\langle \Delta E' \rangle$ decreasing from 1860 to 1415 cm^{-1} with increase of temperature over the range studied. Incubation times increased from 14 to 19 collisions with increase in temperature, and these times, together with values of the conventional relative collision efficiency β and values of $\langle \Delta E' \rangle$, are compared with those of other molecules studied by VEM. The calculated transient population distributions and the associated sequential reaction probabilities are also displayed. The energy transfer efficiency from the hot molecules in down-collisions is lowest in this system and hence the weakest model - exponential - gives a good fit. By contrast, the efficiency of up-transitions, i.e., energy transfer into the cold molecule measured in this work is higher than for other molecules. This is a consequence of the increased

eigenstate density and the important conclusion is reached that this verifies the suitability of a statistical model of energy accommodation to describe vibrational energy transfer involving high energy polyatomic molecules at these temperatures.

17. Technical Report TR17, 15 September 1980

A Reinvestigation of Transients in the Cyclopropane System by the Variable Encounter Method. M. C. Flowers, F. C. Wolters, D. F. Kelley and B. S. Rabinovitch.

The isomerization of cyclopropane to propene was reinvestigated under conditions identical to those used in a previously reported VEM study on cyclobutane. The simultaneous reaction of cyclobutane was also studied in the largest reactor. Some details of the method are amplified; the distribution functions for numbers of collisions in each reactor are displayed as are the contributions to reaction $R(n)$ as a function of the number of consecutive collision n . The present data are in essential agreement with those obtained by Kelley et al [TR 11], although the average size of an internal energy down transition ($\langle \Delta E' \rangle$) for cyclopropane molecules colliding with a hot surface is a little lower than previously estimated. The increasing efficiency of the surface in deactivating energized molecules as the surface temperature decreases is confirmed ($\langle \Delta E' \rangle = 2550 \text{ cm}^{-1}$ at 900 K, on a gaussian model for energy transfer, and $\langle \Delta E' \rangle = 2000 \text{ cm}^{-1}$ at $\sim 1100 \text{ K}$). The surface acts as a somewhat stronger collider for cyclopropane than for cyclobutane and is also a more efficient collider for cyclopropane than are gas-gas cyclopropane collisions. This accords with the results of TR16 and leads to the important conclusion that heterogeneous collision is more efficient than homogeneous collision, in general, for the deactivation of excited molecules.

18. Technical Report TR18, 15 October 1980

Transients in Thermal Isomerization of Cyclobutene by the Variable Encounter. Vibrational Energy Transfer and Relaxation at Lower Temperatures. F. C. Wolters, M. C. Flowers and B. S. Rabinovitch.

Vibrational energy transfer and relaxation in the thermal isomerization of cyclobutene to 1,3-butadiene was studied over the temperature range 429 K - 778 K using the Variable Encounter Method. Reactors characterized by various mean number of collisions, m , per encounter of a substrate molecule with the reactor ($m = 2.6, 5.9, 8.5$ and 27.2) were used. The

significant observation was made that strong collider behavior in collisions of substrate with a seasoned fused quartz wall is reached at lower temperatures (below 450 K), as evidenced by the coincidence of values of the probability of reaction per collision, $\bar{P}_c(m)$, obtained in VEM reactors having different m values. The average energy transferred to the wall per deactivating collision, $\langle \Delta E' \rangle$, increased strongly as temperature decreased, from $\langle \Delta E' \rangle = 2150 \text{ cm}^{-1}$ at 764 K to $> 6000 \text{ cm}^{-1}$ at 460 K. Various forms of models for the probability of energy transfer in down transitions were applied to data. The weak form — exponential — cannot apply, in contrast to the behavior that applies at higher temperatures for looser molecules (higher vibrational eigenstate densities) such as methyl cyclopropane (TR16).

19. Technical Report TR19, 15 March 1981.

Energy Transfer at High Temperatures in the Cyclopropane System.
Temperature Dependence of $\langle \Delta E' \rangle$. T. Kasai, D. F. Kelley and B. S. Rabinovitch.

The study of collisional energy transfer between cyclopropane molecules and a seasoned silica wall by the Variable Encounter Method was extended over a range of more than 500° upward in temperature to 1325 K. The value of $\langle \Delta E' \rangle$, the average energy of a down-transition upon collision of the hot substrate molecule with the wall to decrease monotonically. The data provide the most accurate and well-documented display for a polyatomic molecule of the temperature trend of $\langle \Delta E' \rangle$ presently available. The rate of decrease in $\langle \Delta E' \rangle$ seems to be more rapid at lower temperatures. The data have been explained by the theory of Lin and Rabinovitch on a quasi-statistical accommodation model.

The behavior at still higher temperatures is considered to be not well known and must be considered as in quite unsatisfactory condition. It is of extreme practical and theoretical importance to detect the possible transition of the observed behavior from an attractive potential-dominated accommodation phenomenon to a Landau-Teller type repulsive potential-dominated impulsive phenomenon.

20. Technical Report TR20

Single-Collision Gas-Surface Vibrational Energy Transfer in a Reacting System. D. F. Kelley, T. Kasai and B. S. Rabinovitch.

Gas-surface vibrational energy accommodation in a reactive system has been studied under single collision conditions for the first time for polyatomic molecules. The reaction system was the isomerization of cyclobutene to 1,3-butadiene. Both seasoned pyrex and silica surfaces were used over the temperature ranges 400-775 K and 500-975 K, respectively. Strong collider behavior was observed below ~ 425 K. The vibrational energy accommodation coefficient was found to decrease with rise of temperature from ~ 1.0 to 0.2, while the relative collisional efficiency β_1 declined from ~ 1.0 to 0.008. A stochastic calculation was used to fit the data. This technique provides a more sensitive test between the various model forms for energy transfer than has been possible heretofore. It is clearly shown that a Gaussian or particularly a Boltzmann weighted exponential form provide the best fit.

III. Listing of Publications.

1. Vibrational Energy Transfer in Reaction Systems at Elevated Temperatures, Proceed. Tenth Internat. Shock Tube Sympos. ed. G. Kamimoto, Kyoto, 1976.
2. Intermolecular Vibrational Energy Transfer in Thermal Unimolecular Systems. D. C. Tardy and B. S. Rabinovitch, Chem. Revs. 77, 369 (1977).
3. On the Use of Exact Vibrational State Counting Methods in RRKM Rate Calculations, S. E. Stein and B. S. Rabinovitch, Chem. Phys. Lett. 49, 183 (1977).
4. Vibrational Translational Energy Transfer in Atom-Polyatomic Molecule Collisions in Thermal Reaction Systems. I. Oref and B. S. Rabinovitch, Chem. Phys. 26, 385 (1977).
5. Problems of Diffusion in a Low Pressure Gas Stream as Related to the Polanyi Diffusion Flame (Cloud) Method, D. G. Keil, J. F. Burkhalter and B. S. Rabinovitch, J. Phys. Chem. 82, 355 (1978).
6. Vibrational Energy Transfer in Thermal Unimolecular Systems by the Diffusion Cloud Method. Cyclopropane. E. Kamaratos, J. F. Burkhalter, D. G. Keil and B. S. Rabinovitch, J. Phys. Chem. 83, 984 (1979).
7. Do Highly Excited Reactive Polyatomic Molecules Behave Ergodically? I. Oref and B. S. Rabinovitch, Accts. Chem. Res. 12, 166 (1979).
8. Collisional Relaxation of Non-equilibrium Vibrational Energy Distributions in a Thermal System. Surface Collisions, D. F. Kelley, B. D. Barton, L. Zalotai, and B. S. Rabinovitch, J. Chem. Phys. 71, 538 (1979).
9. Transients in Vibrational Relaxation of Polyatomic Molecules at High Temperatures by Variable Encounter Method. M. C. Flowers, B. D. Barton, F. C. Wolters and B. S. Rabinovitch. Am. Soc. Mass. Spectrom. Proceedings, ed. A. E. Harrison, 27, 661 (1979).
10. Collisional Relaxation of Transient Vibrational Energy Distributions in a Thermal Unimolecular System. The Variable Encounter Method. D. F. Kelley, L. Zalotai, and B. S. Rabinovitch. Chem. Phys. 46, 379 (1980).
11. Vibrational Energy Transfer in a Diffusion-Flow Cyclopropane-d₂ System. J. F. Burkhalter, E. Kamaratos, and B. S. Rabinovitch, J. Phys. Chem. 84 (1980) 476.
12. A Crucial Demonstration of Strong Collisional Behavior of Vibrational Energy for Gas-Surface Collisions. M. C. Flowers, F. C. Wolters, D. F. Kelley and B. S. Rabinovitch, Chem. Phys. Lett. 69, 543 (1980).
13. Transients in the Vibrational Excitation of Cyclobutane Decomposition Using the Variable Encounter Method. M. C. Flowers, F. C. Wolters, B. D. Barton and B. S. Rabinovitch, Chem. Phys. 47 (1980) 189.
14. Gas-Surface Vibrational Energy Transfer in the Transient Region of a Low-Pressure Unimolecular Reaction. B. D. Barton, D. F. Kelley, and B. S. Rabinovitch. J. Phys. Chem. 84, 1299 (1980).

15. Collisional relaxation of vibrational energy transients in the methylcyclopropane system. A variable encounter method study. D. F. Kelley, T. Kasai and B. S. Rabinovitch, 73, 5611 (1980).
16. Transients in Thermal Isomerization of Cyclobutene by VEM. Vibrational Energy Transfer and Relaxation at Lower Temperatures. J. Phys. Chem. 85 (1981) 589.
17. Surface-Gas Energy Transfer in the Cyclopropane/Cyclobutane Isomerization Systems by the Variable Encounter Method. M. C. Flowers, F. C. Wolters, D. F. Kelley, and B. S. Rabinovitch, J. Phys. Chem. 85 (1981) 849.
18. Single-Collision Gas-Surface Vibrational Energy Transfer in a Reacting System, D. F. Kelley, T. Kasai and B. S. Rabinovitch. J. Phys. Chem. 85 (1981) 1100.
19. Energy Transfer at High Temperatures in the Cyclopropane System. Temperature Dependence of $\langle \Delta E' \rangle$. T. Kasai, D. F. Kelley and B. S. Rabinovitch, Chem. Phys. Lett., in press.

IV. Current Work

The following work has been completed (but reports have not yet been circulated), or is in progress.

1. Molecular Structure Effects. A study by VEM of n-propyliodide has been completed. The purpose of the study was to extend the test of the relation between molecular structure and energy transfer efficiency. Propyl iodide is the first polar molecule to be studied in this series. The interesting result has been found that although the vibrational energy level density is greater than that in cyclopropane so that lesser efficiency would be expected, instead the energy transfer model is a stronger one and is attributed to increased attractive intermolecular potential. This will be described in TR21.
2. Nitroalkanes. We are in the middle of a study of the decomposition of nitromethane by the VEM technique. We are using $m = 5$ and 14. We have determined the average collisional reaction probabilities $P_c(m)$. The reaction may be influenced by surface, and this is taking time and effort to sort out. Also, with our comparatively mild sequential activation technique, we have been able to detect at least one higher molecular weight intermediate that has not hitherto been detected in a thermal system.
3. Isotopic Substitution. A study of the secondary molecular isotope effect on efficiency of vibrational energy transfer at high temperatures is nearing completion. The comparison of cyclo-C₃H₆ and cyclo-C₃D₆ provides a very refined test of the relationship between transfer efficiency and molecular structure.
4. Variation of Initial Vibrational Distribution. The study of vibrational energy transfer efficiency as a function of the initial vibrational population vector has not been studied previously for polyatomic molecules/thermal systems. We are completing a study of the isomerization of cyclobutene over the range of reaction temperatures from 600 K to 900 K and initial population temperatures from 273 K to 600 K.

V. Future Work.

The following studies are planned in the year ahead:

1. Nitroalkanes. The study of the vibrational collisional energy transfer efficiency with nitromethane will be continued. This molecule has a large dipole moment and appears to show high efficiency in our preliminary work. The biggest complication in this study may be catalytic effects, but this will, hopefully, not prevent us from reaching a successful analysis of the data. Past work between 800 K and 1000 K will be extended. Other nitroalkanes will also be considered.
2. Initial Energy Distribution. We will complete the present investigation of changing initial temperature (average vibrational energy) of the cyclobutene species on the energy transfer process. We will then extend the work to a molecule having a higher vibrational level density pattern such as methyl cyclobutene.
3. Surface Variation. We hope to examine the effect of variation of the nature of the wall surface and propose to investigate next a metal (gold) surface. We have hoped to initiate this work earlier but were held off by the other work in progress.
4. Variation of Molecule Parameters. We should like to move part of the work to the study of inorganic species such as NOCl , NO_2Cl , etc. as examples of plausible candidates. This would widen the scope of this aspect of the study considerably. Virtually no studies of the kind we propose have ever been made.

VI. Personnel

The following graduate students and fellows have been involved on this project in the period.

Dr. Ryuichi Arakawa
Prof. K. J. Chao (Sabbatical visitor, Tsing Hua University, Taiwan)
Dr. Toshio Kasai
Dr. David Kelley
Reiko Tosa
Wei Yuan (Sabbatical visitor, Qinghua University, Peking)

INT.

6/80

ENERGETIC MATERIALS RESEARCHDISTRIBUTION LIST

Aug. 7, 1981

	<u>No. Copies</u>		<u>No. Copies</u>
Assistant Secretary of the Navy (R, E, and S) Attn: Dr. R.E. Reichenbach Room 5E787 Pentagon Washington, DC 20350	1	AFATL Eglin AFB, FL 32542 Attn: Dr. Otto K. Heiney	1
Office of Naval Research Code 473 Arlington, VA 22217 Attn: Dr. R. Miller	10	AFRPL Code PACC Edwards AFB, CA 93523 Attn: Mr. W. C. Andrepont	1
Office of Naval Research Code 2008 Arlington, VA 22217 Attn: Dr. J. Enig	1	AFRPL Code CA Edwards AFB, CA 93523 Attn: Dr. R. R. Weiss	1
Office of Naval Research Code 260 Arlington, VA 22217 Attn: Mr. D. Siegel	1	Code AFRPL MKPA Edwards AFB, CA 93523 Attn: Mr. R. Geisler	1
Office of Naval Research Western Office 1030 East Green Street Pasadena, CA 91106 Attn: Dr. R. J. Marcus	1	Code AFRPL MKPA Edwards AFB, CA 93523 Attn: Dr. F. Roberto	1
Office of Naval Research Eastern Central Regional Office 495 Summer Street Boston, MA 02210 Attn: Dr. L. Peebles Dr. A. Wood	2	AFSC Andrews AFB, Code DLFP Washington, DC 20334 Attn: Mr. Richard Smith	1
Office of Naval Research San Francisco Area Office One Hallidie Plaza Suite 601 San Francisco, CA 94102 Attn: Dr. P. A. Miller	1	Air Force Office of Scientific Research Directorate of Chemical & Atmospheric Sciences Bolling Air Force Base Washington, DC 20332	1
Defense Technical Information Center DTIC-DDA-2 Cameron Station Alexandria, VA 22314	12	Air Force Office of Scientific Research Directorate of Aero- space Sciences Bolling Air Force Base Washington, DC 20332 Attn: Dr. L. H. Caveny	1
		Anal-Syn Lab Inc. P.O. Box 547 Paoli, PA 19301 Attn: Dr. V. J. Keenan	1

	<u>No. Copies</u>		<u>No. Copies</u>
Army Ballistic Research Labs Code DRDAR-BLP Aberdeen Proving Ground, MD 21005 Attn: Mr. L. A. Watermeier	1	Hercules Inc. Eglin AFATL/DL DL Eglin AFB, FL 32542 Attn: Dr. Ronald L. Simmons	1
Army Ballistic Research Labs ARRADCOM Code DRDAR-BLP Aberdeen Proving Ground, MD 21005 Attn: Dr. Ingo W. May	1	Hercules Inc. Magna Bacchus Works P.O. Box 98 Magna, UT 84044 Attn: Mr. E. H. DeButts	1
Army Ballistic Research Labs ARRADCOM Code DRDAR-BLT Aberdeen Proving Ground, MD 21005 Attn: Dr. Philip Howe	1	Hercules Inc. Magna Bacchus Works P.O. Box 98 Magna, UT 84044 Attn: Dr. James H. Thacher	1
Army Missile Command Code DRSME-RK Redstone Arsenal, AL 35809 Attn: Dr. R. G. Rhoades Dr. W. W. Wharton	2	HQ US Army Material Development Readiness Command Code DRCDE-DW 5011 Eisenhower Avenue Room 8N42 Alexandria, VA 22333 Attn: Mr. S. R. Matos	1
Atlantic Research Corp. 5390 Cherokee Avenue Alexandria, VA 22314 Attn: Dr. C. B. Henderson	1	Johns Hopkins University APL Chemical Propulsion Information Agency Johns Hopkins Road Laurel, MD 20810 Attn: Mr Theodore M. Gilliland	1
Ballistic Missile Defense Advanced Technology Center P.O. Box 1500 Huntsville, AL 35807 Attn: Dr. David C. Sayles	1	Lawrence Livermore Laboratory University of California Livermore, CA 94550 Attn: Dr. M. Finger	1
Ballistic Research Laboratory USA ARRADCOM DRDAR-BLP Aberdeen Proving Ground, MD 21005 Attn: Dr. A. W. Barrows	1	Lawrence Livermore Laboratory University of California Livermore, CA 94550 Attn: Dr. R. McGuire	1
Hercules Inc. Cumberland Aerospace Division Allegany Ballistics Lab P.O. Box 210 Cumberland, MD 21502 Attn: Dr. Rocco Musso	2	Lockheed Missiles and Space Co. P.O. Box 504 Sunnyvale, CA 94088 Attn: Dr. Jack Linsk Org. 83-10 Bldg. 154	1

	<u>No. Copies</u>		<u>No. Copies</u>
Lockheed Missile & Space Co. 3251 Hanover Street Palo Alto, CA 94304 Attn: Dr. H. P. Marshall Dept. 52-35	1	Naval Research Lab Code 6100 Washington, DC 20375	1
Los Alamos Scientific Lab P.O. Box 1663 Los Alamos, NM 87545 Attn: Dr. R. Rogers, WX-2	1	Naval Sea Systems Command Washington, DC 20362 Attn: Mr. G. Edwards, Code 62R3 Mr. J. Murrin, Code 62R2 Mr. W. Blaine, Code 62R	1
Los Alamos Scientific Lab P.O. Box 1663 Los Alamos, NM 87545 Attn: Dr. B. Craig, M Division	1	Naval Sea Systems Command Washington, DC 20362 Attn: Mr. R. Beauregard SEA 64E	1
Naval Air Systems Command Code 330 Washington, DC 20360 Attn: Mr. R. Heitkotter Mr. R. Brown	1	Naval Surface Weapons Center Code R11 White Oak, Silver Spring, MD 20910 Attn: Dr. H. G. Adolph	1
Naval Air Systems Command Code 310 Washington, DC 20360 Attn: Dr. H. Mueller Dr. H. Rosenwasser	1	Naval Surface Weapons Center Code R13 White Oak, Silver Spring, MD 20910 Attn: Dr. R. Bernecker	1
Naval Explosive Ordnance Disposal Facility Indian Head, MD 20640 Attn: Lionel Dickinson Code D	1	Naval Surface Weapons Center Code R10 White Oak, Silver Spring, MD 20910 Attn: Dr. S. J. Jacobs	1
Naval Ordnance Station Code 5034 Indian Head, MD 20640 Attn: Mr. S. Mitchell	1	Naval Surface Weapons Center Code R11 White Oak, Silver Spring, MD 20910 Attn: Dr. M. J. Kamlet	1
Naval Ordnance Station Code PM4 Indian Head, MD 20640 Attn: Mr. C. L. Adams	1	Naval Surface Weapons Center Code R04 White Oak, Silver Spring, MD 20910 Attn: Dr. D. J. Pastine	1
Dean of Research Naval Postgraduate School Monterey, CA 93940 Attn: Dr. William Tolles	1	Naval Surface Weapons Center Code R13 White Oak, Silver Spring, MD 20910 Attn: Dr. E. Zimet	1
Naval Research Lab Code 6510 Washington, DC 20375 Attn: Dr. J. Schnur	1		

6/80

4

	<u>No. Copies</u>		<u>No. Copies</u>
Naval Surface Weapons Center Code R101 Indian Head, MD 20640 Attn: Mr. G. L. MacKenzie	1	Naval Weapons Center Code 388 China Lake, CA 93555 Attn: D. R. Derr	1
Naval Surface Weapons Center Code R17 Indian Head, MD 20640 Attn: Dr. H. Haiss	1	Naval Weapons Center Code 388 China Lake, CA 93555 Attn: Dr. R. Reed Jr.	1
Naval Surface Weapons Center Code R11 White Oak, Silver Spring, MD 20910 Attn: Dr. K. F. Mueller	1	Naval Weapons Center Code 385 China Lake, CA 93555 Attn: Dr. A. Nielsen	1
Naval Surface Weapons Center Code R16 Indian Head, MD 20640 Attn: Dr. T. D. Austin	1	Naval Weapons Center Code 3858 China Lake, CA 93555 Attn: Mr. E. Martin	1
Naval Surface Weapons Center Code R122 White Oak, Silver Spring, MD 20910 Attn: Mr. L. Roslund	1	Naval Weapons Center China Lake, CA 93555 Attn: Mr. R. McCarten	1
Naval Surface Weapons Center Code R121 White Oak, Silver Spring, MD 20910 Attn: Mr. M. Stosz	1	Naval Weapons Support Center Code 5042 Crane, Indiana 47522 Attn: Dr. B. Douda	1
Naval Weapons Center Code 3853 China Lake, CA 93555 Attn: Dr. R. Atkins	1	Rohm and Haas Company 723-A Arcadia Circle Huntsville, Alabama 35801 Attn: Dr. H. Shuey	1
Naval Weapons Center Code 3205 China Lake, CA 93555 Attn: Dr. L. Smith	1	Strategic Systems Project Office Dept. of the Navy Room 901 Washington, DC 20376 Attn: Dr. J. F. Kincaid	1
Naval Weapons Center Code 3205 China Lake, CA 93555 Attn: Dr. C. Thelen	1	Strategic Systems Project Office Dept. of the Navy Room 1048 Washington, DC 20376 Attn: Mr. E. L. Throckmorton Mr. R. Kinert	2
Naval Weapons Center Code 385 China Lake, CA 93555 Attn: Dr. A. Amster	1	Thiokol Chemical Corp. Brigham City Wasatch Division Brigham City, UT 84302 Attn: Dr. G. Thompson	1

6/80

5

	<u>No. Copies</u>		<u>No. Copies</u>
USA ARRADCOM DRDAR-LCE Dover, NJ 07801 Attn: Dr. R. F. Walker	1	University of California Department of Chemistry 405 Hilgard Avenue Los Angeles, CA 90024 Attn: Professor M. Nicol	1
USA ARRADCOM DRDAR-LCE Dover, NJ 07801 Attn: Dr. N. Slagg	1	University of California Energy Center Mail Code B-010 La Jolla, CA 92093 Attn: Prof. S.S. Penner	1
U.S. Army Research Office Chemistry Division P.O. Box 12211 Research Triangle Park, NC 27709	1		
Washington State University Dept. of Physics Pullman, WA 99163 Attn: Professor G.D. Duval	1	Dr. P. Rentzepis Bell Laboratories Murray Hill N.J. 07971	1
Space Sciences, Inc. 135 West Maple Avenue Monrovia, CA 91016 Attn: Dr. M. Farber	1	University of Southern CA Department of Electrical Engineering University Park Los Angeles, CA 90007 Attn: C. Wittig	1
SRI International 333 Ravenswood Avenue Menlo Park, CA 94025 Attn: Mr. M. Hill	1	MIT Dept. of Chemistry Cambridge, MA 02139 Attn: Prof. John Deutsch	1
Office of Naval Research Code 421 Arlington, VA 22217 Attn: Dr. B. Junker	1		
The Johns Hopkins University Department of Chemistry Baltimore, MD 21218 Attn: Dr. Joyce J. Kaufman	1		
University of California Department of Chemistry Berkeley, CA 94720 Attn: Professor Y.T. Lee	1		
Office of Naval Research Code 472 Arlington, VA 22217 Attn: Dr. G. Neece	1		