

AD-A103 816 DUKE UNIV MEDICAL CENTER DURHAM NC DEPT OF ANESTHESIOLOGY F/G 6/19  
PULMONARY EXTRAVASCULAR FLUID DYNAMICS IN OXYGEN TOXICITY.(U)  
1980 J N MILLER; E M CAMPORESI; R COGGIN N00014-76-C-0923

UNCLASSIFIED

NL

1 of 1  
AD-A  
10-816



END  
DATE  
FILMED  
10-81  
DTIC



# DUKE UNIVERSITY MEDICAL CENTER

Department of Anesthesiology

**DISTRIBUTION STATEMENT A**  
Approved for public release;  
Distribution Unlimited

**LEVEL II**

**12**

**DTIC ELECTE**

SEP 8 1981

**S D D**

AD A103816

Office of Naval Research Contract Number: <sup>15</sup> N00014-76-C-0923 <sup>1400</sup>

<sup>6</sup> Title: Pulmonary Extravascular Fluid Dynamics in Oxygen Toxicity.

Principal Investigator: John N. Miller, M.D.

Co-Investigator: <sup>12</sup> Enrico M. Camporesi, M.D.

Technical Assistance: Robin Coggin, B.S.  
Melissa McLeod, M.S.

<sup>7</sup> FINAL TECHNICAL REPORT <sup>12 3</sup>

Introduction. Pulmonary oxygen toxicity in man is heralded by the onset of symptoms primarily referred to the upper airway (retrosternal discomfort, feelings of chest tightness and coughing on deep inspiration). Much has been made of the dose/response importance of quantified decrements in vital capacity as evidencing parenchymal injury (1), yet such decrements also can be interpreted in terms of airway irritation. Early indications of pulmonary parenchymal injury generally involve manifestations of increased lung water content (2). We have made comparative studies of lung water content ( $V_{tiss}$ ), pulmonary capillary blood flow ( $Q_C$ ) and pulmonary diffusing capacity ( $D_{LCO}$ ) with vital capacity ( $V_C$ ) and onset of symptoms during the development of early pulmonary oxygen toxicity in man.

Methods. Fourteen male volunteer subjects aged between 21 and 38 were included in the study. Four subjects were heavy smokers. Each subject underwent a series of measurements performed during air breathing, 100% oxygen breathing over six hours at normal atmospheric pressure (1 ATA); six hours breathing humidified 100%  $O_2$  at 2 ATA; six hours breathing 100%  $O_2$  dry at 2 ATA; and six hours breathing 100% humidified  $O_2$  at 2.25 ATA. In addition, each subject was studied at 24, 48 and 72 hours after each 2 and 2.25 ATA exposure. Studies on individual subjects were spaced so that no effects of any exposure would be cumulative with respect to other exposures. The protocol was approved by the Clinical Investigation Committee and informed consent was obtained. Duplicate measurements of  $V_{tiss}$ ,  $Q_C$  and  $D_{LCO}$  were made using a rebreathing technique with simultaneous gas analysis by quadrupole mass spectrometry (2,3). Vital capacities were obtained by spirometry, using in each case, the best of three measurements. Each battery of studies was repeated at 30 minute intervals throughout each exposure, and at 24, 48 and 72 hours following hyperbaric oxygen exposures.

DATE FILE COPY

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By: Per Ltr. on File	
Distribution/	
Availability Codes	
Dist	Special
<b>A</b>	

Box 3094 • Durham, North Carolina 27710 • Telephone (919) 684-3708

81 8 21 069  
412 501

2000

**Results.** No symptoms of pulmonary oxygen toxicity were reported during control exposures, but began to develop from the fourth hour of exposure at 2.0 and 2.25 ATA. In each case, symptoms developed first in the four smokers, being the only subjects to report respiratory discomfort during the 2.0 ATA humidified O<sub>2</sub> exposure. Twelve subjects, including the smokers, developed symptoms during the dry O<sub>2</sub> exposure at 2 ATA, again from the fourth hour on. Six subjects developed symptoms during their 2.25 ATA humidified O<sub>2</sub> exposures. The severity of symptoms progressively increased with the exposure.

Despite careful measurements of  $V_C$ , we found no consistent decrement caused by the O<sub>2</sub> exposures. Variability of measurement in each subject was  $\pm 2\%$ , so that we felt confident of detecting a 5% decrement as predicted from previous work (1).

Only in the final hour of the dry O<sub>2</sub> exposure at 2.0 ATA and that of the humidified O<sub>2</sub> exposure at 2.25 ATA was any increase in  $V_{tiss}$  detected. This was seen in the same 4 subjects, 2 of whom were smokers, and was of the order of a 50% increase ( $673 \pm 248$  to  $960 \pm 328$  ml). No such increase was seen in the remaining subjects. Moreover, in those subjects whose  $V_{tiss}$  increased, the increase persisted for at least 24 hours following the exposures, but returned to control values by 48 and 72 hours.

Pulmonary diffusing capacity was diminished in all subjects throughout each hyperbaric oxygen exposure ( $28 \pm 4$  ml/min/mm Hg down to  $19.2 \pm 1.5$ ). This was primarily caused by reduced CO uptake by hemoglobin because of the increased number of competing O<sub>2</sub> molecules. However, in those subjects who developed an increase in  $V_{tiss}$  in the final hour of the 2 ATA dry O<sub>2</sub> exposure and the humidified O<sub>2</sub> exposure at 2.25 ATA,  $D_{LCO}$  was further decreased to  $16 \pm 1.2$  ml/min/mm Hg. Again, in the same subjects, a reduction in  $D_{LCO}$  when compared with control values was seen during the 24 hour post exposure studies, but not at 48 or 72 hours.

Measurements of  $Q_C$  showed no consistent differences with respect to controls during or after any of the hyperbaric exposures. Values ranged between  $5.6 \pm 1.0$  and  $8.9 \pm 0.8$  L/min.

**Discussion.** The pathophysiology of early pulmonary oxygen toxicity centers around injury to the Type II pneumocyte with a progressive disruption of surfactant structure, and a concomitant increase in intracellular and interstitial water content. We reasoned that the measurement of  $V_{tiss}$  and  $D_{LCO}$  during the onset of early pulmonary O<sub>2</sub> toxicity would provide a more insightful, yet non-invasive, indicator of the underlying pathophysiology than a decrement in  $V_C$ , since this would be caused by acute airway irritation, as by a reduction in pulmonary compliance, our studies were originally designed to determine the contribution of an early accumulation of lung water to the observed decrement in  $V_C$ . Because of previous authoritative reports of a dose/response relationship between hyperbaric oxygen exposure and diminution in vital capacity (1), we were surprised to find no such relationship in our studies.

*V<sub>tiss</sub> data*

The rebreathing technique for the determination of  $V_{tiss}$  is fraught with great variability, so that quite large increases must be present to be reliably detected. Inherent in the method is the requirement for complete gas mixing within the lung within several rebreathing breaths. Further, the technical problems associated with the technique are compounded by the hyperbaric  $O_2$  environment. Hence, a consistent increase in  $V_{tiss}$  in the same subjects over at least 4 separate measurements, persisting at normal atmospheric pressure over at least 2 further measurements 24 hours, and subsequently returning to control values in the following 4 to 6 measurements, indicates that water does indeed accumulate in the lungs of normal man during the earliest stages of pulmonary  $O_2$  toxicity. Symptoms develop well in advance of objectively measurable and reversible pathophysiology, therefore represent the earliest reliable indicator of impending pulmonary  $O_2$  toxicity.

#### References.

1. Clark, JM and CJ Lambertsen, J. Applied Physiol. 30:739, 1971.
2. Bachofen, M and ER Weibel, Chest, 65:145, 1974.
3. Sackner, MA, D. Greeneltch, MS Heiman, S Epstein and N Atkins, Am. Rev. Respir. Dis. 111(2):157, 1875.

#### Publications and Papers Presented:

1. Miller, J.N. and Camporesi, E.M.: Rebreathing of  $D_{LCO}$  with  $^{13}CO$  Feder. Proc. 36, No. 3, Abstract No. 1888, 1977.
2. Miller, J.N., Camporesi, E.M. and Fagraeus, L.: Rebreathing Assessment of  $V_{tiss}$ ,  $V_C$ ,  $Q_C$  and  $D_{LCO}$  with  $N_2O$  and  $^{13}CO$ . Proc. Internat. Union Physiol. Sci. 13:1517, 1977.
3. Miller, J.N., Miller, Chris and Coggin, Robin: Pulmonary tissue volume and vital capacity in early pulmonary oxygen toxicity (Abstract). Undersea Med. Soc., Undersea Biomed. Res., sup. Vol. 5, No. 1, Abstract No. 9, p. 16, March, 1978.
4. Miller, J.N.: Pulmonary Tissue Volume, Blood Flow, Diffusing Capacity and Vital Capacity in Early Pulmonary Oxygen Toxicity in Man. Proc. Am. Soc. Anesthesiologists, Ann. Mtg. (1978):61.
5. Miller, J.N.: Pulmonary Oxygen Toxicity. Proceedings of Annual Meeting of South Pacific Underwater Medicine Society, Singapore, 1980 (In Press).
6. Miller, J.N., Camporesi, E.M., McLeod, M. and M. Feezor: Studies of Early Pulmonary Oxygen Toxicity in Man. Accepted for presentation at the 1981 Annual Meeting of ASA, New Orleans, La.

The research reported here has been supported under the Office of Naval Research contract N00014-76-C-0923 with funds provided by the Naval Medical Research and Development Command.

**DAT  
FILM  
0-8**