

AD-A104 262 MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF NUTRITI--ETC F/8 6/13
FUNCTION OF THE PEPTIDE ANTIBIOTIC, GRAMICIDIN S, IN ITS PRODUCE--ETC(U)
SEP 81 J M PIRET DAA629-78-C-0015
NL

UNCLASSIFIED

ARO-14926.2-L

For
AD-A104 262



END
DATE
FILMED
40-811
DTIC

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

LEVEL II

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER 14926.2-L	2. GOVT ACCESSION NO. AD-A104262	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Function of the Peptide Antibiotic, Gramicidin S, in its Producer, <u>Bacillus Brevis</u> Nagano		5. TYPE OF REPORT & PERIOD COVERED Final Report: 15 Mar 78 - 14 Jun 81
7. AUTHOR(s) Jacqueline Marie Piret		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Massachusetts Institute of Technology Cambridge, MA 02139		8. CONTRACT OR GRANT NUMBER(s) DAAG29 78 C 0015
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE Sep 81
		13. NUMBER OF PAGES 6
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) antibiotics mutants ultraviolet radiation peptides microbiology solvents sporulation germination spores heat		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The natural function(s) of the peptide antibiotic, gramicidin S (GS) in its producer, <u>Bacillus brevis</u> Nagano, was investigated. Particular attention was paid to the possible role of GS in the differentiation process: sporulation, spore properties and germination. The GS-producing parental strain and a GS-negative mutant of this strain were compared.		

AD A104262

DIN FILE COPY

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

81 3

10 000

Accession For	
NTIS GRA&I	☒
DTIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Availability Codes	
Dist	
Special	
A	

DAAG29-78-C-0015

FUNCTION OF THE PEPTIDE ANTIBIOTIC, GRAMICIDIN S,
IN ITS PRODUCER, BACILLUS BREVIS NAGANO

by

JACQUELINE MARIE/PIRET

Submitted to the Department of Nutrition and Food Science
on March 17, 1981 in partial fulfillment of the
requirements for the Degree of Doctor of Philosophy in
Nutritional Biochemistry and Metabolism

ABSTRACT

The natural function(s) of the peptide antibiotic, gramicidin S (GS) in its producer, Bacillus brevis Nagano, was investigated. Particular attention was paid to the possible role of GS in the differentiation process: sporulation, spore properties and germination. The GS-producing parental strain and a GS-negative mutant of this strain were compared.

Sporulation rate and efficiency, sporulation-associated events (except for GS production) were similar in both strains. Mature parental and mutant spores were equally resistant to heat, ultraviolet irradiation and solvents. Germination initiation in both strains was rapid and both responded similarly to a spectrum of germinants tested. Thus the lack of GS synthesis impaired none of these properties.

Outgrowth, however, was affected. In the presence of exogenous GS, spore outgrowth was inhibited. Whereas mutant spores proceeded through outgrowth and entered vegetative growth quickly, parental spores remained in outgrowth several hours longer. Extraction of GS from the parental spores reduced the outgrowth delay. GS apparently acted from the outside on spores.

Outgrowth delay was dependent upon the concentration of GS present. There was a distinct time during outgrowth when mutant spores were no longer sensitive to exogenous GS. Outgrowth inhibition by GS was permanent. Alanine uptake experiments suggested that GS affects the entry of nutrients into the outgrowing spore.

Thesis Advisor: Dr. Arnold L. Demain

Title: Professor of Industrial Microbiology

Approved for publication
Distribution Unlimited

VI. SUMMARY AND CONCLUSIONS

The role of the peptide antibiotic gramicidin S in its producer Bacillus brevis Nagano was studied. The parent strain and its GS-negative mutant, BI-7, were compared. The study focused on the postulated function of GS in differentiation: sporulation, spore properties and germination. Spores were produced in media supporting low, moderate and high GS synthesis. The extent of sporulation was inversely related to antibiotic production.

Growth and sporulation in parent and BI-7 were followed. The rate, extent and sequence of events of sporulation were similar in both strains in all media used.

The properties of mature spores of the parent and BI-7 were also compared. UV, solvent and heat resistances were determined for spores produced in a range of media. DPA contents were also measured. The absence of GS synthesis by the mutant did not impair the "quality" of its spores.

The response of spores to heat activation and to

germinants was measured by loss of absorbance of spore suspensions. The GS-negative mutant and its parent behaved similarly, regardless of the medium from which they were harvested.

Spore outgrowth, however, was affected by GS. Endogenous GS, associated with parental spores or exogenous GS, added to mutant spores, extended outgrowth by several hours. In the absence of GS or when GS was removed from spores, outgrowth was rapid. All or most of the GS was at or near the spore surface. Outgrowth inhibition by GS was probably permanent. At a distinct time during outgrowth, GS no longer exerted its effect. The uptake of labelled L-alanine by outgrowing spores was inhibited by GS. This inhibition was at a step earlier than the incorporation of L-alanine into protein.

Gramicidin S, made during sporulation by Bacillus brevis Nagano, remains associated with the sporulating, dormant and initiating organism but apparently without affecting these stages of development. During spore outgrowth, however, the antibiotic prolongs this process, inhibiting the onset of vegetative growth. In this GS-sensitive period, nutrient uptake is impaired by GS. The data suggest a mechanism of action for GS whereby it

DAAG29-78-C-0015

interferes in membrane function, such as transport or
energy metabolism, in outgrowing spores.

PUBLICATIONS

- Demain, A.L. and J.M. Piret. 1977. Speculations on the role(s) of antibiotics in spore-forming microorganisms. In Spores VII. G. Chambliss and J.C. Vary, eds. Amer. Soc. Microbiol. Washington, D.C. p. 17.
- Demain, A.L. and J.M. Piret. 1978. Relationship between antibiotic biosynthesis and sporulation. In Regulation of Secondary Product and Plant Hormone Metabolism. M. Luckner and K. Schreiber, eds. Pergamon Press. New York. p. 183.
- Piret, J.M. and A.L. Demain. 1981. Function of the peptide antibiotic gramicidin S in the producer organism, Bacillus brevis. In Sporulation and Germination. Proc. 8th Intl. Spores Conf. Amer. Soc. Microbiol. (in press)
- Piret, J.M., J. Millet and A.L. Demain. Intracellular protease in Bacillus brevis ATCC 9999. (in preparation)
- Piret, J.M. and A.L. Demain. Function of gramicidin S in its producer, Bacillus brevis Nagano. (in preparation)

PERSONNEL

Professor Arnold L. Demain
Professor of Industrial Microbiology
Massachusetts Institute of Technology
Cambridge, Mass. 02139

Dr. Jacqueline M. Piret
Laboratory of Industrial Microbiology
Massachusetts Institute of Technology
Cambridge, Mass. 02139

Dr. Piret obtained her Ph.D. on this project in June, 1981.