

AD-A105 974

NAVAL SURFACE WEAPONS CENTER DAHLGREN VA

F/6 8/3

THE NSWC GLOBAL OCEAN TIDE DATA TAPE (GOTD), ITS FEATURES AND A--ETC(U)

JUN 81 E W SCHWIDERSKI, L T SZETO

UNCLASSIFIED

NSWC/TR-81-254

NL

1 OF 1
AD A
105 974

END
DATE
FILMED
11-81
DTIC

AD A105974

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 NSWC/TR-81-254	2. GOVT ACCESSION NO. 41-4102	3. RECIPIENT'S CATALOG NUMBER 974
4. TITLE (and Subtitle) 6 THE NSWC GLOBAL OCEAN TIDE DATA TAPE (GOTD), ITS FEATURES AND APPLICATION, RANDOM-POINT TIDE PROGRAM	5. TYPE OF REPORT & PERIOD COVERED Final Repts.	
7. AUTHOR(s) 10 E. W. Schwiderski L. T. Szeto	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Surface Weapons Center (K104) Dahlgren, Virginia 22448	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 17 61152N ZR000-01-01 16 ZR00004	
11. CONTROLLING OFFICE NAME AND ADDRESS Chief of Naval Material Department of the Navy Washington, DC 20360	12. REPORT DATE June 1981	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 43 12 42	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES S SELECTED SEP 23 1981 A		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ocean Tides and Currents Numerical Modeling Tidal Charts		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In the following report, the authors highlight briefly the major qualitative and quantitative features of the NSWC ocean tide models, which now include the nine components (M_2 , S_2 , N_2 , K_2), (K_1 , O_1 , P_1 , Q_1), and M_f with models for M_m and S_{sa} nearing completion. Special characteristics of those models are pointed out, which require indispensable attention in various applications particularly in coastal areas with rapidly varying tides. Since the tide models interpolate (see back)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered) **424567**

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

(20)

hydrodynamically empirical tide data at over 2000 continental and island stations, certain simple corrections in the numerical procedures are presented. Plans for more extensive local and perhaps global refinements and improvements of the tide models are mentioned.

In the final part of this report, the authors describe the NSW Global Ocean Tide Data (GOTD) tape including the data arrangement and the standard format. The report ends with a program that computes from the taped harmonic tidal constants instantaneous oceanic and geocentric tides at specified random points and instances in the Random-Point Tide (RPTIDE) program.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

FOREWORD

In this report the authors elaborate the major features of the NSWC ocean tide models, which are significant in various applications. Simple improvements in coastal waters are suggested. The harmonic tidal constants are arranged on the NSWC Global Ocean Tide Data (GOTD) tape, which is described for possible users. The report concludes with the description of the NSWC Random-Point Tide (RPTIDE) program that computes tidal elevations at given random points and instances.

This project was supported by the Naval Surface Weapons Center's Independent Research Fund and by a grant from the National Geodetic Survey of the Department of Commerce/NOS/NOAA.* It is the author's most pleasant obligation to acknowledge the sustained and generous sponsorship of Mr. R. T. Ryland, Jr., Head of the Strategic Systems Department, his Associate, Dr. R. J. Anderle, and Mr. D. R. Brown, Jr., Head of the Space and Surface Systems Division. Many critical and stimulating suggestions were gratefully received from the author's colleagues, Drs. C. J. Cohen, C. Oesterwinter, and B. Zondek.

The date of completion was June 20, 1981.

Released by

David R. Brown Jr.

DAVID R. BROWN, JR., Acting Head
Strategic Systems Department

*National Ocean Survey (NOS)
National Oceanographic and Atmospheric Administration (NOAA)

Accession For	
NSWC GRA&I	☒
ERIC TAB	☐
Unannounced	☐
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

CONTENTS

	Page
FOREWORD.....	iii
ABSTRACT	vii
1. INTRODUCTION.....	1
2. MAJOR FEATURES OF THE NSWC OCEAN TIDE MODELS	3
3. APPLICATION OF THE NSWC OCEAN TIDE DATA	6
4. NSWC GLOBAL OCEAN TIDE DATA (GOTD) TAPE.....	9
5. NSWC RANDOM-POINT TIDE PROGRAM (RPTIDE)	12
REFERENCES	16
APPENDIXES	
A. RPTIDE PROGRAM USER'S GUIDE	
B. RPTIDE PROGRAM LISTING	
DISTRIBUTION	

ABSTRACT

In the following report, the authors highlight briefly the major qualitative and quantitative features of the NSWC ocean tide models, which now include the nine components (M_2 , S_2 , N_2 , K_2), (K_1 , O_1 , P_1 , Q_1), and M_f with models for M_m and S_{sa} nearing completion. Special characteristics of those models are pointed out, which require indispensable attention in various applications particularly in coastal areas with rapidly varying tides. Since the tide models interpolate hydrodynamically empirical tide data at over 2000 continental and island stations, certain simple corrections in the numerical procedures are presented. Plans for more extensive local and perhaps global refinements and improvements of the tide models are mentioned.

In the final part of this report, the authors describe the NSWC Global Ocean Tide Data (GOTD) tape including the data arrangement and the standard format. The report ends with a program that computes from the taped harmonic tidal constants instantaneous oceanic and geocentric tides at specified random points and instances in the Random-Point Tide (RPTIDE) program.

1. INTRODUCTION

Although the mathematical modeling of ocean tides enjoyed since Newton a tricentennial fascinating history (Schwiderski 1980a), due to the measured digital knowledge of the bathymetry of the real ocean basins, the solution of the problem was destined to await the advent of modern digital computers. Fortunately, along with the necessary computational tool, significantly improved numerical techniques and hydrodynamical notions of turbulent dissipation processes became available for the mathematically discrete and realistic definition of the ocean model (Schwiderski 1980a). Nevertheless, the strictly mathematical formulation of the problem failed to describe the true tides over some important ocean areas where low tides were predicted for observed high tides. The computed tidal charts conspicuously displayed none of the long-known strong distortions and retardations of ocean tides by shallow continental shelves (see, e.g., Figure 2) and narrow ocean ridges.

To overcome those fundamental shortcomings, the author (Schwiderski 1978a, b, 1979a - d, and 1980a) introduced a unique hydrodynamical interpolation technique, which incorporates into the model, empirical tide data collected from harmonically analyzed tide gauge measurements along shorelines and at some other ocean bottom irregularities. The combined mathematical-experimental method removed essentially all purely mathematical deficiencies and made the desired accurate charting of ocean tides possible (Schwiderski 1979e, 1980b, 1981a - h, j, and 1982a - d).

All earlier investigations started from the basic Laplace tidal equations (LTEs), which included only the astronomical equilibrium tide as its sole tide-forcing potential. Following suggestions of Proudman (1928) that there might exist significant interactions between the oceanic and terrestrial tides, Grace (1930) augmented the LTEs by including simple approximations of the terrestrial tide, the terrestrial response to the oceanic tidal load, and the corresponding three gravity perturbations of the equilibrium tide.

Remarkably, all numerical tidalists disregarded those investigations till Farrell (1972) derived a more accurate approximation of the same interaction terms, which were then considered by Hendershott (1972), Zahel (1977), Estes (1979), Parke and Hendershott (1979) and others. In order to avoid the involved computation of the loading effects, Accad and Pekeris (1978) assumed a uniform proportionality between the oceanic tide and its loading effects. Though this type of simplification was earlier suggested by Takahasi (1929) and Grace (1930), Accad and Pekeris justified this approach by displaying the striking resemblance of the corresponding global maps. The author (Schwiderski 1978a, 1979c, d, and 1980a) adopted the same approximation for all NSWC ocean tide models, and Goad (1980b) computed for the M_2 ocean tide the corresponding terrestrial dip, which also resembles closely its cause.

Since the first publication of the author's NSWC ocean tide model about three years ago, an unexpected wide range of applications to various problems in oceanography, geophysics, astronomy, and space technology has been conducted. Fortunately, most of those applications

were very gratifying as they confirmed pointwise and/or globally the high accuracy and usefulness of the computed partial tides (see, e.g., Goad 1980a, b, Melchior et al. 1980, and Wahr and Sasao 1980). Since numerical global tide models are naturally limited in detailed accuracy in certain local ocean areas, some foreseeable difficulties emerged particularly in applications to compute loading effects of ocean tides on the earth gravity field.

In order to avoid unnecessary complications in such applications, relevant qualifying and limiting features of the NSWC ocean tide models will be pointed out in Section 2. Simple immediate and future improvements of the models will be mentioned and discussed in Section 3. The arrangement of the gridded ocean tide data on the NSWC magnetic tape is described in Section 4. Finally Section 5 describes a program to compute from the taped harmonic ocean tide constants instantaneous ocean and geocentric tidal heights at a given set of random geographical points and specified instances. The corresponding user's guide and program listing are presented in Appendixes A and B.

2. MAJOR FEATURES OF THE NSWC OCEAN TIDE MODELS

In applications, the following special characteristics of the NSWC ocean tide models require scrupulous attention and evaluation. All these features are directly discernible from the author's various publications listed in the References.

(a) **Modes of the NSWC Ocean Tide Models:** The NSWC ocean tide model includes now (see Table 1) nine completed harmonic partial tides of the semidiurnal (M_2 , S_2 , N_2 , K_2), diurnal (K_1 , O_1 , P_1 , Q_1) and long-period (Mf) species with models of the additional long-period components (Mm, Ssa) nearing completion. All partial tides are represented by their harmonic constants, that is, amplitudes and phases which are recorded gridwise on magnetic tape (Schwiderski 1980d, and 1981j). Also, gridwise tabulated data charts and corange and cotidal maps of all eleven constituents are presented in NSWC technical reports (Schwiderski 1979e, 1981a - g, and 1982a - c).

(b) **The Gridded Ocean Basins:** For the recording of the harmonic tidal constants, the simply connected global ocean basins are divided by integral-degree longitudes and colatitudes into uniform $1^\circ \times 1^\circ$ grid cells. Naturally, the model excludes all inland waters such as the Great Lakes and the Black and Dead Seas, as well as the following gridwise disconnected bordering waters: the Baltic, Kattegat, Irish, Mediterranean, Red, Ceram, Sulu, and Japan Seas; the Gulfs of California, Persia, Chihli, and Huraki; and the Hudson and Korean Bays. Similarly, many shallow (less than 5-m depth) and/or narrow waters such as the entire Barrier Reef area, the Gironde Estuary, and the Fjords of Norway could not be modeled.

(c) **Computed and Empirical Tide Data:** The recorded harmonic constants include all computed and hydrodynamically interpolated empirical tide data. The latter values are specially labeled in the printed charts (Schwiderski 1979e, 1981a - g and 1982a - c) for accuracy verification and for proper applications (see Section 3). All data are tabulated as representative for the centers of the corresponding grid cells; which is strictly realistic only for all open ocean areas where the tides vary negligibly over the mesh area. In fact, the computed data are rigorously defined as cell averages, which may deviate somewhat from the true point values where the tides suffer strong distortions and/or retardations over short distances.

The empirical data are by definition (Schwiderski 1978a, 1979d, and 1980a) aligned boundary values at continental or island shore stations or point values at some bottom irregularities. In addition to their obvious displacement (see Figure 1 below), many of those data must be viewed as crude averages of drastically varying high tides. This applies, for instance, to the Skagerak; the English Channel; the Bays of Fundy, Baffin, and Bristol; the Gulfs of St. Lawrence, Aden, Oman, Thailand, Tonkin, and Carpentaria; the Straits of Florida, Hudson, Formosa, Bass, and Cook; as well as most Seas of the Indonesian Archipelago, where mostly empirical input data are listed.

Table 1. Constants of Major Tidal Modes

Tidal Mode	K (m)	σ (10^{-4} /sec)	χ (deg)
Semidiurnal Species			
M_2 = Principal Lunar	0.242 334	1.405 19	$2h_0 - 2s_0$
S_2 = Principal Solar	0.112 841	1.454 44	0
N_2 = Elliptical Lunar	0.046 398	1.378 80	$2h_0 - 3s_0 + p_0$
K_2 = Declination Luni-Solar	0.030 704	1.458 42	$2h_0$
Diurnal Species			
K_1 = Declination Luni-Solar	0.141 565	0.729 21	$h_0 + 90$
O_1 = Principal Lunar	0.100 574	0.675 98	$h_0 - 2s_0 - 90$
P_1 = Principal Solar	0.046 843	0.725 23	$-h_0 - 90$
Q_1 = Elliptical Lunar	0.019 256	0.649 59	$h_0 - 3s_0 - 90$
Long-Period Species			
M_f = Fortnightly Lunar	0.041 742	0.053 234	$2s_0$
M_m = Monthly Lunar	0.022 026	0.026 392	$s_0 - p_0$
S_{sa} = Semiannual Solar	0.019 446	0.003 8921	$2h_0$

K = amplitude of the partial tide

σ = frequency of the partial tide

χ = astronomical argument of the partial tide.

(h_0, s_0, p_0) = mean longitudes of sun, moon, and lunar perigee at Greenwich midnight

$h_0 = 279.696\ 68 + 36\ 000.768\ 930\ 485T + 3.03 \cdot 10^{-4}T^2$

$s_0 = 270.434\ 358 + 481\ 267.883\ 141\ 37T - 0.001\ 133T^2 + 1.9 \cdot 10^{-6}T^3$

$p_0 = 334.329\ 653 + 4\ 069.034\ 032\ 957\ 5T - 0.010\ 325T^2 - 1.2 \cdot 10^{-5}T^3$

where

$T = [27\ 392.500\ 528 + 1.000\ 000\ 035\ 6D]/36\ 525$

$D = d + 365(y - 1975) + \text{Int}[(y - 1973)/4]$

d = day number of year ($d = 1$ for January 1)

$y \geq 1975$ = year number

and

$\text{Int}[x]$ = integral part of x

(d) **Accuracy in Open Oceans:** In the publications (Schwiderski (1978a, b, 1979a - c, 1980a, 1981a - h, and 1982a - d) a tide prediction accuracy of better than 5 cm anywhere in the open oceans has been claimed for the dominant M_2 tide. This accepted estimate is based on the worldwide agreement of the computed tide data with empirical data which carry the same accuracy. Indeed, due to the somewhat controversial harmonic analysis of tide gauge measurements, no higher accuracy can be claimed even by the most recent deep-sea tide data (see, e.g., Schwiderski 1981b). All lesser computed partial tides (see Table 1) carry relatively the same component accuracy as M_2 . Thus, in superposition a total tide prediction accuracy of better than 10 cm can now be assumed over all open oceans.

(e) **Accuracy in Coastal Waters:** Due to the a priori resolution limits of the $1^\circ \times 1^\circ$ grid system mentioned under (b) and (c) above, the NSWC tide models may lose some of their open ocean accuracy (d) in coastal waters of rapidly varying tides. Of course, a similar loss of accuracy may be encountered in shore areas where the available empirical data are marginal in quantity and/or quality. This applies particularly to the Arctic and Antarctic coastlines, where large ice sheets may also cause some tidal distortions and retardations which have been assumed as negligible.

(f) **Interactions of Oceanic and Terrestrial Tides:** All NSWC partial tide models include effects of tide-generated terrestrial and oceanic mass perturbations in the simplified forms described in Section 1. In agreement with other investigations, the inclusion of the earth tides produced significant first-order effects on the tidal amplitudes. As expected useful second-order corrections of amplitudes and phases were registered when oceanic load effects were introduced, which is also in agreement with other numerical experiments conducted for instance by Zahel (1977), Accad and Pekeris (1978), Estes (1979), and Parke and Hendershott (1979).

(g) **Hydrodynamical Properties:** The ocean tidal equations are the basic LTEs augmented by physically meaningful lateral eddy dissipation and linear bottom friction terms with novel eddy and friction coefficients depending linearly on the respective vertical and horizontal cell wall areas. Accordingly, eddy dissipation was found significant in deep oceans while bottom friction dominated in shallow waters. It may be emphasized that both terms were needed to achieve realistic decay and dispersion features and to bring the computed data in good agreement with empirical data. They were not necessary to enforce stability of the numerical procedure. In particular, the friction coefficients were meshwise modified by the unique hydrodynamical interpolation in order to achieve a realistically smooth acceptance of over 2000 empirical tide data pairs.

(h) **Mathematical Features:** The finite difference analogue of the ocean tidal equations uses the Hansen-Zahel (Zahel 1970) staggered difference scheme in space and improved differences in time. At boundaries, the free-slip condition is strictly imposed while the no-crossflow condition is acceptably violated by the hydrodynamical interpolation of empirical boundary data. To avoid the polar singularity of the tidal equations, a new series expansion is incorporated, which will be described in a forthcoming paper (Schwiderski 1981i).

3. APPLICATIONS OF THE NSWC OCEAN TIDE DATA

Various applications of the NSWC ocean tide models require indispensable consideration of the major features summarized in Section 2. This is particularly true in applications to compute interactions of oceanic and terrestrial tides, which manifest themselves as anomalies in the earth's gravity field. Though such perturbations amount to only a few microgals, they can now be determined by precision measurements such as those carried out by Kuo et al. (1970) and more recently on a global scale by Melchior et al. (1980). On the other hand, gravity anomalies caused by ocean tidal loading can be computed from accurate ocean tide (e.g., Schwiderski 1980b and 1981j) and solid earth (e.g., Farrell 1972) models (see, for instance, Goad 1980a, b and Melchior et al. 1980). Unfortunately, both the experimentally derived and computed gravity perturbations of ocean tidal loads are subject to the following difficulties.

(a) **Empirical Gravity Anomalies of Ocean Tidal Loads:** The measured gravity anomalies include dominant perturbations caused by the direct terrestrial tides, which are usually modeled by neglecting all lateral density and elasticity variations of the solid earth. Yet, those heterogeneities exist and may not be negligible especially in tectonically active areas. Furthermore, after subtraction of the modeled terrestrial tide effects, the remaining gravity anomalies must be subjected to the same harmonic analysis as the measured ocean tide, which is particularly controversial for short-time observations.

(b) **Computed Gravity Anomalies of Ocean Tidal Loads:** The computation of the yielding of the solid earth under the oceanic tide is also based on a highly simplified lateral homogeneous earth model as the terrestrial tide model (a) above. Moreover, as has been pointed out by various investigators (e.g., Pekeris 1978, Accad and Pekeris 1978, Goad 1980a, b, and Schwiderski 1980a), the purely numerical computation of gravity anomalies from ocean tidal loads poses considerable difficulties and may be the cause of significant errors. This is particularly true at coastal and oceanic stations where the convolution of the numerical ocean tide with a singular Green's function must be computed. Indeed, depending on the actual numerical procedure chosen, the singularity may amplify in various ways any or all of the deficiencies of the NSWC ocean tide models discussed in Section 2.

In spite of all the possible error sources, Goad (1980a, b) and Melchior et al. (1980) arrive at excellent agreements for the bulk of their compared empirical and computed gravity anomalies. These results lend convincing support to the accuracy claim of Melchior et al. concerning their worldwide gravity measurements. Vice versa, these investigations verify indirectly the oceanwide accuracy and usefulness of the NSWC tide models. In fact, a detailed analysis of the massive Melchior et al. data indicates that all of the remaining discrepancies between the empirical and computed data can probably be traced to one or more of the modeling and computing difficulties elaborated above.

(c) **Immediate Improvements:** In coastal and oceanic areas some of the deficiencies of the NSW tide models (Section 2, b, c, e) can be immediately removed by preparing more detailed limited-area tide models on a locally refined grid system. Models of this sort have already been constructed by various researchers especially for excluded or insufficiently resolved bordering waters. Where no refined models are available, the refinement may be accomplished by linear or higher-order interpolation of the listed $1^\circ \times 1^\circ$ data and by using additional empirical data, say, from the huge collections of the British Admiralty (1977) and the International Hydrographic Bureau (1978), which are in possession of the author. In such a refining process the hydrodynamically interpolated empirical tide data, which have been displaced to the centers of the grid cells (Section 2, c), should be first returned as closely as possible to the original tide gauge locations. Specifically, in boundary cells the empirical data should be moved from the cell centers to the centers of the boundary segments (Figure 1a). For small oceanic (unresolved) islands, the tabulated empirical data should be moved from the cell centers to the original stations (Figure 1b). This relatively simple local procedure is strongly recommended where the ocean tides suffer drastic distortions and retardations from the deep oceans to the shallow areas.




Figure 1. Replacing of Empirical Tide Data

The important effect of the replacement of empirical data may be illustrated by the incredibly sharp decay of the tidal ranges (= double amplitudes) across the continental shelf of the New York Bight. As can be seen in Figure 2, the tidal ranges measured by Gill and Porter (1980) drop from 4.6 ft at the coast over a shelf width of less than 200 km to an almost constant height of about 3 ft in the deep ocean. The global NSW ocean tide model approximates this distortion very closely, but only when the empirical boundary datum $\bar{x}$ is returned to the original boundary position x . Obviously, simple linear interpolations between the adjusted model data x and x' could significantly improve computations of corresponding gravity anomalies in that neighborhood.




Figure 2. Sharp Decay of Tidal Ranges Across the Continental Shelf of New York Bight: $\odot$ Gill-Porter Measurements, $(\bar{x}, x')$ NSWC Tide Model, x Replaced Empirical Boundary Datum $\bar{x}$

(d) **Future Improvements:** Refined local tide models are presently under consideration by the author. The results of those models will subsequently be used in connection with forthcoming new and better empirical data to improve the global NSWC tide models. An error reduction from 10 cm to 3 cm is already being called for in various applications. Of course, a model of this accuracy will require more and better measured and analyzed empirical data especially in critical areas.

4. NSWG GLOBAL OCEAN TIDE DATA (GOTD) TAPE

The NSWG GOTD-1981 tape contains the amplitudes $\xi_{m,n}^i$ (in meters) and Greenwich phases $\delta_{m,n}^i$ (in degrees, $0^\circ = 360^\circ$) of the leading harmonic ocean tidal modes (M_2 , S_2 , K_1 , O_1 , N_2 , P_1 , K_2 , Q_1 , Mf , Mm , Ssa) on a $1^\circ \times 1^\circ$ spherical grid system. This GOTD-1981 tape is identical to the earlier GOTD-1980 tape, which did not contain the Mm and Ssa data.

The entire data set ($\xi_{m,n}^i$, $\delta_{m,n}^i$) is arranged by modes according to the mode number $i = 1, 2, 3, \dots, 11$ as defined in Table 2, which also lists the corresponding species number ν (compare also Table 1).

Table 2. Entire Data Set (see Table 1)

Tidal Symbol	M_2	S_2	K_1	O_1	N_2	P_1	K_2	Q_1	Mf	Mm	Ssa
Mode Number: i	1	2	3	4	5	6	7	8	9	10	11
Species Number: ν	2	2	1	1	2	1	2	1	0	0	0

In each mode ($i = 1, 2, \dots, 11$), the data are arranged by colatitude numbers $n = 1, 2, \dots, 168$ (see Equation 2) in consecutive pairs of blocks each of which contain 361 words of amplitudes or phases arranged by longitude numbers $m = 1, 2, \dots, 360$ (see Equation 1) with the last (361-st) word indicating the colatitude number n . The first block in each pair contains the amplitudes $\xi_{m,n}^i$ in Format F5.3 and the second one the phases $\delta_{m,n}^i$ in Format F5.1. The data have been blockwise generated by the BUFFER-OUT statement on the CDC 6700 computer. The magnetic tape has the following properties: seven track, BCD form, even parity, 556 bpi, and unlabeled.

The tidal constants ($\xi_{m,n}^i$, $\delta_{m,n}^i$) are representative for the $1^\circ \times 1^\circ$ square with the geographical center point (λ_m, θ_n) , where

$$\lambda_m = (m - 0.5)^\circ = \text{longitude east } (m = 1, 2, \dots, 360) \quad (1)$$

$$\theta_n = (n - 0.5)^\circ = \text{colatitude } (n = 1, 2, \dots, 168) \quad (2)$$

On land (continental or large islands; see Schwiderski 1978c) all tidal constants have been set to:

$$\xi_{m,n}^i = 9.999, \quad \delta_{m,n}^i = 999.9 \text{ for land} \quad (3)$$

The precise definition of those data, their accuracy and limitation, and their proper application and improvement has been discussed in Sections 2 and 3. To avoid possible misleading conclusions any application of those data *must* evaluate realistically the listed features and suggestions.

The instantaneous ocean tide ζ_i (in meters) of any model i ($= 1, 2, \dots, 11$) can be found at any given instant (y = year, d = day number of year y , and t = universal time of day d in seconds) and any given ocean point (λ, θ) by linearly interpolating the amplitude ξ_i and phase δ_i belonging to the point (λ, θ) in the corresponding partial tide tables and computing

$$\zeta_i = \zeta_i(\lambda, \theta, t) = \xi_i \cos [\sigma_i t + \pi (\chi_i - \delta_i)/180] \quad (4)$$

In this formula σ_i and χ_i denote, respectively, the frequency and the astronomical argument of the partial tide listed in Table 1.

The corresponding instantaneous astronomical partial equilibrium tide η_i (or tide-generating potential G_i ; see Schwiderski 1978a, 1979c, d, or 1980a) is determined by

$$\eta_i = \eta_i(\lambda, \theta, t) = K_i f_{\nu_i}(\theta) \cos [\sigma_i t + \pi (\chi_i + \nu_i \lambda)/180] \quad (5)$$

where $K_i = K$ and $\nu_i = \nu$ denote, respectively, the tidal amplitude and species number listed in Table 1. The species function of colatitude θ is defined by

$$f_{\nu}(\theta) = \begin{cases} \sin^2 \theta & \text{for } \nu = 2 \\ \sin 2\theta & \text{for } \nu = 1 \\ \frac{1}{2}(3 \sin^2 \theta - 2) & \text{for } \nu = 0 \end{cases} \quad (6)$$

In a simple second-order Love-number approximation, the corresponding solid-earth tide ζ_i^e is given by (see Schwiderski 1978a, 1979c, d, or 1980a)

$$\zeta_i^e = \zeta_i^e(\lambda, \theta, t) \approx 0.612 \eta_i \quad (7)$$

Similarly, in a simple Accad-Pekeris approximation (see Schwiderski 1978a, 1979c, d, or 1980a), the corresponding dip (yielding) of the solid earth $\zeta_i^{e,o}$ in response to the oceanic tidal load ζ_i is given by

$$\zeta_i^{e,o} = \zeta_i^{e,o}(\lambda, \theta, t) \approx -0.0667 \zeta_i \quad (8)$$

Thus, the corresponding instantaneous geocentric partial tide ζ_i^c (relative to the geoidal sea surface at rest) is determined by linear superposition to

$$\zeta_i^c = \zeta_i^c(\lambda, \theta, t) = \zeta_i + \zeta_i^e + \zeta_i^{e,o} \quad (9)$$

Finally, by linear superposition of all corresponding partial tides, one finds the instantaneous local tides:

$$\xi = \xi(\lambda, \theta, t) = \sum_i \xi_i = \text{total oceanic tide} \quad (10a)$$

$$\xi^c = \xi^c(\lambda, \theta, t) = \sum_i \xi_i^c = \text{total geocentric tide} \quad (10b)$$

Note to Users: Copies of the NSWC GOTD tape in its discribed standard format (or any other desired format) may be requested from the authors:

E. W. Schwiderski, Code K104, Tel. (703) 663-8406

L. T. Szeto, Code K14, Tel. (703) 663-7741

NSWC, Dahlgren, Virginia 22448

5. NSWC RANDOM-POINT TIDE PROGRAM (RPTIDE)

The following NSWC RPTIDE computes from the harmonic tidal constants listed on the NSWC GOTD 1981 (or GOTD 1980) tape (Section 4) the instantaneous oceanic and/or geocentric tides at given random points and instances. The corresponding Users' Guide and Program Listing are given in Appendixes A and B. A fast program to compute geocentric tides along satellite tracks is described in Schwiderski, E. W. and Szeto, L. T. 1981.

1) Input Data

- (1) $(\xi_{m,n}^i, \delta_{m,n}^i)$ = ocean tide amplitudes (in m) and Greenwich phases (in deg) from GOTD 1981 (or 1980) tape, where

$i = 1, 2, 3, \dots, 11$ = mode numbers for GOTD 1981
 $(0 < i < 10$ for GOTD 1980)

$m = 1, 2, \dots, 360$ ($\neq 0$) = longitude numbers

$n = 1, 2, \dots, 168$ = colatitude numbers

Note: $\xi_{m,n}^i = 9.999$, $\delta_{m,n}^i = 999.9$ on land

$\delta_{m,n}^i = 360^\circ = 0^\circ$ (phase jump)

- (2) y = year ≥ 1975 (fixed for one run!)

- (3) d = day of year y ($d = 1$ for January 1st, also fixed for one run!)

- (4) t_j = specified instances (in sec after Greenwich midnight of day d , where $j = 1, 2, 3, \dots, J$)

- (5) (λ_j, θ_j) = longitudes (East) and latitudes (+ North, - South) in deg. of random points corresponding to instances t_j

- (6) J = number of random points

- (7) I = number of superposed tidal modes if $1 \leq I \leq 11$ with option
 $-1 \leq I \leq -11$ for computation of single tidal mode $i = |I|$, $|I| < 10$
for GOTD 1980 tape)

- (8) $\alpha = 0$ or $= 1$ = earth tide parameter ($\alpha = 1$ earth tide included, $\alpha = 0$ earth tide excluded)

(9) $\beta = 0$ or $= 1$ = ocean load parameter ($\beta = 1$ ocean load included, $\beta = 0$ ocean load excluded)

(10) $\gamma = 0$ or $= 1$ = ocean tide parameter ($\gamma = 1$ ocean tide included, $\gamma = 0$ ocean tide excluded)

II) Preliminary Computations

With the tidal parameters $\nu = \nu_i$, $\sigma = \sigma_i$, $K = K_i$, h_o , s_o , p_o specified in Tables 1 and 2 compute:

$$h = \pi h_o / 180, \quad s = \pi s_o / 180, \quad p = \pi p_o / 180$$

$$\chi = \chi_i \text{ for } i = 1, 2, 3, \dots, 11$$

and set ($E_i = 0.612 K_i$)

$$\begin{array}{lll} E_1 = 0.148\,308, & E_2 = 0.069\,059, & E_3 = 0.086\,638, \\ E_4 = 0.061\,515, & E_5 = 0.028\,396, & E_6 = 0.028\,668, \\ E_7 = 0.018\,791, & E_8 = 0.011\,785, & E_9 = 0.025\,546, \\ E_{10} = 0.013\,480, & E_9 = 0.011\,901, & \end{array}$$

also

$$\bar{\gamma} = \gamma - 0.0667 \beta$$

III) Main Computation

(1) Set $j = 1$

(2) Set $\xi_j = 0$ and

$$i \left\{ \begin{array}{l} = 1 \text{ for } I > 0, \\ = |I| \text{ for } I \geq 0 \text{ (single mode!)} \end{array} \right.$$

(3) Compute:

$$m = \text{Int} [\lambda_j + 0.5] + 1, \psi = m - (\lambda_j + 0.5)$$

$$n = \text{Int} [90.5 - \phi_j] + 1, \theta = \bar{\theta} = n - (90.5 - \phi_j)$$

and transfer to core memory

$$\begin{array}{l} (\xi_{m-1, n-1}^i, \delta_{m-1, n-1}^i); (\xi_{m, n-1}^i, \delta_{m, n-1}^i) \\ (\xi_{m-1, n}^i, \delta_{m-1, n}^i); (\xi_{m, n}^i, \delta_{m, n}^i) \end{array} \quad (*)$$

where $m - 1 = 0 \rightarrow 360$

Check for land points and replace

- (a) if $\xi_{m, n-1}^i = 9.999$, replace $\theta \rightarrow 0$
- (b) if $\xi_{m, n}^i = 9.999$, replace $\theta \rightarrow 1$
- (c) if (a) and (b) hold, replace $\psi \rightarrow 1$
- (d) if $\xi_{m-1, n-1}^i = 9.999$, replace $\bar{\theta} \rightarrow 0$
- (e) if $\xi_{m-1, n}^i = 9.999$, replace $\bar{\theta} \rightarrow 1$
- (f) if (d) and (e) hold, replace $\psi \rightarrow 0$
- (4) Interpolate ξ_j^i or the ruled second-order surface in ψ and θ :

$$\begin{aligned} \xi_j^i = & (1 - \psi) [\theta \xi_{m, n-1}^i + (1 - \theta) \xi_{m, n}^i] \\ & + \psi [\bar{\theta} \xi_{m-1, n-1}^i + (1 - \bar{\theta}) \xi_{m-1, n}^i] \end{aligned} \quad (**)$$

Test for 360° phase jumps and replace

$$\begin{aligned} \text{if } \delta_{m, n-1}^i - \delta_{m, n}^i & \begin{cases} > 180, \text{ replace } \delta_{m, n-1}^i \rightarrow \delta_{m, n-1}^i - 360 \\ < -180, \text{ replace } \delta_{m, n}^i \rightarrow \delta_{m, n}^i - 360 \end{cases} \\ \text{if } \delta_{m-1, n-1}^i - \delta_{m, n}^i & \begin{cases} > 180, \text{ replace } \delta_{m-1, n-1}^i \rightarrow \delta_{m-1, n-1}^i - 360 \\ < -180, \text{ replace } \delta_{m, n}^i \rightarrow \delta_{m, n}^i - 360 \end{cases} \\ \text{if } \delta_{m-1, n}^i - \delta_{m, n}^i & \begin{cases} > 180, \text{ replace } \delta_{m-1, n}^i \rightarrow \delta_{m-1, n}^i - 360 \\ < -180, \text{ replace } \delta_{m, n}^i \rightarrow \delta_{m, n}^i - 360 \end{cases} \\ \text{if } \delta_{m-1, n-1}^i - \delta_{m, n-1}^i & \begin{cases} > 180, \text{ replace } \delta_{m-1, n-1}^i \rightarrow \delta_{m-1, n-1}^i - 360 \\ < -180, \text{ replace } \delta_{m, n-1}^i \rightarrow \delta_{m, n-1}^i - 360 \end{cases} \\ \text{if } \delta_{m-1, n}^i - \delta_{m, n-1}^i & \begin{cases} > 180, \text{ replace } \delta_{m-1, n}^i \rightarrow \delta_{m-1, n}^i - 360 \\ < -180, \text{ replace } \delta_{m, n-1}^i \rightarrow \delta_{m, n-1}^i - 360 \end{cases} \end{aligned}$$

$$\text{if } \delta_{m-1, n}^i - \delta_{m-1, n-1}^i \begin{cases} > 180, \text{ replace } \delta_{m-1, n}^i \rightarrow \delta_{m-1, n}^i - 360 \\ < -180, \text{ replace } \delta_{m-1, n-1}^i \rightarrow \delta_{m-1, n-1}^i - 360 \end{cases}$$

Use adjusted δ s to interpolate δ_j^i by formula (**) with ξ replaced by δ .

Now, compute and replace

$$\begin{aligned} \zeta_j &\rightarrow \zeta_j + \bar{\gamma} \xi_j^i \cos [\sigma_i t_j + \chi_i - \pi \delta_j^i / 180] \\ &+ \alpha E_i(\phi_j) \cos [\sigma_i t_j + \chi_i + \pi \nu_i \lambda_j / 180] \end{aligned}$$

where

$$E_i(\phi_j) = \begin{cases} E_i \cos^2 \phi_j & \text{for } \nu_i = 2 \\ E_i \sin^2 \phi_j & \text{for } \nu_i = 1 \\ \frac{1}{2} E_i (3 \cos^2 \phi_j - 2) & \text{for } \nu_i = 0 \end{cases}$$

If $i < |I|$, replace $i \rightarrow i + 1$, transfer to core memory new data (*), and repeat routine (4).

If $i \leq |I|$, print λ_j , ϕ_j , t_j , and ζ_j .

If $j < J$, replace $j \rightarrow j + 1$ and repeat (2) through (4).

If $j \leq J$ stop program.

Note: This program does not consider any special features or suggestions made in Sections 2 and 3. Hence, if a higher accuracy near coastal boundaries is needed, an appropriate local computation should be used following the suggestions of Section 2.

REFERENCES

1. Accad, Y. and Pekeris, C. L., 1978. "Solution of the Tidal Equations for the M_2 and S_2 Tides in the World Oceans from a Knowledge of the Tidal Potential Alone," *Phil. Trans. Roy. Soc., London, Ser. A*, 290, p. 235.
2. British Admiralty Tide Tables, 1977. Vols. 1, 2, and 3.
3. Estes, R. H., 1979. "A Simulation of Global Ocean Tide Recovery Using Altimeter Data with Systematic Orbit Error," *Marine Geodesy*, 3, p. 75.
4. Farrell, W. E., 1972. "Deformation of the Earth by Surface Loads," *Rev. Geophys. Space Phys.*, 10, p. 261.
5. Gill, S. K. and Porter, D. L., 1980. "Theoretical Offshore Tide Range Derived from a Simple Defant Tidal Model Compared with Observed Offshore Tides," *International Hydrographic Review, Monaco* LVII, p. 155.
6. Goad, C. C., 1980a. "Gravimetric Tidal Loading Computed from Integrated Green's Functions," *J. Geophysical Research*, 85, B5, p. 2679.
7. Goad, C. C., 1980b. "The Computation of Tidal Loading Effects with Integrated Green's Functions," Proceedings of the Second International Symposium on Problems Related to the Redefinition of North American Vertical Geodetic Networks, May 26-30, Ottawa, Canada.
8. Grace, S. F., 1930. "The Semidiurnal Lunar Tidal Motion of the Red Sea," *Monthly Notices of the Roy. Astr. Soc., Geophysical Supplement*, II, p. 273.
9. Hendershott, M. C., 1972. "The Effects of Solid-Earth Deformation on Global Ocean Tides," *Geophys. J. Roy. Astr. Soc.*, 29 p. 380.
10. International Hydrographic Bureau, 1978. *Harmonic Constants*, Computer Tape, Monaco.
11. Kuo, J. T., Jachens, R. C., White, G., and Ewing M., 1970. "Tidal Gravity Measurements Along a Transcontinental Profile Across the United States," *Comm. Observ. Roy. Belg.*, 9, p. 96.
12. Melchior, P., Moens, M., Ducarme, B., and VanRuymbeke, M., 1980. "Tidal Loading Along a Profile Europe-East Africa-South Asia-Australia and the Pacific Ocean," *Physics of the Earth and Planetary Interiors*.

13. Parke, M. E. and Hendershott, M. C., 1979. " M_2 , S_2 , K_1 Models of the Global Ocean Tide on an Elastic Earth," *Marine Geodesy* 3, p. 379.
14. Pekeris, C. L., 1978. "The Bodily Tide and the Yielding of the Earth Due to Tidal Loading," *Geophys. J. Roy. Astr. Soc.*, 52, p. 471.
15. Proudman, J., 1928. "Deformation of Earth-Tides by Means of Water-Tides in Narrow Seas," *Bull. No. 11, Sect. Oceanogr., Cons. de Recherches, Venedig.*
16. Schwiderski, E. W., 1978a. *Global Ocean Tides, Part I: A Detailed Hydrodynamical Interpolation Model*, NSWC/DL TR-3866.
17. Schwiderski, E. W., 1978b. "A Detailed Hydrodynamical Interpolation Model of Worldwide Ocean Tides," presented at the Int. Symp. on Interaction of Marine Geodesy and Ocean Dynamics, Miami, Florida, October 10-15.
18. Schwiderski, E. W., 1978c. *Hydrodynamically Defined Ocean Bathymetry*, NSWC/DL TR-3888.
19. Schwiderski, E. W., 1979a. "NSWC Ocean Tide Program," presented at the NASA SEASAT ALT/POD Calibration Workshop, Austin, Texas, June 11-15.
20. Schwiderski, E. W., 1979b. "Detailed Ocean Tide Models of (N_2 , M_2 , S_2 , K_2) and (K_1 , P_1 , O_1 , Q_1) Including an Atlas of Tidal Charts and Maps," presented at the XVIIth General Assembly of the International Union of Geodesy and Geophysics in Canberra, Australia, December 2-15.
21. Schwiderski, E. W., 1979c. "Ocean Tides, Part I: Global Tidal Equations," *Marine Geodesy*, 3, p. 161.
22. Schwiderski, E. W., 1979d. "Ocean Tides, Part II: A Hydrodynamical Interpolation Model," *Marine Geodesy*, 3, p. 219.
23. Schwiderski, E. W., 1979e. *Global Ocean Tides, Part II: The Semidiurnal Principal Lunar Tide (M_2), Atlas of Tidal Charts and Maps*, NSWC TR 79-414.
24. Schwiderski, E. W., 1980a. "On Charting Global Ocean Tides," *Reviews of Geophysics and Space Physics*, 18, p. 243.
25. Schwiderski, E. W., 1980b. "NSWC Ocean Tide Data Tape, GOTD 1980," Available at the Naval Surface Weapons Center Dahlgren, Virginia 22448.
26. Schwiderski, E. W., 1981a. *Global Ocean Tides, Part III: The Semidiurnal Principal Solar Tide (S_2), Atlas of Tidal Charts and Maps*, NSWC TR 81-122.

27. Schwiderski, E. W., 1981b. *Global Ocean Tides, Part IV: The Diurnal Luni-Solar Declination Tide (K_1)*, *Atlas of Tidal Charts and Maps*, NSWC TR 81-142.
28. Schwiderski, E. W., 1981c. *Global Ocean Tides, Part V: The Diurnal Principal Lunar Tide (O_1)*, *Atlas of Tidal Charts and Maps*, NSWC TR 81-144.
29. Schwiderski, E. W., 1981d. *Global Ocean Tides, Part VI: The Semidiurnal Elliptical Lunar Tide (N_2)*, *Atlas of Tidal Charts and Maps*, NSWC TR-218.
30. Schwiderski, E. W., 1981e. *Global Ocean Tides, Part VII: The Diurnal Principal Solar Tide (P_1)*, *Atlas of Tidal Charts and Maps*, NSWC TR 81-220.
31. Schwiderski, E. W., 1981f. *Global Ocean Tides, Part VIII: The Semidiurnal Luni-Solar Declination Tide (K_2)*, *Atlas of Tidal Charts and Maps*, NSWC TR 81-222.
32. Schwiderski, E. W., 1981g. *Global Ocean Tides, Part IX: The Diurnal Elliptical Lunar Tide (Q_1)*, *Atlas of Tidal Charts and Maps*, NSWC TR 81-224.
33. Schwiderski, E. W., 1981h. *Similarities of the Leading Semidiurnal (N_2 , M_2 , S_2 , K_2) and Diurnal (K_1 , P_1 , O_1 , Q_1) Ocean Tides*, Naval Surface Weapons Center Technical Report in preparation.
34. Schwiderski, E. W., 1981i. *Exact Expansions of Arctic Ocean Tides*, Naval Surface Weapons Center Technical Report in preparation.
35. Schwiderski, E. W., 1981j. "NSWC Ocean Tide Data Tape, GOTD 1981," Available at the Naval Surface Weapons Center, Dahlgren, Virginia 22448.
36. Schwiderski, E. W., 1982a. *Global Ocean Tides, Part X: The Fortnightly Lunar Tide (M_f)*, *Atlas of Tidal Charts and Maps*, Naval Surface Weapons Center Technical Report in preparation.
37. Schwiderski, E. W., 1982b. *Global Ocean Tides, Part XI: The Monthly Lunar Tide (M_m)*, *Atlas of Tidal Charts and Maps*, Naval Surface Weapons Center Technical Report in preparation.
38. Schwiderski, E. W., 1982c. *Global Ocean Tides, Part XII: The Semiannual Solar Tide (S_{sa})*, *Atlas of Tidal Charts and Maps*, Naval Surface Weapons Center Technical Report in preparation.
39. Schwiderski, E. W., 1982d. *Similarities of the Leading Long-Period (M_f , M_m , S_{sa}) Ocean Tides*, Naval Surface Weapons Center Technical Report in preparation.

40. Schwiderski, E. W. and Szeto L. T., 1981. *On Computing Instantaneous Geocentric Tides Along Satellite Tracks, the NSWC STT Program*, Naval Surface Weapons Center, Technical Report in preparation.
41. Takahasi, R., 1929. "Tilting Motion of the Earth Crust Caused by Tidal Loading," *Bull. Earthquake Res. Inst.*, 6, p. 85.
42. Wahr, J. M. and Sasao, T., 1980. "A Diurnal Resonance in the Ocean Tide and in the Earth's Load Response Due to the Resonant Free Core Nutation," *Geophys. J. Roy. Astr. Soc.*
43. Zahel, W., 1970. "Die Reproduktion Gezeitenbedingter Bewegungsvorgänge im Weltozean Mittels des *Hydrodynamisch-Numerischen Verfahrens*," *Mitteilungen des Inst. f. Meereskunde der Univ. Hamburg*, XVII.
44. Zahel, W., 1977. "The Influence of Solid Earth Deformations on Semidiurnal and Diurnal Oceanic Tides," *Proc. IRLA Int. Collog. on Numerical Methods of Science and Technical Computation*, Springer, Berlin.

APPENDIX A

RPTIDE PROGRAM USER'S GUIDE

APPENDIX A

RPTIDE PROGRAM USER'S GUIDE

The Random-Point Tide (RPTIDE) program computes instantaneous oceanic and/or geocentric tides at randomly given geographical points and specified instances as described in Section 5. The program listing in Appendix B shows RPTIDE written in FORTRAN IV extended for the CDC 6700 computer under the SCOPE 3.4 operating system. The BUFFER IN statement buffers in GOTD 1980 or 1981 tide constants (Section 4), which are in binary mode. The GOTD magnetic tape is in coded mode; its files are attached as follows for the example $I = 6$, $L = IT1$ (see Sections 4 and 5):

ATTACH. TAPE L	$\xi_{m,n}^1$	} GOTD files
ATTACH. TAPE L + 1,	$\delta_{m,n}^1$	
ATTACH. TAPE L + 2,	$\xi_{m,n}^2$	
ATTACH. TAPE L + 3,	$\delta_{m,n}^2$	
⋮		
ATTACH. TAPE L + 10,	$\xi_{m,n}^6$	
ATTACH. TAPE L + 11,	$\delta_{m,n}^6$	

All other input data are entered using data cards consisting of one type A data card one or more type B data cards. The type A card contains 8 right-justified integers with the format of 8I5. The integers represent the following ordered variables (Section 5 symbols in parentheses):

YEAR = (y) = year > 1975 (fixed for one run!)

DAY = (d) = day of YEAR (d = 1 for January 1st, also fixed for one run!)

JPTS = (J) = total number of random points or instances with same YEAR and DAY

MTIDE = (I) = total number of tidal modes to be superposed ($1 \leq i \leq I$ where $I < 10$ for GOTD 1980 and $I < 12$ for GOTD 1981). Option if MTIDE = - N with $0 < N < 10$ or 12, the computed tide is the single partial tide $i = N$

ETINC = (α) = earth tide parameter (ETINC = 0 excludes and ETINC = 1 includes earth tide)

LOAD = (β) = ocean load parameter (LOAD = 0 excludes and LOAD = 1 includes ocean load)

OCEAN = (γ) = ocean tide parameter (OCEAN = 0 excludes and OCEAN = 1 includes ocean tide)

IT1 = beginning unit number of $2 * \lfloor \text{MTIDE} \rfloor$ consecutive unit numbers to which GOTD files are attached (see example above)

Each type B data card contains three numbers with the format 3F20.8. These numbers are, respectively:

TIME = (t) = time (in sec) of given instant after Greenwich midnight of day DAY (may be several days later)

LONG = (λ) = Longitude (East; in degrees)

LAT = (ϕ) = Latitude (+ North, - South; in degrees)

Other major notations corresponding to Section 5 are:

D = (δ) = Oceanic tidal phase

X = (ξ) = Oceanic tidal amplitude

T = (ζ) = instantaneous tidal height

NU = (ν) = tidal species number

FREQ = (σ) = tidal frequency

ASTRO = (χ) = astronomical argument

ETIDE = (E) = earth tide constant amplitude

ET = (E(ϕ)) = earth tide geographical amplitude

Note: For land points in squares with 9.999 vertex amplitudes only the earth tide is meaningful if ETINC = 1, LOAD = OCEAN = 0. For improved accuracy in coastal waters, special computations should incorporate the suggestions of Section 3 c and d.

APPENDIX B
RPTIDE PROGRAM LISTING

08/27/81 14.17.22

FTN 4.6+433

SUBROUTINE COMPUT 74774 OPT=1

```

1      SUBROUTINE COMPUT
      *      I M, X, D, XN, DN, LONG
      *      NU, TIME, ASTRO, NAL, ETINC, GAMMAR
      *      Y, OTT, OLL, ETT
      *      P, PPG, LAT, ETIDE, T)
      *      DIMENSION ASTRO(11), X(360), D(360), XN(360), DN(360)
      *      FREZ(11), ETIDE(11), NU(11)
      *      INTEGER ETINC
      *      REAL LONG, LAT
      *      DATA CTR/Q.01745329252/
      *      XLAND=9.999
      *      M1=M-1
      *      IF(M.EQ.1) M1=360

15      C      COSECTION TC P-4- OF FORMULATIONS(11-1-80)
      C      P= M-(LONG+.5)
      C      TR= N*1.0-( 90.5-LAT)
      C      TE=TR
      C      IF( XN( M).EQ.XLAND) TE=0
      C      IF( X( M).EQ.XLAND) TE=1
      C      IF( XN(M1).EQ.XLAND) TB=0
      C      IF( X(M1).EQ.XLAND) TB=1
      C      IF( XN( M).EQ.XLAND.AND. X(M).EQ.XLAND) P=1

25      C      IF( XN(M1).EQ.XLAND.AND. X(M1).EQ.XLAND) P=0
      C      INTERPOLATE
      *      XLJ=(1-P)*(XN(M)*TE+ (1-TE)*X(M))
      *      + P*( XN(M1)*TB + (1-TB)*X(M1))
      *      DMN1= DN(M)
      *      DMN= 0(M)
      *      DM1N1= DN(M1)
      *      DM1N= D(M1)
      *      D1= DN(M)- D(M)
      *      D2= DN(M1)- D(M)
      *      D3= D(M1)- D(M)
      *      D4= DN(M1)- DN(M)
      *      D5= D(M1)- DN(M)
      *      D6= D(M1)- D(M1)
      *      IF(D1.GT. 180.0) DMN1 =DMN1 -360.0
      *      IF(D1.LT.-180.0) DMN =DMN -360.0
      *      IF(D2.GT. 180.0) DM1N1=DM1N1-360.0
      *      IF(D2.LT.-180.0) DMN =DMN -360.0
      *      IF(D3.GT. 180.0) DM1N =DM1N -360.0
      *      IF(D3.LT.-180.0) DMN =DMN -360.0
      *      IF(D4.GT. 180.0) DM1N1=DM1N1-360.0
      *      IF(D4.LT.-180.0) DMN1 =DMN1 -360.0
      *      IF(D5.GT. 180.0) DM1N =DM1N -360.0
      *      IF(D5.LT.-180.0) DM1N =DM1N -360.0
      *      IF(D6.GT. 180.0) DM1N =DM1N -360.0
      *      IF(D6.LT.-180.0) DM1N1=DM1N1-360.0
      *      INTERPOLATE
      *      DLJ=(1-P)*( DMN1*TE+ (1-TE)*DMN)
      *      + P*( DM1N1*TB+ (1-TB)*DM1N)
      *      COMPUTE EARTH TIDE
      *      COSLAT= COS(LAT*CTR)
      *      COS2= COS(LAT*CTR)
      *      ET=COS2

```


08/27/81 14.17.22

FTN 4.6+433

SUBROUTINE PRINTC 74/74 OPT=1

```

1      SUBROUTINE PRINTC
      ( T, LONG, LAT, J, TJ, TIME
      , OT, OL, ETD
      , MTIDE)
      REAL LONG, LAT
      IF(.NOT.(J.EQ.1)) GO TO 200
      PRINT10C
      FORMAT(*1*, I20,*(DEGREES) * , T42, *TOTAL*
      , T82, *TOTAL TIDE*
      , T96, *OCEAN TIDE OCEAN LOADING EARTH TIDE*
      )
      PRINT104
      FORMAT(* , T17, *LONG*, T34, *LAT*
      , T42, *COMPONENT TIDES*, T63, *TIME*
      , T85, *( METERS*, T126, *)*
      )
      CONTINUE
      ENDIF
      PRINT300, LONG, LAT, MTIDE, TIME, TJ, OT, OL, ETD
      FORMAT(*0*, 2F20.6, 15, F20.2, 9X, 4F14.4)
      RETURN
      ENDC
200    C
      100  *
      *
      *
      104  *
      *
      *
      15   *
      *
      *
      200  C
      300
  
```

SYMBOLIC REFERENCE MAP (R=3)

ENTRY POINTS	DEF LINE	REFERENCES
3 PRINTC	1	21

VARIABLES	SN	TYPE	RELOCATION	REFS	DEFINED	DEFINED
0 ETD		REAL	F.P.	19	DEFINED	1
0 J		INTEGER	F.P.	6	DEFINED	1
0 LAT		REAL	F.P.	5	DEFINED	1
0 LONG		REAL	F.P.	5	DEFINED	1
0 MTIDE		INTEGER	F.P.	19	DEFINED	1
0 OL		REAL	F.P.	19	DEFINED	1
0 OT		REAL	F.P.	19	DEFINED	1
0 T		REAL	F.P.	1	DEFINED	1
0 TIME		REAL	F.P.	19	DEFINED	1
0 TJ		REAL	F.P.	19	DEFINED	1
*UNUSED						

FILE NAMES	MODE	WRITES	7	12	19
CUTPLT	FMT				

STATEMENT LABELS	DEF LINE	REFERENCES
21 100 FMT	8	7
40 104 FMT	13	12
13 200 FMT	17	6
66 300 FMT	20	19

STATISTICS	PROGRAM LENGTH
	730 50

DISTRIBUTION

Library of Congress
Washington, DC 20540
ATTN: Gift and Exchange Division

Defense Technical Information Center
Cameron Station
Alexandria, VA 22314

Director
Defense Mapping Agency, HQ
Washington, DC 20360
ATTN: Dr. C. F. Martin
P. W. Schwimmer

Defense Mapping Agency
Hydrographic-Topographic Center
6500 Brooks Lane
Washington, DC 20315
ATTN: MAJ J. Jerome
Randy Smith
Mrs. I. Fischer

Defense Mapping Agency
Aerospace Center
St. Louis, MO 63118
ATTN: William P. Wall
Don McEntee

Office of Naval Research
800 N. Quincy St.
Arlington, VA 22203
ATTN: G. R. Hamilton
Dr. W. S. Wilson
J. G. Heacock
R. S. Andrews

Oceanographer of the Navy
U. S. Naval Observatory
34 and Massachusetts Ave. NW
Washington, DC 20390

Naval Oceanographic Office
NSTL Station
Bay St. Louis, MS 39522
ATTN: Dr. T. Davis
L. B. Bourquin

(12) Technical Director (SP-20)
Strategic Systems Project Office
Washington, DC 20390

Naval Postgraduate School
Monterey, CA 93940
ATTN: Prof. R. L. Haney
Prof. M. L. Elsberry
Prof. D. C. Gallacher
Prof. R. W. Garwood, Jr.
Prof. N. K. Mooers

Fleet Numerical Oceanography Center
Monterey, CA 93940
ATTN: CDR B. Schramm

Director
Naval Research Laboratory
Washington, DC 20360
ATTN: V. E. Noble
B. S. Yapple
A. Shapiro
D. T. Chen

Army Engineers Topographic Laboratory
Ft. Belvoir, VA 22060
ATTN: Dr. A. Mancini

Air Force Geodetic Laboratory
L. G. Hanscom Field
Bedford, MA 01730

DISTRIBUTION (Continued)

Scripps Institution of Oceanography
University of California at San Diego
LaJolla, CA 92037

ATTN: Dr. W. H. Munk
Dr. M. C. Hendershott
Prof. B. D. Zetler
Prof. S. M. Smith
Prof. H. W. Menard
Dr. J. H. Filloux
Dr. B. Bernstein
Dr. D. S. Luther

Dr. C. Wunsch
MIT/Dept. Earth & Planetary Sciences
Cambridge, MA 02139

Woods Hole Oceanographic Institute
Woods Hole, MA 02543

ATTN: Dr. H. M. Stommel
Dr. G. Veronis
Dr. N. P. Fofonoff
Dr. J. Whitehead
Dr. P. G. Brewer

Battelle Columbus Laboratories
505 King Ave.
Columbus, OH 43201
ATTN: A. G. Mourad

Dr. J. W. Chamberlain
Rice University
Houston, TX 77001

Dr. R. H. Rapp
Ohio State University
Dept. of Geodetic Science
1958 Neil Ave.
Columbus, OH 43210

Dr. R. O. Reid
Texas A&M University
College Station, TX 77843

Florida State University
Dept. of Oceanography
Tallahassee, FL 32306
ATTN: Dr. J. J. O'Brien
Dr. W. Sturges
Ms. L. Vasant

Prof. F. E. Snodgrass
Inst. of Geophysics and Planetary Physics
University of California at San Diego
LaJolla, CA 92037

Prof. K. Wyrtki
University of Hawaii
Honolulu, HI 96822

Prof. B. Tapley
Dept. of Aerospace Eng. & Eng. Mechanics
WRW 402
University of Texas
Austin, TX 78712

Prof. D. Lynch
Thayer School of Engineering
Dartmouth College
Hanover, NH 03755

Prof. T. B. Curtin
North Carolina State Univ. at Raleigh
Dept. of Marine Science & Engineering
P.O. Box 5923
Raleigh, NC 27650

University of Washington
Dept. of Oceanography WB-10
Seattle, WA 98195
ATTN: Dr. A. J. Clarke
Prof. D. Winter
Prof. M. Jamart

DISTRIBUTION (Continued)

David T. Haislip
U. S. Coast Guard
400 7th Street, S.W.
Washington, DC 20590

NOAA/National Ocean Survey
National Geodetic Survey
Rockville, MD 20852
ATTN: Dr. B. Chovitz
Dr. J. M. Diamante
Dr. B. C. Douglas
Dr. C. C. Goad
Dr. F. Morrison

NOAA/National Ocean Survey
Oceanographic Division
Rockville, MD 20852
ATTN: D. C. Simpson
D. L. Porter
R. A. Smith
B. B. Parker

NOAA/Atlantic Oceanographic and
Meteorological Lab.
Physical Oceanography Laboratory
15 Rickenbacker Causeway
Miami, FL 33149
ATTN: G. A. Maul
H. M. Byrne

NOAA/Pacific Marine Environmental Lab.
Seattle, WA 98105
ATTN: Dr. J. R. Apel
H. O. Mofjeld
C. A. Pearson
M. Byrne

NOAA/Geophysical Fluid Dynamics Lab.
Princeton University
Princeton, NJ 08540
ATTN: Dr. J. Smagorinsky
Dr. K. Bryan
Dr. M. D. Cox

NOAA/National Center for Atmospheric
Research
Boulder, CO 80303
ATTN: Dr. W. R. Holland

NASA/Goddard Space Flight Center
Greenbelt, MD 20771
ATTN: Dr. J. W. Siry
D. E. Smith
J. G. Marsh
T. L. Felsentreger
J. Zwally

NASA/Wallops Station
Information Processing and Analysis
Branch
Wallops Island, VA 23337
ATTN: C. D. Leitao
N. E. Huang
W. B. Krabill
B. Speidel

Director
U.S. Army Ballistic Research Laboratory
Aberdeen Proving Ground, MD 21005
ATTN: DRDAR-TBS-S (STINFO)

Smithsonian Astrophysical Observatory
60 Garden St.
Cambridge, MA 02138
ATTN: Dr. E. M. Gaposchkin
Dr. G. C. Wiffenbach
B. Stevens

National Science Foundation
1951 Constitution Ave., N.W.
Washington, DC 20550
ATTN: Mathematical Sciences Division
J. G. Gross
R. E. Wall

DISTRIBUTION (Continued)

Jet Propulsion Laboratory
California Institute of Technology
4800 Oak Grove Drive
Pasadena, CA 91103
ATTN: Dr. M. Parke
Dr. J. Lorell
Dr. G. Born

Prof. J. T. Kuo
Lamont-Doherty Geological Observatory
Columbia University
Palisades, NY 10964

Prof. W. J. Pierson, Jr.
1641 Rosalind Ave.
Elmont, NY 11003

Prof. Morris Schulkin
Applied Physics Laboratory
University of Washington
Seattle, WA 98105

Dr. R. H. Estes
Business and Technological Systems, Inc.
Aerospace Building, Suite 605
10210 Greenbelt Rd.
Seabrook, MD 20801

T. V. Martin
Sci. Res. and Appl. Group
Washington Analytical Services Center, Inc.
6801 Kenilworth Ave.
Riverdale, MD 20840

Dr. S. K. Jordan
The Analytic Sciences Corporation
6 Jacob Way
Reading, MA 01867

The Rand Corporation
Santa Monica, CA 90406
ATTN: Director, Climate Program

Local:

C
D
E31 (GIDEP)
E41
K
K02
K04
K05
K10
K102
K1040
K104U
K104S (400)
K104Z
K12
K12G
K12M
K12S
K12T
K12 W
K13
K13H
K14
K14S
K20
K204
K21
K21D
K30
K40
K404A
K404S
K41
K42
K44
K50
K51B
R

DISTRIBUTION (Continued)

R02

R04

R31G

R40

R44

R44VT

X210 (6)