


α

AD-A107992

LEVEL III




395-568

20 OCT 1977

R. M. Bell

MSNW



AD A107992

KINETIC STUDIES FOR XeF AND KrF LASERS

DTIC
ELECTE
NOV 30 1981
H

Prepared for
OFFICE OF NAVAL RESEARCH

September 1977

DTIC FILE COPY

DISTRIBUTION STATEMENT
Approved for public release
Distribution Unlimited



MATHEMATICAL SCIENCES NORTHWEST, Inc.

10 10 19

11 14 September 1977

14 Program Code Number TE20
MSNW-1062

9 Quarterly Research and development rept. no. 3

1

A042910

QUARTERLY R&D STATUS REPORT NO. 3

6 TITLE: KINETIC STUDIES FOR XeF AND KrF LASERS

15 Contract Number: N00014-76-C-1066
Effective Date: 1 September 1976
Expiration Date: 30 November 1977
Amount: \$99,909

DTIC ELECTED
NOV 30 1981
H

Sponsored by
Defense Advanced Research Projects Agency
DARPA Order 1806

10 Principal Investigator: R. E. Center (206) 827-0460
C. H. Fisher (206) 827-0460

Scientific Officer: Director, Physics Programs
Physical Sciences Division
Office of Naval Research
Department of the Navy
800 North Quincy Street
Arlington, Virginia 22117

MATHEMATICAL SCIENCES NORTHWEST, INC.
P. O. Box 1887
Bellevue, Washington 98009

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U. S. Government.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

404058

B

TABLE OF CONTENTS

| SECTION | PAGE |
|---|------|
| I INTRODUCTION | 1 |
| II EXPERIMENTAL RESULTS | 2 |
| 2.1. XeF(B ^{1/2}) Radiative Lifetime | 2 |
| 2.2. XeF(B ^{1/2}) Collisional Quenching | 4 |
| REFERENCES | 9 |

| | |
|--------------------|-------------------------------------|
| Accession For | |
| NTIS GRA&I | <input checked="" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By | |
| Distribution/ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A | |

LIST OF FIGURES

| FIGURE | | PAGE |
|--------|--|------|
| 1 | The 353 nm Fluorescence from a Gas Mixture of He + 10 Torr Xe + 0.7 Torr UF ₆ at a Total Pressure of 125 Torr | 3 |
| 2 | Decay Frequency as a Function of Pressure for a He + 4% Xe + 0.3% UF ₆ Gas Mixture | 5 |
| 3 | Variation of 353 nm Fluorescence Decay as a Function of Fluorine Partial Pressure | 6 |

LIST OF TABLES

| TABLE | | PAGE |
|-------|--|------|
| 1 | Summary of XeF(B $\frac{1}{2}$) Radiative Lifetime and Quenching Data | 7 |

I. INTRODUCTION

yield

The goal of this program is to determine the radiative lifetime and the collisional quenching rate coefficients for the XeF and KrF upper laser levels using a state selective, laser excited fluorescence technique. Details of the experimental procedure were described in our semi-annual report.¹ During this reporting period, we have determined the XeF(B^{1/2}) radiative lifetime by monitoring the decay of the 353 nm XeF(B^{1/2} → X) fluorescence as a function of pressure. Preliminary collisional quenching rate constants have been determined for He, Xe, and F₂ from the variation of the fluorescence decay time constant as a function of partial pressure of the collision partner. Collisional quenching by Ar was investigated; however, no bimolecular rate constant was determined because of nonexponential decay of the XeF(B → X) emission in the presence of Ar. This behavior suggests collisional coupling to some other excited state, most likely the XeF(C^{3/2}) state.

Bimolecular collisional quenching rate coefficients by Ne and NF₃¹ remain to be determined during the next reporting period. In addition, termolecular quenching by He, Ne, and Ar will be investigated by increasing the partial pressures of these constituents to the 1 to 3 atm range. Collisional coupling between the B^{1/2} and C^{3/2} states will also be investigated by looking for fluorescence at 460 nm on the XeF(C → X) band.

- | -

II. EXPERIMENTAL RESULTS

2.1. XeF(B $\frac{1}{2}$) Radiative Lifetime


Oscilloscope traces showing the 351 nm exciting pulse and the 353 nm fluorescence signal are presented in Figure 1. This signal was obtained with a He + 10 torr Xe + 0.7 torr UF₆ gas mixture at 125 torr total pressure. We have switched to UF₆ as the F atom source because its UV absorption cross section (resulting in dissociation to produce an F atom) is two orders of magnitude larger than that of F₂, which yields a substantial increase in the XeF fluorescence. However, the UF₅ produced during the flash has a low vapor pressure resulting in the formation of solid particles which increased the scattered light from the 351 nm exciting pulse on subsequent shots. Therefore, it was necessary to pump out the cell and refill it with fresh gas after every shot.

If the XeF(B $\frac{1}{2}$) excited state is lost only by radiative decay and bimolecular collisions, it will decay exponentially once the exciting laser pulse has terminated. A semilogarithmic plot of the 353 nm fluorescence decay read from the oscilloscope trace is presented in Figure 1. The slope of the straight line fit to the data yields the decay time constant τ^{-1} given by

$$\begin{aligned}\tau^{-1} &= \tau_B^{-1} + k_{\text{He}} [\text{He}] + k_{\text{Xe}} [\text{Xe}] + k_{\text{UF}_6} [\text{UF}_6] \\ &= \tau_B^{-1} + (k_{\text{He}} X_{\text{He}} + k_{\text{Xe}} X_{\text{Xe}} + k_{\text{UF}_6} X_{\text{UF}_6}) p\end{aligned}$$


where τ_B is the radiative lifetime of the XeF(B $\frac{1}{2}$) state, k_i and X_i are the quenching rate coefficient and the mole fraction of the i^{th} component of the mixture, and p is the total pressure. Thus, a plot of the decay time constant τ^{-1} as a function of pressure for a

a)



PHOTODIODE (XeF Laser)
PHOTOMULTIPLIER (353 nm
Fluorescence)

b)



77 01035


Figure 1. The 353 nm Fluorescence from a Gas Mixture of He+10 Torr Xe + 0.7 torr UF₆ at a Total Pressure of 125 torr. a) Oscilloscope traces showing the 351 nm exciting laser pulse and the 353 nm fluorescence signal. b) Semi-logarithmic plot of the 353 nm fluorescence decay.

gas mixture with fixed composition should follow a straight line with the $\text{XeF}(B \frac{1}{2})$ radiative lifetime as the zero pressure intercept. A plot of the data obtained for a He + 4% Xe + 0.13% UF_6 gas mixture is shown in Figure 2. The zero pressure intercept of the straight line fit to the data gives a $\text{XeF}(B \frac{1}{2})$ radiative lifetime of 18.7 ± 1.4 nsec. This result is in good agreement with that obtained by Burnham and Harris from the dissociative excitation of XeF_2 .²

2.2. $\text{XeF}(B \frac{1}{2})$ Collisional Quenching


Bimolecular quenching rate constants for various molecules are obtained by measuring the $\text{XeF}(B \frac{1}{2})$ decay time constant as a function of the partial pressure of that constituent while holding the partial pressures of the other constituents fixed. The slope of the straight line fit to the data gives the two-body quenching rate constant for that molecule. To illustrate the procedure, a plot of the 353 nm decay time constant τ^{-1} as a function of F_2 partial pressure is shown in Figure 3. The partial pressures of the other constituents of the gas mixtures were held fixed at 250 torr He + 10 torr Xe + 0.3 torr UF_6 . The quenching rate constant for F_2 determined from the slope of the straight line fit to the data is $k_{\text{F}_2} = 1.2 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$. Using this same procedure, quenching rate constants for He and Xe have also been determined. These rate constants are listed in Table 1 along with the pressure range over which the quenching was studied. The measured rate constants are in fairly good agreement with those inferred by Brashears, Setser and Desmarreau from Stern-Volmer plots of the quenching of $\text{XeF}(B \frac{1}{2} \rightarrow X)$ emission produced by photolysis of XeF_2 .³

No quenching rate constant is listed in the table for Ar even though it is a primary constituent of e-beam excited XeF laser



77 01036

Figure 2. Decay Frequency as a Function of Pressure for a He + 4% Xe + 0.13% UF₆ Gas Mixture. The zero pressure intercept of the straight line fit to the data gives a XeF (B_{1/2}) radiative lifetime of 18.7 nsec.



77 01037

Figure 3. Variation of 353 nm Fluorescence Decay as a Function of Fluorine Partial Pressure. The slope of the straight line fit to the data gives a quenching rate constant of $1.2 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$.

Table 1
Summary of XeF (B $\frac{1}{2}$) Radiative Lifetime
and Quenching Data

XeF(B $\frac{1}{2}$) $\tau_r = 18.7 \pm 1.4$ nsec
XeF(B $\frac{1}{2}$) 2-Body Quenching Rate Constants

| Molecule | $k_q(\text{cm}^3 \text{ Molecule}^{-1} \text{ sec}^{-1})$ | Pressure Range (torr) |
|----------------|---|--------------------------|
| He | 2×10^{-12} | 75 - 750 |
| Xe | 6×10^{-11} | 5 - 80 |
| F ₂ | 1.2×10^{-10} | 0.5 - 24 |

mixtures. This is due to the fact that the 353 nm fluorescence in the presence of Ar exhibited nonexponential decay over much of the pressure range studied, making it difficult to extract a rate constant without a more complicated analysis. This behavior suggests collisional transfer from the $B^{1/2}$ state during the decay period. The most likely candidate is the $XeF(C^{3/2})$ state which is nearly degenerate with the $B^{1/2}$ state.⁴ During the next reporting period, the possibility of collisional coupling between the $B^{1/2}$ and $C^{3/2}$ states will be investigated by looking for broadband fluorescence at 460 nm on the $XeF(C^{3/2} \rightarrow X)$ transition.

REFERENCES

1. R.E. Center and C.H. Fisher, "Kinetic Studies for XeF and KrF Lasers," MSNW Semi-Annual Report, May 1977.
2. R. Burnham and N.W. Harris, *J. Chem. Phys.* 66, 2742 (1977).
3. H.C. Brashears, Jr., D.W. Setser, and D. Desmarteau, *Chem. Phys. Lett.* 48, 84 (1977).
4. T.H. Dunning and P.J. Hay, to be published.